

Hermite Expansions of C -regularized cosine Functions

Rachid Ameziane Hassani ¹

Aziz Blali,²

Youssef Bajjou³

and Abdelkhalek El Amrani,^{4,5}

Abstract

The aim of this paper is to approximate the exponentially bounded C -regularized cosine function by the Hermite series, recalling the notions and the results used.

AMS Mathematics Subject Classification (2010): Primary 47D60; Secondary 34G10, 47D03, 33C45.

Key words and phrases: Hermite functions, Hermite Expansions, C -regularized semigroup, C -regularized cosine functions.

1 Introduction and preliminaries

The series expansion of Hermite orthogonal polynomials have been an important tool in quantum mechanics and statistical studies, both theoretical and applied. The study of sufficient conditions for the convergence of Hermite series has been the subject of numerous works ; for more details see [9]. In 2015, Luciano and Pedro studied in their article [4], the Hermite expansion of C_0 -groups and cosine functions, in this works we will be interested in Hermite expansion of C -regularized cosine function, starting with reminding the notations, concepts and results used.

Throughout this paper E denotes a non-trivial complex Banach space, $\mathfrak{F}(E, F)$ denotes the set of all applications from E to another Banach space F , $B(E)$ denotes the space of bounded linear operators from E into itself, and $L_{loc}^1(E)$ the set of all $f \in \mathfrak{L}(\mathbb{R}, E)$ locally integrable. For a closed linear operator A on E , $\mathcal{D}(A)$, $R(A)$ and $\rho(A)$ denote its domain, range and resolvent set, respectively. $\mathcal{D}(A)$ equipped with the graph norm $\|x\|_{\mathcal{D}(A)} = \|x\|_E + \|Ax\|_E$ become Banach space . Throughout this paper, $C \in B(E)$ will be an injective operator. The C -resolvent set of A , denoted by $\rho_C(A)$, is defined by

¹Sidi Mohamed Ben Abdellah University, Dhar El mahraz Faculty of Sciences, Department of Mathematics, Atlas Fès, Morocco.
e-mail: ra.ameziane@yahoo.fr

²Sidi Mohamed Ben Abdellah University, Ecole Normale Supérieure, Department of Mathematics, B. P. 5206 Bensouda-Fès, Morocco.
e-mail: aziz.blali@usmba.ac.ma

³Sidi Mohamed Ben Abdellah University, Dhar El Mahraz Faculty of Sciences , Department of mathematics and computer science, Atlas Fès, Morocco.
e-mail: youssefbajjou2017@gmail.com

⁴Sidi Mohamed Ben Abdellah University , Dhar El Mahraz Faculty of Sciences, Department of mathematics and computer science, Atlas Fès, Morocco.
e-mail: abdelkhalek.elamrani@usmba.ac.ma

⁵Corresponding author

$\rho_C(A) := \{\lambda \in \mathbb{C} \mid R(C) \subseteq R(\lambda I - A) \text{ and } \lambda I - A \text{ is injective in } B(E)\}$ and by $R_C(\lambda, A) = (\lambda I - A)^{-1}C$ ($\lambda \in \rho_C(A)$) the C -resolvent.

Hermite functions and Hermite Expansions on Banach spaces

For all $n \in \mathbb{N}$, the classical Hermite polynomial, defined by Rodrigues formula:

$$(\forall x \in \mathbb{R}) \quad H_n(x) = e^{x^2} (-1)^n \frac{d^n}{dx^n} e^{-x^2}.$$

H_n is a polynomial with the of degree n , the same parity as n , whose highest monomial degree is $2^n X^n$ and have real coefficients. Furthermore, they verify the following condition of orthogonality:

$$\int_{\mathbb{R}} H_n(x) H_m(x) e^{-x^2} dx = \delta_{n,m} n! 2^n \sqrt{\pi},$$

where $\delta_{n,m}$ is the Kronecker delta. We also have recurrence relations, differential equations and satisfied the Muckenhoupt estimates:

$$(\exists c > 0) (\forall (n_0, t) \in \mathbb{N} \times \mathbb{R} : t^2 \leq 2(2n_0 + 1)) (\forall n \geq n_0) \mid H_n(t) \mid \leq c \left(\frac{e^{\frac{t^2}{2}} \sqrt{2^n n!}}{n^{\frac{1}{12}}} \right). \quad (1)$$

For more details of the classical theory of orthogonal polynomials see [21], [13], [3], [4] and [2].

The Hermite functions on \mathbb{R} are defined by:

$$\phi_n(x) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} H_n(x) e^{-\frac{x^2}{2}} = (-1)^n \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} e^{\frac{x^2}{2}} \frac{d^n}{dx^n} e^{-x^2}. \quad (2)$$

$(\phi_n)_{n \in \mathbb{N}}$ is an orthonormal basis in the Hilbert space $L^2(\mathbb{R})$ and satisfied some recurrence relations, equality and inequality. For more details See [21], [13], [3], [4] and [2].

For $n \in \mathbb{N}$, we denote by φ_n , the function on \mathbb{R} defined by :

$$(\forall t \in \mathbb{R}) \quad \varphi_n(t) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} e^{-\frac{t^2}{2}} \phi_n(t) = \frac{1}{2^n n! \sqrt{\pi}} e^{-t^2} H_n(t) = \frac{(-1)^n}{2^n n! \sqrt{\pi}} \frac{d^n}{dt^n} e^{-t^2}. \quad (3)$$

φ_n has the same parity as n , satisfies recurrence relations and differential equations, for example:

$$(\forall (n, m) \in \mathbb{N}^2) \quad \varphi_n^{(m)} = (-1)^m 2^m (n+1) \dots (n+m) \varphi_{n+m}. \quad (4)$$

And the following useful inequality given by Cauchy-Schwartz inequality:

$$\| \varphi_n \|_1 \leq \frac{1}{\sqrt{2^n n!}}. \quad (5)$$

For more details see [4].

Of the most important properties of $(\varphi_n)_{n \in \mathbb{N}}$ family is that if $f : \mathbb{R} \rightarrow E$ be a differentiable function such that $\int_{-\infty}^{+\infty} e^{-t^2} \|f(t)\|^2 dt < +\infty$, then the series $\sum_{n \in \mathbb{N}} c_n(f)H_n(t)$ converges pointwise to $f(t)$ for each $t \in \mathbb{R}$, where

$$c_n(f) = \int_{-\infty}^{+\infty} \varphi_n(t)f(t)dt.$$

For more details see [13], [19], [4] and [9].

C -regularized semigroup and C -regularized cosine functions

C -regularized semigroup

A map $T : [0, +\infty[\rightarrow B(E)$ is called C -regularized semigroup or C -semigroup, if

1. $T(t+s)C = T(t)T(s)$ for all $t, s \in \mathbb{R}^+$.
2. $T(0) = C$.
3. $t \mapsto T(t)x$ is continuous on \mathbb{R}^+ for every $x \in E$.

Its generator W is defined by

$$\mathcal{D}(W) = \left\{ x \in E : \lim_{s \rightarrow 0^+} \frac{T(s)x - Cx}{s} \text{ exists in } R(C) \right\}$$

and

$$(\forall x \in \mathcal{D}(W)) \quad Wx = C^{-1} \lim_{s \rightarrow 0} \frac{T(s)x - Cx}{s}.$$

We say that $(T(t))_{t \geq 0}$ is exponentially bounded C -regularized semigroup if $(\exists M \geq 0) (\exists \omega \geq 0) : \|T(t)\| \leq Me^{\omega t}$, for all $t \geq 0$.

$(T(t))_{t \geq 0}$ is no degenerate if $T(t)x = 0$ for all $t \geq 0$, then $x = 0$. see [6], [18] and [17].

Since we have for all (t, x) fixed in $\mathbb{R}^+ \times \mathcal{D}(W)$:

$$\lim_{s \rightarrow 0^+} \frac{T(s)T(t)x - CT(t)x}{s} = T(t) \lim_{s \rightarrow 0^+} \frac{T(s)x - Cx}{s} = T(t)CWx = CT(t)Wx \in R(C),$$

this means that $T(t)x \in \mathcal{D}(W)$ and $WT(t)x = T(t)Wx$.

The following results can be found in [6] and [5].

Let $(T(t))_{t \in \mathbb{R}^+}$ be a strongly continuous family such that $\|T(t)\| \leq Me^{\omega t}$ for all $t \geq 0$ and W is closed linear operator, to conclude that $(T(t))_{t \in \mathbb{R}^+}$ is C -regularized semigroup generated by W , it is sufficient that :

$$W = C^{-1}WC, (\omega, +\infty) \subset \rho_C(W), \text{ and } R_C(\lambda, W)x = \int_0^{+\infty} e^{-\lambda t} T(t)x dt \text{ for } \lambda > \omega, x \in E. \tag{6}$$

We refer to [25], [5] and [18] for more details.

Let $0 < \alpha \leq \frac{\pi}{2}$, $\Sigma_\alpha = \{\lambda \in \mathbb{C}/\lambda \neq 0 \text{ and } |\arg(\lambda)| < \alpha\}$ and let $(T(t))_{t \geq 0}$ be a C -regularized semigroup. Then we say that $(T(t))_{t \geq 0}$ is an analytic C -regularized semigroup of angle α , if there exists an analytic function $\mathbf{T} : \Sigma_\alpha \rightarrow B(E)$ which satisfies:

1. $\mathbf{T}(t) = T(t)$ for all $t > 0$.
2. $\lim_{z \rightarrow 0, z \in \Sigma_\gamma} \mathbf{T}(z)x = 0$ for all $\gamma \in]0, \alpha[$ and $x \in E$.

For more details see [25].

C -regularized cosine function

A map $T : \mathbb{R} \rightarrow B(E)$ is called C -regularized cosine function or C -cosine function, if

1. $T(t+s)C + T(t-s)C = 2T(t)T(s)$ for all $t, s \in \mathbb{R}$.
2. $T(0) = C$.
3. $t \rightarrow T(t)x$ is continuous on \mathbb{R} for every $x \in E$.

We say that $(T(t))_{t \in \mathbb{R}}$ is exponentially bounded C -regularized cosine if

$$(\exists M > 0) (\exists \omega \geq 0) : \|T(t)\| \leq Me^{\omega|t|}, \text{ for all } t \in \mathbb{R}.$$

The associated sine operator function $S(\cdot)$ is defined by $S(t) := \int_0^t T(s)ds$ for all $t \in \mathbb{R}$. The operator W defined by

$$\mathcal{D}(W) = \left\{ x \in E : \lim_{s \rightarrow 0} \frac{2(T(s)x - Cx)}{s^2} \text{ exists in } R(C) \right\}$$

and

$$Wx = C^{-1} \lim_{s \rightarrow 0} \frac{2(T(s)x - Cx)}{s^2} \text{ for all } x \in \mathcal{D}(W).$$

is called the generator of $(T(t))_{t \in \mathbb{R}}$. See [12] and [17].

We notice that W is closed linear operator in E , T is odd, $T(t)x \in \mathcal{D}(W)$, $S(t)x \in \mathcal{D}(W)$, $\int_0^t S(s)x ds \in \mathcal{D}(W)$ and $W \int_0^t S(s)x ds = T(t)x - Cx$ for all $x \in E$ and $t \in \mathbb{R}$, $C^{-1}WC = W$ and if $(T(t))_{t \in \mathbb{R}}$ is exponentially bounded C -regularized cosine then the function sine $S := S(t)_{t \in \mathbb{R}}$ is also exponentially bounded, so for all $k \in \mathbb{N}$ and $x \in \mathcal{D}(W)$,

$$\lim_{t \rightarrow \pm\infty} t^k e^{-t^2} T(t)x = \lim_{t \rightarrow \pm\infty} t^k e^{-t^2} S(t)x = \lim_{t \rightarrow \pm\infty} t^k e^{-t^2} WS(t)x = 0, \quad (7)$$

because, if $\|T(t)\| \leq Me^{\omega t}$ for all $t \in \mathbb{R}$, then for all $x \in \mathcal{D}(W)$:

$$\|t^k e^{-t^2} T(t)x\| \leq |t|^k e^{-|t|^2} \|T(t)x\| \leq |t|^k e^{\frac{\omega^2}{4}} e^{-(|t| - \frac{\omega}{2})^2} \|x\|. \text{ See [12,}$$

proposition 1.1 and 1.2] for more properties and more details.
Consider in E the well-posed Cauchy problem

$$((ACP(W, u_0, u_1, 0)_2) \left\{ \begin{array}{l} u''(t) = Wu(t), \quad t \in \mathbb{R} \\ u(0) = u_0 \\ u'(0) = u_1. \end{array} \right.$$

Where W generates a C -cosine operator function $(T(t))_{t \in \mathbb{R}}$ then

$$u(t) = C^{-1}T(t)u_0 + C^{-1}S(t)u_1$$

is the unique solution of the above Cauchy problem for every pair (u_0, u_1) of initial values in $C(\mathcal{D}(A))$. For more details see [12].

A strongly continuous family $(T(t))_{t \in \mathbb{R}^+}$ such that $\|T(t)\| \leq Me^{\omega t}$, for all $t \geq 0$ is C -regularized cosine with generator the closed linear operator W if and only if the following condition holds:

$$C^{-1}WC = W, \lambda^2 \in \rho_C(W) \text{ and } \lambda R_C(\lambda^2, W) = \int_0^{+\infty} e^{-\lambda t} T(t) dt \text{ on } E \text{ for all } \lambda > \omega.$$

For more details see [11], [12] and [23].

If W is the generator of exponentially bounded and C -regularized cosine $(T(t))_{t \in \mathbb{R}}$, then W is the generator of analytic C -regularized semigroup of angle $\frac{\pi}{2}$ defined by

$$(\forall z \in \Sigma_{\frac{\pi}{2}}, T_1(z)x = \int_0^{+\infty} \frac{e^{-\frac{s^2}{4z}}}{\sqrt{\pi z}} T(s)x ds \text{ (for all } z \in \Sigma_{\frac{\pi}{2}}, \operatorname{Re}(z) > 0) \quad (8)$$

whose proof is similar for C_0 -cosine operator-valued function. (See [10] and [1]).

2 Main results

Lemma 2.1. *Let $(T(t))_{t \in \mathbb{R}}$ be an exponentially bounded C -cosine function on a Banach space E with generator $(W, \mathcal{D}(W))$. For all $n \in \mathbb{N}$ and $x \in \mathcal{D}(W)$, we have :*

1. $\int_{-\infty}^{+\infty} \varphi_{2n+1}(t)T(t)x dt = 0$
2. $\int_{-\infty}^{+\infty} \varphi_{2n}(t)T(t)x dt = \frac{1}{2^{2n}(2n)!} W^n T_1(\frac{1}{4})x.$
3. *In the case where $\sup_{t \in \mathbb{R}} \|T(t)\| < +\infty$, we have :*
 - (i) $\|W^n T_1(\frac{1}{4})x\| \leq 2^n \sqrt{(2n)!} \sup_{t \in \mathbb{R}} \|T(t)\| \|x\|.$
 - (ii) $\|\int_{-\infty}^{+\infty} \varphi_{2n}(t)T(t)x dt\| \leq \frac{c_1 \sqrt{(2(n-1))!}}{2^{n+1} (2n)!} \|Wx\|$ with c_1 a positive constant.

Proof

1. Let $n \in \mathbb{N}$ and $x \in E$. The function $t \mapsto \varphi_{2n+1}(t)T(t)x$ is continuous and odd, integrable in the sens of Bochner, then $\int_{-\infty}^{+\infty} \varphi_{2n+1}(t)T(t)xdt = 0$.
2. Let $x \in \mathcal{D}(W)$. For all $n \in \mathbb{N}$, let $I_n(x) = \int_{-\infty}^{+\infty} \frac{d^{2n}}{dt^{2n}}(e^{-t^2})T(t)xdt$, then:

$$\begin{aligned}
 I_n(x) &= \left[\frac{d^{2n-1}}{dt^{2n-1}}(e^{-t^2})T(t)x \right]_{-\infty}^{+\infty} - \int_{-\infty}^{+\infty} \frac{d^{2n-1}}{dt^{2n-1}}(e^{-t^2}) \frac{d}{dt}T(t)xdt \\
 &= 0 - \left[\frac{d^{2n-2}}{dt^{2n-2}}(e^{-t^2})WS(t)x \right]_{-\infty}^{+\infty} + \int_{-\infty}^{+\infty} \frac{d^{2n-2}}{dt^{2n-2}}(e^{-t^2}) \frac{d^2}{dt^2}T(t)xdt \\
 &= 0 - 0 + \int_{-\infty}^{+\infty} \frac{d^{2n-2}}{dt^{2n-2}}(e^{-t^2})WT(t)xdt \\
 &= W \int_{-\infty}^{+\infty} \frac{d^{2n-2}}{dt^{2n-2}}(e^{-t^2})T(t)xdt \\
 &= W I_{n-1}(x).
 \end{aligned}$$

A simple recurrence on n gives

$$I_n(x) = W^n I_0(x) = W^n \int_{-\infty}^{+\infty} e^{-t^2}T(t)xdt$$

From where by definition of φ_n we obtain:

$$\begin{aligned}
 \int_{-\infty}^{+\infty} \varphi_{2n}(t)T(t)xdt &= \int_{-\infty}^{+\infty} \frac{(-1)^{2n}}{2^{2n}(2n)!\sqrt{\pi}} \frac{d^{2n}}{dt^{2n}}e^{-t^2}T(t)xdt \\
 &= \frac{1}{2^{2n}(2n)!\sqrt{\pi}} I_n(x) \\
 &= \frac{W^n}{2^{2n}(2n)!\sqrt{\pi}} \int_{-\infty}^{+\infty} e^{-t^2}T(t)xdt \\
 &= \frac{W^n}{2^{2n}(2n)!\sqrt{\pi}} 2 \int_0^{+\infty} e^{-t^2}T(t)xdt \\
 &= \frac{W^n}{2^{2n}(2n)!} T_1\left(\frac{1}{4}\right)x.
 \end{aligned}$$

3. In the case where $\sup_{t \in \mathbb{R}} \|T(t)\| < +\infty$, we obtain:

- (i) For all $x \in \mathcal{D}(W)$:

$$\begin{aligned}
 \| W^n T_1\left(\frac{1}{4}\right)x \| &\leq 2^{2n}(2n)! \int_{-\infty}^{+\infty} \| \varphi_{2n}(t)T(t)x \| dt \\
 &\leq 2^{2n}(2n)! \left(\sup_{t \in \mathbb{R}} \| T(t) \| \right) (\| x \|)(\| \varphi_{2n}(t) \|_1) \\
 &\leq 2^{2n}(2n)! \sup_{t \in \mathbb{R}} \| T(t) \| \times \| x \| \times \frac{1}{\sqrt{2^{2n}(2n)!}} \quad (\text{by 5}) \\
 &\leq 2^n \sqrt{(2n)!} \| x \| \sup_{t \in \mathbb{R}} \| T(t) \| .
 \end{aligned}$$

- (ii) Let $x \in \mathcal{D}(W)$ and $n \geq 1$, if we posed $B = \| \int_{-\infty}^{+\infty} \varphi_{2n}(t)T(t)x dt \|$ then using (i) and integration by parts we get

$$\begin{aligned}
 B &= \| \frac{1}{4(2n-1)(2n)} \int_{-\infty}^{+\infty} \varphi''_{2n-2}(t)T(t)x dt \| \quad (\text{by 4}) \\
 &= \| -\frac{1}{4(2n-1)(2n)} \int_{-\infty}^{+\infty} \varphi'_{2n-2}(t)T'(t)x dt \| \\
 &= \| \frac{1}{4(2n-1)(2n)} \int_{-\infty}^{+\infty} \varphi_{2n-2}(t)T''(t)x dt \| \\
 &= \| \frac{1}{4(2n-1)(2n)} \int_{-\infty}^{+\infty} \varphi_{2n-2}(t)WT(t)x dt \| \quad (\text{like 1}) \\
 &\leq \frac{1}{4(2n-1)(2n)} \left(\int_{-\infty}^{+\infty} | \varphi_{2n-2}(t) | dt \right) \left(\sup_{t \in \mathbb{R}} \| T(t) \| \right) (\| Wx \|) \\
 &\leq \left(\sup_{t \in \mathbb{R}} \| T(t) \| \right) \frac{1}{4(2n-1)(2n)} \| \varphi_{2n-2} \|_1 \| Wx \| \\
 &\leq \frac{c_1}{2^2(2n-1)(2n)} \frac{1}{2^{n-1}\sqrt{2(n-1)!}} \| Wx \| \quad (\text{by 5}) \quad (c_1 = \sup_{t \in \mathbb{R}} \| T(t) \|) \\
 &\leq \frac{c_1 \sqrt{2(n-1)!}}{(2n-1)(2n)} \frac{1}{2^{n+1}(2(n-1)!)} \| Wx \| \\
 &\leq \frac{c_1 \sqrt{2(n-1)!}}{2^{n+1}(2n)!} \| Wx \| .
 \end{aligned}$$

Theorem 2.1. Let $(T(t))_{t \in \mathbb{R}}$ be an exponentially bounded C -cosine function on a Banach space E ($T(t) \leq Me^{\omega|t|}$, $M > 0$ and $\omega \geq 0$) with a generator $(W, \mathcal{D}(W))$.

1. For any $x \in \mathcal{D}(W)$,

$$T(t)x = \sum_{n=0}^{+\infty} \frac{1}{2^{2n}(2n)!} W^n T_1\left(\frac{1}{4}\right)x H_{2n}(t), \quad \text{for all } t \in \mathbb{R}.$$

2. In the case where $\sup_{t \in \mathbb{R}} \| T(t) \| < +\infty$, we have for any $x \in \mathcal{D}(W)$:

- (i)

$$T_1(z)x = \sum_{n=0}^{+\infty} \frac{1}{2^{2n}n!} W^n T_1\left(\frac{1}{4}\right)x (4z-1)^n, \text{ for all } z \in \mathbb{C}, \left|z - \frac{1}{4}\right| < \frac{1}{4}.$$

- (ii) For all $t \in \mathbb{R}$, there is $m_t \in \mathbb{N}$ and $c_t > 0$ such that:

$$(\forall m \geq m_t) \quad \left\| T(t)x - \sum_{n=0}^m \frac{1}{2^{2n}(2n)!} W^n T_1\left(\frac{1}{4}\right)x H_{2n}(t) \right\| \leq \frac{C_t}{m^{\frac{1}{2}}} \|Wx\|.$$

Moreover, the convergence is uniform on any compact of \mathbb{R} .

Proof

1. Let $x \in \mathcal{D}(W)$, which implies that $Cx \in \mathcal{D}(W)$ because $WCx = CWx$. Let's remember that $T(\cdot)x : \mathbb{R} \rightarrow E$ is in $C^2(\mathbb{R}, E)$ and like

$$\begin{aligned} \int_{-\infty}^{+\infty} e^{-t^2} \|T(t)x\|^2 dt &\leq \int_{-\infty}^{+\infty} e^{-t^2} \|T(t)\|^2 \|x\|^2 dt \\ &\leq M^2 \int_{-\infty}^{+\infty} e^{-t^2} e^{2\omega|t|} \|x\|^2 dt \\ &\leq M^2 \|x\|^2 \int_{-\infty}^{+\infty} e^{-|t|^2 + 2\omega|t|} dt \\ &\leq M^2 \|x\|^2 e^{\omega^2} \int_{-\infty}^{+\infty} e^{-(|t|-\omega)^2} dt \\ &\leq M^2 \|x\|^2 e^{\omega^2} \int_{-\infty}^{+\infty} e^{-u^2} du \\ &< +\infty, \end{aligned}$$

then, the series $\sum_{n \in \mathbb{N}} c_n(T(\cdot)x)H_n$ with

$$c_n(T(\cdot)x) = \int_{-\infty}^{+\infty} \varphi_n(t) T(t)x dt = \frac{1}{2^n n!} W^n T_1\left(\frac{1}{4}\right)x$$

converges pointwise to $T(t)x$ for $t \in \mathbb{R}$, it is $T(t)x = \sum_{n=0}^{+\infty} \frac{1}{2^n n!} W^n T_1\left(\frac{1}{4}\right)x H_n(t)$.

2. • (i)

According to 3.(i) of Lemma 2.1, for all $x \in \mathcal{D}(W)$ and $z \in \mathbb{C}$, we have

$$\begin{aligned} \sum_{n=0}^{+\infty} \frac{\|W^n T_1\left(\frac{1}{4}\right)x\|}{2^{2n}n!} |4z-1|^n &\leq \left(\sup_{t \in \mathbb{R}} \|T(t)\| \right) \|x\| \sum_{n=0}^{+\infty} \frac{2^n \sqrt{(2n)!}}{2^{2n}n!} |4z-1|^n \\ &\leq c_1 \sum_{n=0}^{+\infty} \frac{\sqrt{(2n)!}}{2^n n!} |4z-1|^n \quad \left(c_1 = \left(\sup_{t \in \mathbb{R}} \|T(t)\| \right) \right) \end{aligned}$$

Stirling's formula $\frac{\sqrt{(2n)!}}{n!} \sim_{+\infty} 2^n n^{-\frac{1}{4}} (2\pi)^{-\frac{1}{4}} 2^{\frac{1}{4}}$ implies that:

$$c_1 \sum_{n=0}^{+\infty} \frac{\sqrt{(2n)!}}{2^n n!} |4z - 1|^n \leq c \sum_{n=0}^{+\infty} \frac{|4z - 1|^n}{n^{\frac{1}{4}}} < \infty, \text{ for } |z - \frac{1}{4}| < \frac{1}{4}.$$

Finally let a analytic family of operators :

$$V : D(\frac{1}{4}, \frac{1}{4}) = \{z \in \mathbb{C} / |z - \frac{1}{4}| < \frac{1}{4}\} \rightarrow B(\mathcal{D}(W)), z \mapsto V(z) :$$

$$\mathcal{D}(W) \rightarrow \mathcal{D}(W), x \mapsto V(z)x = \sum_{n=0}^{+\infty} \frac{W^n T_1(\frac{1}{4})x}{2^{2n} n!} (4z - 1)^n.$$

So For $x \in D(W)$:

$$\begin{aligned} V'(z)x &= \sum_{n=1}^{+\infty} \frac{W^n T_1(\frac{1}{4})x}{2^{2n} n!} 4n(4z - 1)^{n-1} \\ &= \sum_{n=0}^{+\infty} \frac{W^n T_1(\frac{1}{4})Wx}{2^{2n} n!} (4z - 1)^n \\ &= V(z)Wx \\ &= WV(z)x. \end{aligned}$$

Like W a generator of analytic C -semigroups $(T_1(t))_{t \geq 0}$ of angle $\frac{\pi}{2}$ and $V(\frac{1}{4})x = T_1(\frac{1}{4})x \in \mathcal{D}(W)$ then $V(t)x = T_1(t)x$ for $t \in]0, \frac{1}{2}[$ because the first order abstract Cauchy problem

$$\begin{cases} u'(t) = Wu(t), & t \in]0, \frac{1}{2}[\\ u(\frac{1}{4}) = T_1(\frac{1}{4})x \end{cases}$$

has unique solution defined in $]0, \frac{1}{2}[$. On the other hand the two functions $z \mapsto T_1(z)x$ and $z \mapsto V(z)x$ are holomorphic on the connected open set $D(\frac{1}{4}, \frac{1}{4})$, likewise $]0, \frac{1}{2}[\subset \{u \in D(\frac{1}{4}, \frac{1}{4}), V(u)x = T_1(u)x\}$, by the principle of analytical continuation we get: $V(z)x = T_1(z)x$ for $|z - \frac{1}{4}| < \frac{1}{4}$.

- (ii) Let $t \in \mathbb{R}$ be fixed. We are looking to increase the quantity

$$\left\| \sum_{n=0}^{+\infty} \frac{1}{2^{2n} (2n)!} W^n T_1(\frac{1}{4})x H_{2n}(t) - \sum_{n=0}^m \frac{1}{2^{2n} (2n)!} W^n T_1(\frac{1}{4})x H_{2n}(t) \right\|. \quad (9)$$

There exists $m_t \in \mathbb{N}^*$ such that $0 \leq t^2 \leq 2(2m_t + 1)$, according to inequality (1)

$$(\forall n \geq m_t) \quad |H_{2n}(t)| \leq c \left(\frac{e^{\frac{t^2}{2}} \sqrt{2^{2n} (2n)!}}{(2n)^{\frac{1}{2}}} \right) \quad (\text{the } c \text{ is independent of } t \text{ and it is the inequality constant constant (1)),$$

let $x \in \mathcal{D}(W)$, according to the inequality 3 - (ii) of lemma 2.1,

there is a constant $c_1 > 0$ such as for all $m \geq m_t$ from where:

$$\begin{aligned}
(9) &\leq \sum_{n=m+1}^{+\infty} \left\| \frac{1}{2^{2n}(2n)!} W^n T_1\left(\frac{1}{4}\right) x H_{2n}(t) \right\| \\
&\leq \sum_{n=m+1}^{+\infty} \left\| \int_{-\infty}^{+\infty} \varphi_{2n}(s) T(s) x ds \right\| \cdot \|H_{2n}(t)\| \\
&\leq \sum_{n=m+1}^{+\infty} \frac{c_1 \sqrt{(2n-2)!}}{2^{n+1}(2n)!} \|Wx\| c \frac{e^{\frac{t^2}{2}} \sqrt{2^{2n}(2n)!}}{(2n)^{\frac{1}{12}}} \quad (3.ii \text{ of lemma 2.1}) \\
&\leq \sum_{n=m+1}^{+\infty} \frac{cc_1 e^{2m_t+1} \sqrt{(2n-2)!(2n)!} 2^n}{2^{n+1}(2n)! 2^{\frac{1}{12}} n^{\frac{1}{12}}} \|Wx\| \\
&\leq \sum_{n=m+1}^{+\infty} c'_t \sqrt{\frac{(2n-2)!}{(2n)!}} \frac{1}{n^{\frac{1}{12}}} \|Wx\| \quad (c'_t = cc_1 e^{2m_t+1}) \\
&\leq \sum_{n=m+1}^{+\infty} \frac{c'_t}{\sqrt{2n(2n-1)} n^{\frac{1}{12}}} \|Wx\| \\
&\leq \sum_{n=m+1}^{+\infty} \frac{c'_t}{n^{\frac{13}{12}}} \|Wx\| \quad (\text{because } \frac{1}{\sqrt{2n(2n-1)}} \leq \frac{1}{n}).
\end{aligned}$$

The Riemann serie $\sum_{n \geq 1} \frac{1}{n^{\frac{13}{12}}}$ is convergent, so

$$\sum_{n=m+1}^{+\infty} \frac{1}{n^{\frac{13}{12}}} \sim_{m \rightarrow +\infty} \frac{1}{\frac{13}{12} - 1} \frac{1}{m^{\frac{13}{12} - 1}} = \frac{1}{\frac{1}{12}} \frac{1}{m^{\frac{1}{12}}} \text{ from where}$$

$$\begin{aligned}
\|T(t)x - \sum_{n=0}^m \frac{1}{2^{2n}(2n)!} W^n T_1\left(\frac{1}{4}\right) x H_{2n}(t)\| &\leq c'_t c_2 \frac{12}{m^{\frac{1}{12}}} \|Wx\| \\
&\leq \frac{c_t}{m^{\frac{1}{12}}} \|Wx\|.
\end{aligned}$$

- If $K \subset \mathbb{R}$ is a compact set then there is $m_0 \in \mathbb{N}$ for all $t \in K$, $0 \leq t^2 \leq 2(2m_0 + 1)$, so the constant c'_t (from where c_t) is independent of t therefore, the reminder tends uniformly on K towards 0.

Example 2.2. Let $m : \mathbb{R} \rightarrow \mathbb{R}^-$ be an even measurable function. In the Banach space $L^1(\mathbb{R})$, we consider the family $T := (T(t))_{t \in \mathbb{R}} \subset \mathfrak{F}(L^1(\mathbb{R}))$ defined by $T : \mathbb{R} \rightarrow B(L^1(\mathbb{R}))$, $t \mapsto T(t) : L^1(\mathbb{R}) \rightarrow L^1(\mathbb{R})$, $f \mapsto T(t)(f) : \mathbb{R} \rightarrow \mathbb{R}$, $s \mapsto T(t)(f)(s) = \cos(t\sqrt{-m(s)}) f(-s)$.

Clearly, $(T(t))_{t \in \mathbb{R}} \subset B(L^1(\mathbb{R}))$. If we put $T(0) = C$, then for all $s, t \in \mathbb{R}$ $T(t)(f)(s) = \cos(t\sqrt{-m(s)}) C(f)(s)$. We also have:

- (i) C is an injective element of $B(L^1(\mathbb{R}))$.

- (ii) For all $t, s, z \in \mathbb{R}$:

$$\begin{aligned}
 D(z) &= (T(t+s) + T(t-s))(C(f))(z) \\
 &= T(t+s)(C(f))(z) + T(t-s)(C(f))(z) \\
 &= \cos((t+s)\sqrt{-m(z)})C(f)(-z) + \cos((t-s)\sqrt{-m(z)})C(f)(-z) \\
 &= \cos((t+s)\sqrt{-m(z)})f(z) + \cos((t-s)\sqrt{-m(z)})f(z) \\
 &= (\cos(t\sqrt{-m(z)} + s\sqrt{-m(z)}) + \cos(t\sqrt{-m(z)} - s\sqrt{-m(z)}))f(z) \\
 &= 2\cos(t\sqrt{-m(z)})\cos(s\sqrt{-m(z)})f(z) \\
 &= 2\cos(t\sqrt{-m(z)})\cos(s\sqrt{-m(z)})(C(f))(-z) \\
 &= 2\cos(t\sqrt{-m(z)})T(s)(C(f))(z) \\
 &= 2\cos(t\sqrt{-m(z)})C(T(s)(f))(z) \quad (m \text{ is an even function}) \\
 &= 2T(t)(T(s)(f))(z) \\
 &= 2T(t)T(s)(f)(z)
 \end{aligned}$$

- (iii) For each f fixed in $L^1(\mathbb{R})$. The function $\mathbb{R} \rightarrow L^1(\mathbb{R}), t \mapsto T(t)(f)$ is continuous on \mathbb{R} .

- (iv) For all $f, g \in L^1(\mathbb{R})$ such that $C^{-1} \lim_{t \rightarrow 0} \frac{2(T(t)f - C(f))}{t^2} = g$, we have :

$$\begin{aligned}
 0 &= \lim_{t \rightarrow 0} \left\| \frac{2(T(t)(f) - C(f))}{t^2} - C(g) \right\|_{L^1(\mathbb{R})} \\
 &= \lim_{t \rightarrow 0} \int_{\mathbb{R}} \left| \frac{2(\cos(t\sqrt{-m(s)})C(f)(s) - C(f)(s))}{t^2} - C(g)(s) \right| ds \\
 &= \lim_{t \rightarrow 0} \int_{\mathbb{R}} \left| \frac{2(\cos(t\sqrt{-m(s)}) - 1)}{t^2} C(f)(s) - C(g)(s) \right| ds \\
 &= \int_{\mathbb{R}} \lim_{t \rightarrow 0} \left| \frac{2(\cos(t\sqrt{-m(s)}) - 1)}{t^2} C(f)(s) - C(g)(s) \right| ds \quad (\text{Fatou's lemma}) \\
 &= \int_{\mathbb{R}} |m(s).C(f)(s) - C(g)(s)| ds \\
 &= \int_{\mathbb{R}} |C(C(m).f - g)(s)| ds \quad (\text{because } \forall h \in \mathfrak{F}(\mathbb{R}, \mathbb{R}) \quad C \circ C(h) = h) \\
 &= \|C(m.f - g)\|_{L^1(\mathbb{R})}, \quad (C(m) = m \text{ because } m \text{ is an even function})
 \end{aligned}$$

- (v) Let $t \in \mathbb{R}$ fixed, we have for all $f \in L^1(\mathbb{R})$:

$$\|T(t)(f)\|_{L^1(\mathbb{R})} = \int_{\mathbb{R}} |\cos(t\sqrt{-m(s)})f(-s)| ds \leq \|f\|_{L^1(\mathbb{R})}.$$

Ultimately, $(T(t))_{t \in \mathbb{R}}$ is uniformly bounded C -regularized cosine function with generator $(W, \mathcal{D}(W))$ defined by W :

$\mathcal{D}(W) = \{f \in L^1(\mathbb{R}) / m.f \in L^1(\mathbb{R})\} \rightarrow L^1(\mathbb{R}), f \mapsto W(f) = m.f$. On the other hand, for all $s \in \mathbb{R}$ and $f \in L^1(\mathbb{R})$,

$$\begin{aligned} \frac{2}{\sqrt{\pi}} \int_0^{+\infty} e^{-t^2} T(t)(f)(s) dt &= \frac{2}{\sqrt{\pi}} \int_0^{+\infty} e^{-t^2} \cos(t\sqrt{-m(s)}) f(-s) dt \\ &= \frac{2}{\sqrt{\pi}} \int_0^{+\infty} e^{-t^2} \operatorname{Re} \left(e^{i(t\sqrt{-m(s)})} \right) f(-s) dt \\ &= \operatorname{Re} \left(\frac{2}{\sqrt{\pi}} \int_0^{+\infty} e^{-t^2 + it\sqrt{-m(s)}} f(-s) dt \right) \\ &= \operatorname{Re} \left(\frac{2}{\sqrt{\pi}} \int_0^{+\infty} e^{\frac{m(s)}{4}} e^{-\left(t - i\frac{\sqrt{-m(s)}}{2}\right)^2} f(-s) dt \right) \\ &= e^{\frac{m(s)}{4}} \left(\frac{2}{\sqrt{\pi}} \int_0^{+\infty} e^{-u^2} du \right) \cdot f(-s) \\ &= e^{\frac{m(s)}{4}} f(-s), \end{aligned}$$

then for all $s \in \mathbb{R}$ and $f \in L^1(\mathbb{R})$

$$T_1\left(\frac{1}{4}\right)(f)(s) = \left(\frac{2}{\sqrt{\pi}} \int_0^{+\infty} e^{-t^2} \cos(t\sqrt{-m(s)}) dt \right) \cdot f(-s) = e^{\frac{m(s)}{4}} f(-s),$$

Theorem 2.1 gives for $f \in \mathcal{D}(W)$ and $s, t \in \mathbb{R}$:

$$\begin{aligned} T(t)(f)(s) &= \sum_{n=0}^{+\infty} \frac{1}{2^{2n}(2n)!} W^n T_1\left(\frac{1}{4}\right)(f)(s) H_{2n}(t) \\ &= \sum_{n=0}^{+\infty} \frac{(m(s))^n}{2^{2n}(2n)!} e^{\frac{m(s)}{4}} f(-s) H_{2n}(t). \end{aligned}$$

$$\text{So } T(t)(f) = \sum_{n=0}^{+\infty} \frac{(m(\cdot))^n}{2^{2n}(2n)!} e^{\frac{m(\cdot)}{4}} H_{2n}(t) C(f).$$

References

- [1] W. ARENDT, C. J. K. BATTY, M. HIEBER, F. NEUBRANDER, Vector-valued Laplace Transforms and Cauchy Problems, Convolved C -cosine functions and semigroups. Birkh user Verlag, Basel, (2001).
- [2] R. ASKEY AND S., Mean convergence of expansions in Laguerre and Hermite series, Amer. J. Math. 87, 695-708 (1965).
- [3] S.S. BONAN, D. S. CLARK, Estimates of the Hermite and the Freud Polynomials, Journal of approximation theory 63, 210-244 (1990).
- [4] L.L. DAN, Hermite expansions of C_0 -groups and cosine functions, Journal of Mathematical Analysis and Applications, UniversidaddeZaragoza, 50009 Zaragoza, Spain (2015).

- [5] R. DELAUBENFELS, S. GUOZHENG, S. WANG, R. AFTABIZADEH, Regularized semigroups, Existence Families And the abstract Cauchy Problem, Differential and Integral Equations, Volume 8, p. 1477-1496 (1995).
- [6] M. KALTENBAVK, F. LEOPOLD, Generalisations of Semigroups of Operators in the View of Linear Relations, Analysis und Scientific Computing der Technischen Universitat Wien.
- [7] M. KALTENBACK, F.L. SCHWENNINGER, Generalisations of Semigroups of Operators in the View of Linear Relations, Analysis und Scientific Computing der Technischen Universitat Wien, Matr. Nr. 0625539 Wahringer Strasse 125/15-1180 Wien. Vienna University of Technology.
- [8] V. KEYANTUO, C. LIZAMA, P.J. MIANA, Algebra homomorphisms defined via convoluted semigroups and cosine functions, Journal of Functional Analysis 257. 3454-3487 (2009).
- [9] E. KOGBETLIANTZ, Recherches sur la sommabilite des series d'Hermite, Annales scientifiques de l'E.N.S. 3^e serie, tome 49 (1932).
- [10] M. KOSTIC, (a, k) -Regularized C -Resolvent Families: Regularity and Local Properties, Hindawi Publishing Corporation Abstract and Applied Analysis, Volume 2009, Article ID 858242, 27 pages doi:10.1155/2009/858242.
- [11] C.C. KUO, Perturbations of local C -cosine functions, Stud. Univ. Babeş-Bolyai Math. 65, No.4, 585-597 (2020).
- [12] C.C. KUO, S.Y. SHAW, C -Cosine Functions and the Abstract Cauchy Problem, Journal of Mathematical Analysis and Applications 210, 632-646 (1997).
- [13] N.N. LEBEDEV, Special functions and their applications, prentice-hall, inc (1965).
- [14] F. LI, J. LIANG, T.J. XIAO, J. ZHANG, On Perturbation of Convoluted-Regularized Operator Families, Hindawi Publishing Corporation Journal of Function Spaces and Applications, Volume 2013, Article ID 579326.
- [15] F. LI, J.H. LIU, Note on multiplicative perturbation of local C -regularized cosine functions with nondensely defined generators, Electronic Journal of Qualitative Theory of Differential Equations, No. 57, 1-12 (2010).
- [16] M. LI, Q. ZHENG, J. ZHANG, Regularized resolvent families, Taiwanese Journal of Mathematics, vol.11, no.1, p. 117-133 (2007).
- [17] M. MOSALLANEZHADA, M. JANFADAB, On Mixed C -Semigroups of Operators on Banach Spaces, Filomat 30:10, 2673-2682 (2016).

- [18] P. PREDÁ, A. POGAN, C. PREDÁ, The Perron Problem For C -SEM, *Math. J. Okayama Univ.* 46, 141-151 (2004).
- [19] J. A. SHOHAAT, On the developement of functions in series of orthogonal polynomials, The university of Pennsylvania (1935).
- [20] H. SKOVGAARD, Asymptotic forms of hermite polynomials. Departement of Mathematics California in scitute of Technology Pasadenea (1959).
- [21] G. SZEGŐ, Orthogonal polynomials, PubLished by the American Mathematical Society Providence, Rhode Island (1939).
- [22] S.W. WANG, Properties of subgenerators of C -Regularized semigroups, proceedings of the American Mathematical Society, Volume 126, Number 2, p.453-460 (1998).
- [23] J. ZHANG, Regularized cosine Functions and Polynomials of Group Generators, *Calculi, Scientia iranica*, vol 4, Nos.1, 2, pp 12-18 (1997).
- [24] Q. ZHENG, Matrices of operators and regularized cosine functions, *J. Math. Anal. Appl.*315, 68-75 (2006).
- [25] Q. ZHENG, Y. ZHAO, An analyticity criterion for regularized semigroups, proceedings of the american Mathematical society, Volume 130, Number 8, S 0002-9939(02)06219-6 (2002).