

THE TARSKI-LINDENBAUM ALGEBRA OF THE
CLASS OF PRIME MODELS WITH INFINITE
ALGORITHMIC DIMENSIONS HAVING
OMEGA-STABLE THEORIES

M.G. PERETYAT'KIN 

Communicated by S.V. SUDOPLATOV

Abstract: We study the class of all prime strongly constructivizable models of infinite algorithmic dimensions having ω -stable theories in a fixed finite rich signature. It is proved that the Tarski-Lindenbaum algebra of this class considered together with a Gödel numbering of the sentences is a Boolean Σ_1^1 -algebra whose computable ultrafilters form a dense subset in the set of all ultrafilters; moreover, this algebra is universal with respect to the class of Boolean Σ_1^1 -algebras. This gives a characterization to the Tarski-Lindenbaum algebra of the class of all prime strongly constructivizable models of infinite algorithmic dimensions having ω -stable theories.

Keywords: Tarski-Lindenbaum algebra, strongly constructive model, computable isomorphism, semantic class of models, ω -stable theory, prime model.

The problem of characterizing Tarski-Lindenbaum algebras of some semantic classes of models of a finite rich signature was considered in the works

PERETYAT'KIN M.G., THE TARSKI-LINDENBAUM ALGEBRA OF THE CLASS OF PRIME MODELS WITH INFINITE ALGORITHMIC DIMENSIONS HAVING OMEGA-STABLE THEORIES.

© 2024 PERETYAT'KIN M.G.

This research is funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan, Grant No BR20281002.

Received December, 14, 2023, Published April, 8, 2024.

[10], [7], [11], [12], [13], and others. In this work, we describe the Tarski-Lindenbaum algebra of the class of all prime strongly constructivizable models of infinite algorithmic dimensions having ω -stable theories.

Preliminaries. We consider theories in first-order predicate logic *with equality* and use general concepts of model theory, algorithm theory, Boolean algebras and constructive models found in [6], [14], [15], [1], and [4]. We consider signatures admitting Gödel's numberings of the formulas. Generally, *incomplete theories* are considered.

A finite signature is called *rich*, if it contains at least one n -ary predicate or function symbol for $n > 1$, or two unary function symbols. A theory F is called *finitely axiomatizable* if it is defined by a finite set of axioms, and its signature is finite. The following notations are used: $FL(\sigma)$ is the set of all formulas of signature σ , $FL_n(\sigma)$ is the set of all formulas of signature σ with n free variables x_0, x_1, \dots, x_{n-1} , $SL(\sigma)$ is the set of all sentences (i.e., closed formulas) of signature σ . In the work, we use a finite rich signature σ , and consider a fixed Gödel numbering $\Phi_i, i \in \mathbb{N}$, of the set $SL(\sigma)$. The set of all Gödel numbers of formulas from a set $\Sigma \subseteq SL(\sigma)$ is denoted by $Nom(\Sigma)$. We use notation $\mathcal{P}(A)$ for the power-set of A , and $|A|$ or $Card(A)$ for cardinality of the set A . The set of all finite tuples α of the form $\alpha = \langle \alpha_0, \alpha_1, \dots, \alpha_n \rangle, \alpha_i \in \{0, 1\}$, is denoted by $2^{<\omega}$. The empty tuple is denoted by \emptyset . The *canonical (Gödel) index* of a finite tuple of zeros and ones of the form $\varepsilon = \langle \varepsilon_0, \varepsilon_1, \dots, \varepsilon_{n-1} \rangle, \varepsilon_i \in \{0, 1\}$, is the number $Nom(\varepsilon) = 2^n + \varepsilon_0 2^{n-1} + \varepsilon_1 2^{n-2} + \dots + \varepsilon_{n-1} - 1$. We often write shortly $\langle \varepsilon \rangle$ instead of $Nom(\varepsilon)$.

A countable or finite model \mathfrak{M} together with a numeration $\nu : \mathbb{N} \xrightarrow{onto} |\mathfrak{M}|$ is called a *numerated model* and is denoted by (\mathfrak{M}, ν) . We denote by \mathfrak{M}_ν the model obtained by enriching signature of \mathfrak{M} to $\sigma' = \sigma \cup \{c_i \mid i \in \mathbb{N}\}$ in such a way that each new constant c_i is interpreted by $\nu(n)$ in $|\mathfrak{M}|$. A numerated model (\mathfrak{M}, ν) is said to be *constructive* if the atomic diagram $AD(\mathfrak{M}_\nu)$ is computable, and (\mathfrak{M}, ν) is *strongly constructive* if the theory $Th(\mathfrak{M}_\nu)$ is decidable. Two numerated models (\mathfrak{N}, ν) and (\mathfrak{M}, v) are said to be *computably isomorphic*, denoted $(\mathfrak{N}, \nu) \cong (\mathfrak{M}, v)$, if there is an isomorphism μ between \mathfrak{N} and \mathfrak{M} , and two computable functions $f(x)$ and $g(x)$, such that the following diagram is commutative:

$$\begin{array}{ccc}
 \mathbb{N} & \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{g} \end{array} & \mathbb{N} \\
 \nu \downarrow & & \downarrow v \\
 \mathfrak{N} & \xrightarrow{\mu} & \mathfrak{M}
 \end{array}$$

In a parallel way, we also write $\mathfrak{N} \cong \mathfrak{M}$ to indicate that abstract models \mathfrak{N} and \mathfrak{M} (without numerations) are isomorphic.

We consider Boolean algebras in the signature $\sigma_{BA} = \{\cup, \cap, -, \mathbf{0}, \mathbf{1}\}$. Let \mathcal{B} be a Boolean algebra, and $a \in \mathcal{B}$. By $\mathcal{B}[a]$, we denote the restriction of the Boolean algebra \mathcal{B} on the set of all subelements of the element a counting that $\mathbf{1} = a$ and $-x$ is defined as $a \setminus x$ in $\mathcal{B}[a]$. If b is an element in a Boolean algebra and $\alpha \in \{0, 1\}$, then b^α means b for $\alpha = 1$ and $-b$ for $\alpha = 0$.

Similarly, if Φ is a formula and $\alpha \in \{0, 1\}$, then Φ^α means Φ for $\alpha = 1$ and $\neg\Phi$ for $\alpha = 0$.

Let T be a theory of signature σ . On the set of sentences $SL(\sigma)$, an equivalence relation \sim_T is defined by the rule $\Phi \sim_T \Psi \Leftrightarrow T \vdash (\Phi \leftrightarrow \Psi)$. The logical connectives \vee , $\&$, and \neg generate Boolean operations \cup , \cap , and $-$ on the quotient set $SL(\sigma)/\sim_T$; One can easily check that, these operations are well-defined on the \sim_T -classes. Thereby, we obtain an algebra of the form

$$L(T) = (SL(\sigma)/\sim_T; \cup, \cap, -, \mathbf{0}, \mathbf{1}),$$

that, in fact, is a Boolean algebra. It is called the *Tarski-Lindenbaum algebra* of the theory T ; alternatively, it is called the *Boolean sentence algebra* of T , cf. [3]. By $\mathcal{L}(T)$, we denote the Tarski-Lindenbaum algebra $L(T)$ considered together with a Gödel numbering γ ; thereby, the concept of a computable isomorphism is applicable to such objects. A similar construction with respect to formulas with n free variables gives a definition of the *Tarski-Lindenbaum algebra* $L_n(T)$ of theory T with n free variables, $n < \omega$. Such algebras are often considered in the case when the theory T is complete.

Following Rogers, we use the notation W_n for n th computably enumerable set in Post's numbering of the family of all c.e. sets, [14, Sec. 5.2]. Moreover, W_n^t is a finite part of the set W_n that can be computed in t steps. Furthermore, we denote by W_n^X the computably enumerable set with c.e. index n relative to computability with an oracle $X \subseteq \mathbb{N}$, [14, Sec. 9.3].

We give an example of a complete set in the hierarchy class Π_1^1 .

Lemma 0.1. *There is a computable sequence (\mathcal{L}_i, ν_i) , $i < \omega$, of constructive linear orderings such that*

$$\mathcal{W} = \{n \mid \mathcal{L}_n \text{ is a well-ordered set}\} \tag{0.1}$$

is an m -complete in the class Π_1^1 set.

PROOF. See a construction in [9, p. 44]. □

1 Boolean algebras over classes of hierarchies

Let Ξ be a class of a hierarchy, eg. Σ_n^0 , Δ_n^0 , Π_n^0 , Σ_n^1 , or Π_n^1 with $0 < n < \omega$ (actually, we do not concern other classes of hierarchies in this paper). A numerated Boolean algebra (\mathcal{B}, ν) is called a Ξ -algebra, if all its signature operations are uniformly presentable by computable functions on ν -numbers, while the equality predicate in \mathcal{B} is a Ξ -relation in the numeration ν ; i.e., there exist computable functions $u(x, y)$, $v(x, y)$, $w(x)$, and a relation $E(x, y)$ which represent the Boolean algebra as follows, for any $x, y \in \mathbb{N}$:

$$\begin{aligned} \nu(x) \cup \nu(y) &= \nu(u(x, y)), \\ \nu(x) \cap \nu(y) &= \nu(v(x, y)), \\ -\nu(x) &= \nu(w(x)), \\ \nu(x) = \nu(y) &\Leftrightarrow E(x, y), \quad E \in \Xi. \end{aligned} \tag{1.1}$$

A Boolean algebra \mathcal{B} is said to be a Ξ -algebra, if there is a numeration ν such that (\mathcal{B}, ν) is a Boolean Ξ -algebra.

As it is accepted, cf. [14, Sec. 14.1], the class of all relations on \mathbb{N} expressible by Σ_n^X -forms is denoted by Σ_n^X , while the class of all relations expressible by Π_n^X -forms is denoted by Π_n^X ; we also denote $\Delta_n^X = \Sigma_n^X \cap \Pi_n^X$. In particular, Σ_1^X is the class of all X -enumerable relations on the set \mathbb{N} .

A numerated Boolean algebra (\mathcal{B}, ν) is said to be X -enumerable if all operations in (\mathcal{B}, ν) are presented by computable functions, while the equality relation is X -enumerable in the numeration ν . A Boolean algebra \mathcal{B} is said to be an X -enumerable algebra if there is a numeration ν such that (\mathcal{B}, ν) is an X -enumerable Boolean algebra.

Lemma 1.1. *For an arbitrary numerated Boolean Σ_1^X -algebra (\mathcal{B}, ν) , there is a numeration ν of \mathcal{B} such that (\mathcal{B}, ν) is a Boolean Σ_1^X -algebra whose computable ultrafilters form a dense set in the set of all ultrafilters of the algebra (\mathcal{B}, ν) .*

PROOF. Cf. [12, Lem. 2]. □

2 An isomorphism criterion for Boolean sentence algebras

Generalizing constructions of computable isomorphisms in the works [11], [12], and [13], we obtain the following criterion for isomorphism of a given numerated Boolean algebra to a segment of the Tarski-Lindenbaum algebra of a class of models.

Theorem 2.1. *Given a numerated Boolean algebra (\mathcal{B}, ν) together with a class K of models of a finite rich signature σ and a sentence Φ of the signature σ . Suppose that $g_i, i \in \mathbb{N}$, is a generating set for \mathcal{B} , and $\theta_i, i \in \mathbb{N}$ is a generating set for $\mathcal{L}(\text{Th}(K))$. A partial mapping λ^* defined by rule $\lambda^*(g_i) = \theta_i, i \in \mathbb{N}$, uniquely determines an isomorphism*

$$\lambda : \mathcal{B} \rightarrow \mathcal{L}(\text{Th}(K \cap \text{Mod}_\sigma(\Phi))) \quad (2.1)$$

if and only if for all α in $2^{<\omega}$, $\alpha = \langle \alpha_0, \alpha_1, \dots, \alpha_k \rangle$, we have:

$$g_0^{\alpha_0} \cap \dots \cap g_k^{\alpha_k} \neq \mathbf{0} \Leftrightarrow \Phi \& \theta_0^{\alpha_0} \& \dots \& \theta_k^{\alpha_k} \text{ has a } K\text{-model.} \quad (2.2)$$

Moreover, the mapping λ represents a computable isomorphism

$$\lambda : (\mathcal{B}, \nu) \rightarrow (\mathcal{L}(\text{Th}(K \cap \text{Mod}_\sigma(\Phi))), \gamma) \quad (2.3)$$

with respect to a Gödel numbering γ of $SL(\sigma)$, whenever the sequence $g_i, i \in \mathbb{N}$, is computable in (\mathcal{B}, ν) , while $\theta_i, i \in \mathbb{N}$, is computable in the Gödel numbering γ .

PROOF. As a prototype for this statement, we use a well-known result [15, 13.3].

For any tuple α in $2^{<\omega}$, $\alpha = \langle \alpha_0, \dots, \alpha_k \rangle$, consider elements:

$$\begin{aligned} b_\alpha &= g_0^{\alpha_0} \cap \dots \cap g_k^{\alpha_k}, \\ \theta_\alpha &= \theta_0^{\alpha_0} \& \dots \& \theta_k^{\alpha_k}. \end{aligned} \quad (2.4)$$

For an arbitrary finite set of tuples $\alpha, \beta, \dots, \delta \in 2^{<\omega}$, we put

$$\lambda(b_\alpha \cup b_\beta \cup \dots \cup b_\delta) = \Theta_\alpha \vee \Theta_\beta \vee \dots \vee \Theta_\delta. \tag{2.5}$$

Based on (2.2), it is possible to show that λ is a bijective mapping from \mathcal{B} to $\mathcal{L}(\text{Th}(K \cap \text{Mod}(\Phi)))$ because the expressions involved in (2.5) cover all elements of the algebras we are considering; moreover, λ is an isomorphism from \mathcal{B} into $\mathcal{L}(\text{Th}(K \cap \text{Mod}(\Phi)))$. Obviously, λ is a computable isomorphism in numerations ν and γ provided that the generating sequences are computable in these numerations.

Theorem 2.1 is proved. □

3 Strong constructivizations of prime models

A model \mathfrak{N} is called *prime* if it can be elementarily embedded in any other model of theory $T = \text{Th}(\mathfrak{N})$. Vaught’s criterion in [16] states that a countable complete theory T has a prime model if and only if the family of principal type in of this theory is dense in the family of all types (equivalently, all Tarski-Lindenbaum algebras $L_n(T)$, $n < \omega$, are atomic), while the model \mathfrak{N} of theory T is prime if and only if any finite tuple $\bar{a} \in |\mathfrak{N}|$ realizes a principal type of this theory.

Let us consider some technical facts about prime models.

Lemma 3.1. *Let T be a complete decidable theory. The set \mathcal{A}' of all formulas of the Tarski-Lindenbaum algebras $\mathcal{L}_n(T)$, $n \in \mathbb{N}$ that are not atoms of theory T is computably enumerable.*

PROOF. A formula $\varphi(\bar{x})$ is not an atom of theory T if either it is inconsistent with T or there exists a formula $\psi(\bar{x})$ such that both formulas $\varphi(\bar{x}) \ \& \ \psi(\bar{x})$ and $\varphi(\bar{x}) \ \& \ \neg\psi(\bar{x})$ are compatible with T . Based on this, it is simple to construct an effective enumeration of the set \mathcal{A}' . □

Lemma 3.2. *Let T be a complete decidable theory. The set \mathcal{A} of atomic formulas of the Tarski-Lindenbaum algebras $\mathcal{L}_n(T)$, $n \in \mathbb{N}$ is computably enumerable if and only if \mathcal{A} is computable, i.e. \mathcal{A} is uniformly computable in the parameter n .*

PROOF. Implication \Leftarrow is obvious. As for the implication \Rightarrow , this assertion is a simple consequence of Lemma 3.1. □

A model \mathfrak{N} is said to be *strongly constructivizable* if it admits a strong constructivization. A model \mathfrak{N} has algorithmic dimension 1, symbolically $\text{dim}_{s.c.}(\mathfrak{N}) = 1$, if \mathfrak{N} is strongly constructivizable, moreover, any two strong constructivizations ν_1 and ν_2 \mathfrak{N} are equivalent, that is the numbered models (\mathfrak{N}, ν_1) and (\mathfrak{N}, ν_2) are computably isomorphic, symbolically $(\mathfrak{N}, \nu_1) \cong (\mathfrak{N}, \nu_2)$. If $\text{dim}_{s.c.}(\mathfrak{N}) = 1$, we say that the model \mathfrak{N} is *autostable with respect to strong constructivizations* or simply *autostable*, since the case of weak constructivizations is not considered in this paper. A model \mathfrak{N} has algorithmic dimension ω , symbolically $\text{dim}_{s.c.}(\mathfrak{N}) = \omega$, if there is an infinite sequence of strong constructivizations ν_i , $i < \omega$ of this model such that any two numerated models in the sequence (\mathfrak{N}, ν_i) , $i < \omega$ are not isomorphic.

It was proved in [8] that, for any strongly constructivizable model \mathfrak{N} only cases $\dim_{s.c.}(\mathfrak{N}) = 1$ and $\dim_{s.c.}(\mathfrak{N}) = \omega$ are possible, while the cases of finite algorithmic dimensions $\dim_{s.c.}(\mathfrak{N}) = n$, $1 < n < \omega$, are impossible. Namely, the following assertion is proved.

Theorem 3.3. (Nurtazin criterion on autostability). *Let \mathfrak{M} be a strongly constructivizable model of a complete decidable theory T . Then the following claims are equivalent:*

- (a) \mathfrak{M} is autostable with respect to strong constructivizations,
- (b) there is a finite sequence \bar{a} of elements $|\mathfrak{M}|$ such that enrichment (\mathfrak{M}, \bar{a}) of the model \mathfrak{M} with these constants is a prime model and the family of sets of atoms of Tarski-Lindenbaum algebras $\mathcal{L}_n(T')$, $n \in \mathbb{N}$, of theory $T' = \text{Th}(\mathfrak{M}, \bar{a})$ is computable, i.e., the indicated family is uniformly computable in the parameter n .

Moreover, if conditions (a) and (b) are not satisfied, there exists an infinite sequence ν_i , $i < \omega$ of strong constructivizations of the model \mathfrak{M} such that, for any $i, j < \omega$, $i \neq j$, numbered models (\mathfrak{M}, ν_i) and (\mathfrak{M}, ν_j) are not computably embeddable in each other.

PROOF. See [8, Th. 1]. □

Now, we present two key statements for prime models.

Theorem 3.4. *Let T be a complete decidable theory with a prime model \mathfrak{N} . The model \mathfrak{N} is strongly constructivizable if and only if the set of principal types of the theory T is computable.*

PROOF. See [2, Th. 1] or [5, Sec. 1]. □

Theorem 3.5. *Let T be a complete decidable theory with a strongly constructivizable prime model \mathfrak{N} . The following statements are equivalent to each other:*

- (a) the set of formulas \mathcal{A} presenting atoms in the Tarski-Lindenbaum algebras $\mathcal{L}_n(T)$, $n \in \mathbb{N}$, is computably enumerable,
- (b) the set of formulas \mathcal{A} presenting atoms in the Tarski-Lindenbaum algebras $\mathcal{L}_n(T)$, $n \in \mathbb{N}$, is computable uniformly in n ,
- (c) the model \mathfrak{N} is autostable with respect to strong constructivizations.

PROOF. The proof is based on the Nurtazin criterion [8].

(a) \Rightarrow (b) This implication follows from Lemma 3.2.

(b) \Rightarrow (c) By assumption, the prime model \mathfrak{N} is strongly constructivizable, while the theory T is complete and decidable. We also suppose that the set of atoms \mathcal{A} of theory T is computable. Applying an effective version of Vaught's construction for an isomorphism between elementary equivalent prime models, [16], it is possible to construct a computable isomorphism $\mu : (\mathfrak{N}, \nu) \rightarrow (\mathfrak{N}, \nu')$ for any two strong constructivizations ν and ν' of the model \mathfrak{N} .

(c) \Rightarrow (a) Assume that the prime model \mathfrak{N} is autostable with respect to strong constructivizations, i.e., Part (a) of Theorem 3.3 holds. Then, Part (b) of Theorem 3.3 must also hold, i.e., there is a finite sequence \bar{a} of elements in $|\mathfrak{N}|$ such that the model $\mathfrak{N}^* = (\mathfrak{N}, \bar{a})$ of theory $T^* = \text{Th}(\mathfrak{N}, \bar{a})$ of signature

$\sigma^* = \sigma \cup \{a_i \mid a_i \in \bar{a}\}$ is a prime model and the family of atoms of the Tarski-Lindenbaum algebras $\mathcal{L}_n(\text{Th}(\mathfrak{N}, \bar{a}))$, $n \in \mathbb{N}$, is computable uniformly in n . By condition, the model \mathfrak{N} itself is prime. Since any finite tuple in a prime model realizes a principal type, there is a formula $\theta(\bar{z})$ that is an atom of theory T such that $\mathfrak{N} \models \theta(\bar{a})$ holds.

We consider a technical statement linking the theories T and T^* :

$$\begin{aligned} \phi(\bar{x}) \in FL(\sigma) \text{ is an atom of } T &\Leftrightarrow \text{there is } \varphi(\bar{x}, \bar{z}) \in FL(\sigma) \text{ s.t.} & (3.1) \\ \varphi(\bar{x}, \bar{a}) \text{ is an atom of } T^* &\text{, and } T \vdash \phi(\bar{x}) \leftrightarrow (\exists \bar{z})[\varphi(\bar{x}, \bar{z}) \& \theta(\bar{z})]. \end{aligned}$$

First, we suppose that formula $\phi(\bar{x})$ is an atom of theory T . Consider a tuple \bar{c} in \mathfrak{N} such that $\mathfrak{N} \models \phi(\bar{c})$. Since model \mathfrak{N} is prime, all types realized in \mathfrak{N} are principal. From this, it follows that there is a formula $\varphi(\bar{x}, \bar{z})$, which is an atom of theory T , and, in \mathfrak{N} , we have $\varphi(\bar{c}, \bar{a})$. Moreover, $\varphi(\bar{c}, \bar{a}) \& \theta(\bar{a})$ holds because, by assumption, we have $\mathfrak{N} \models \theta(\bar{a})$. Thus, formula $\varphi'(\bar{x}, \bar{z}) = \varphi(\bar{x}, \bar{z}) \& \theta(\bar{z})$ is satisfied in \mathfrak{N} on the tuple $\xi = (\bar{c}, \bar{a})$. Therefore, the formula $\varphi'(\bar{x}, \bar{z})$ is equivalent in T to the formula $\varphi(\bar{x}, \bar{z})$ since the latter is an atom of theory T . As a result, we obtain that, in theory T , it is provable $\phi(\bar{x}) \sim (\exists \bar{z})[\varphi(\bar{x}, \bar{z}) \& \theta(\bar{z})]$.

Now, we suppose that $\varphi(\bar{x}, \bar{a})$ is an atom of theory T^* . Consider the formula

$$\phi(\bar{x}) = (\exists \bar{z})[\varphi(\bar{x}, \bar{z}) \& \theta(\bar{z})]. \tag{3.2}$$

Our goal is to show that this formula is an atom of the theory T . For this, it is sufficient to show that, for arbitrary tuples \bar{c}_1 and \bar{c}_2 in \mathfrak{N} of appropriate lengths, it follows from $\mathfrak{N} \models \phi(\bar{c}_1)$ and $\mathfrak{N} \models \phi(\bar{c}_2)$ that there is an automorphism μ of the model \mathfrak{N} such that $\mu(\bar{c}_1) = \bar{c}_2$. Since $\mathfrak{N} \models \phi(\bar{c}_1)$, the right-hand side expression in (3.2) ensures that there is a tuple \bar{e}_1 in \mathfrak{N} such that $\mathfrak{N} \models \varphi(\bar{c}_1, \bar{e}_1) \& \theta(\bar{e}_1)$. Similarly, by virtue of $\mathfrak{N} \models \phi(\bar{c}_2)$, expression (3.2) provides that there is a tuple \bar{e}_2 in \mathfrak{N} such that $\mathfrak{N} \models \varphi(\bar{c}_2, \bar{e}_2) \& \theta(\bar{e}_2)$. Since the formula $\theta(\bar{z})$ is an atom of theory T , and, in the model \mathfrak{N} both $\theta(\bar{a})$ and $\theta(\bar{e}_1)$ hold, there must be an automorphism $\mu_1 : \mathfrak{N} \rightarrow \mathfrak{N}$ sending \bar{e}_1 to \bar{a} . Denote $\mu_1(\bar{c}_1)$ by \bar{c}'_1 . Similarly, both $\theta(\bar{a})$ and $\theta(\bar{e}_2)$ are satisfied in the model \mathfrak{N} , so there must be an automorphism $\mu_2 : \mathfrak{N} \rightarrow \mathfrak{N}$ sending \bar{e}_2 to \bar{a} . Denote $\mu_2(\bar{c}_2)$ by \bar{c}'_2 . In view of the properties of automorphisms μ_1 and μ_2 , we have $\mathfrak{N} \models \varphi(\bar{c}'_1, \bar{a}) \& \theta(\bar{a})$ and $\mathfrak{N} \models \varphi(\bar{c}'_2, \bar{a}) \& \theta(\bar{a})$. By assumption, formula $\varphi(\bar{x}, \bar{a})$ is an atom of theory T^* . Therefore, there is an automorphism $\mu_0 : \mathfrak{N} \rightarrow \mathfrak{N}$ that is identical on tuple \bar{a} and maps tuple \bar{c}'_1 into tuple \bar{c}'_2 . It remains to verify that the composition of automorphisms $\mu = \mu_2^{-1} \mu_0 \mu_1$ ensures the relations $\mu(\bar{c}_1) = \bar{c}_2$ and $\mu(\bar{a}) = \bar{a}$. This shows that the formula (3.2) is an atom of theory T .

Thereby, the relation (3.1) indeed takes place, thus allowing us to complete proof of the part (c) \Rightarrow (a). Based on the presentation (3.1), it is possible to organize an effective enumeration of the set \mathcal{A} of atoms of theory $T = \text{Th}(\mathfrak{N})$ that is exactly what is required.

Theorem 3.5 is proved.

4 Compact binary trees and the canonical construction

Definition of the notion of a compact binary tree can be found in [9, Sec, 2.1]. In this work, we use for such trees a more specialized term *compact binary trees* instead.

A *full compact binary tree* is a partially ordered set $\mathcal{D}_0 = \langle \mathbb{N}, \preceq \rangle$ of the form shown in Fig. 4.1(a). In particular, $1 \preceq 8, 6 \preceq 6, \neg(6 \preceq 9)$, and $\neg(9 \preceq 6)$ is satisfied. There are two natural operations within the tree, $L(n) = 2n + 1$ is the *left successor*, and $R(n) = 2n + 2$ is the *right successor* of an element $n \in \mathbb{N}$. A nonempty set $\mathcal{D} \subseteq \mathbb{N}$ is said to be a *compact binary tree* whenever the following conditions are satisfied: $m \preceq n \ \& \ n \in \mathcal{D} \Rightarrow m \in \mathcal{D}$, and $L(n) \in \mathcal{D} \Leftrightarrow R(n) \in \mathcal{D}$, for all $m, n \in \mathbb{N}$. An element n in a compact binary tree \mathcal{D} such that $L(n) \notin \mathcal{D}$ is said to be a *dead end* of the tree \mathcal{D} . The set of all dead-end elements in a tree \mathcal{D} is denoted by $\text{Dend}(\mathcal{D})$. A tree is called *atomic* if, above each of its elements, there is at least one dead-end element.

A few examples of compact binary trees are given in Fig. 4.1(a,b,c).




Fig. 4.1. Examples of compact binary trees

A natural operation of the *direct sum* of two given trees \mathcal{D}_1 and \mathcal{D}_2 is available. The tree $\mathcal{D}_1 \oplus \mathcal{D}_2$ is obtained by attaching isomorphic copies of the trees \mathcal{D}_1 and \mathcal{D}_2 to the two dead-end elements in the three-element tree given in Fig. 4.1(b) as it is schematically shown in Fig. 4.2.




Fig. 4.2. Scheme of the operations \oplus for compact binary trees

A set $\pi \subseteq \mathcal{D}$ is called a *chain* in the tree \mathcal{D} , if $m, n \in \pi \Rightarrow m \preceq n \vee n \preceq m$ and $m \preceq n \ \& \ n \in \pi \Rightarrow m \in \pi$, for all $m, n \in \mathbb{N}$. If \mathcal{D} is a compact binary tree, we denote by $\Pi(\mathcal{D})$ the set of all *maximal chains*, while $\Pi^{fin}(\mathcal{D})$ is the set of all *maximal finite chains* in the tree \mathcal{D} . A tree \mathcal{D} is said to be *superatomic*, if the set $\Pi(\mathcal{D})$ is at most countable.

See [9, Sec. 2.1] for definitions concerning the trees in detail.

Lemma 4.1. *There is a tree \mathcal{D}^* which is a superatomic (the more, atomic), computably enumerable, non-computable compact binary tree for which both families $\Pi^{fin}(\mathcal{D}^*)$ and $\Pi(\mathcal{D}^*)$ are computable.*

PROOF. Let \mathcal{D} be the tree shown in Fig. 4.1(c). Since \mathcal{D} is computable, the set $\text{Dend}(\mathcal{D})$ is also computable. Moreover, $\text{Dend}(\mathcal{D})$ is infinite, thus there is a computable bijective mapping $f : \mathbb{N} \rightarrow \text{Dend}(\mathcal{D})$. Let $K \subseteq \mathbb{N}$ be a computably enumerable non-computable set. Let $\mathcal{D}^* = \mathcal{D} \cup \{L(f(i)) \mid i \in$

$K\} \cup \{R(f(i)) \mid i \in K\}$. It is simple to check that the set \mathcal{D}^* is an atomic computably enumerable compact binary tree. Moreover, both families $\Pi(\mathcal{D}^*)$ and $\Pi^{fin}(\mathcal{D}^*)$ are computable. In particular, $\Pi(\mathcal{D}^*)$ is countable ensuring that the tree \mathcal{D}^* is superatomic. By construction, reducibility $n \in K \Leftrightarrow L(f(n)) \in \mathcal{D}^*$ takes place, thus the tree \mathcal{D}^* is not computable.

Lemma 4.1 is proved. □

Lemma 4.2. *There is an effective operator sending an arbitrary constructive linear ordering (\mathcal{L}, ν) into a tree $\mathcal{D}(\mathcal{L}, \nu)$ such that the following claims hold:*

- (a) *for any constructive linear ordering (\mathcal{L}, ν) , the tree $\mathcal{D}(\mathcal{L}, \nu)$ is computable,*
- (b) *the tree $\mathcal{D}(\mathcal{L}, \nu)$ is superatomic if and only if \mathcal{L} is well ordered,*
- (c) *if \mathcal{L} is well ordered, the family of chains $\Pi(\mathcal{D}(\mathcal{L}, \nu))$ is computable.*

PROOF. See [9, Lem. 2.2.1]. □

By \mathcal{D}_n , we denote the closure of the set W_n up to a compact binary tree, while \mathcal{D}_n^X is the closure of the set W_n^X up to a compact binary tree. It can be checked that the tree \mathcal{D}_n is computably enumerable, while the tree \mathcal{D}_n^X is computably enumerable with oracle X . Moreover, each c.e. tree is presented in the sequence $\mathcal{D}_n, n \in \mathbb{N}$, and each c.e. tree in computation with oracle X is presented in the sequence $\mathcal{D}_n^X, n \in \mathbb{N}$. The number n is considered as a c.e. index for the tree \mathcal{D}_n , and n is considered as a c.e. index for the tree \mathcal{D}_n^X considered in computability with an oracle X .

We denote by \mathfrak{P}_n the truth-table condition with the Gödel number n , cf. [9, Sec. 3.1]. We write $A \models \mathfrak{P}_n$ to indicate that \mathfrak{P}_n is satisfied in the set $A \subseteq \mathbb{N}$. The set of the form

$$\Omega(m) = \{A \subseteq \mathbb{N} \mid (\forall i \in W_m) A \models \mathfrak{P}_i\}, \tag{4.1}$$

is said to be the *parametric Stone space* with an index m , cf. definition of \mathcal{R}_m in [9, Sec. 3.1].

Main statement on the *canonical construction* of finitely axiomatizable theories can be found in [9]. Its part involved in construction of this work states the following:

Theorem 4.3. [9, Th. 3.1.1] *Effectively in a pair of integers (m, s) and a finite rich signature σ , it is possible to construct a finitely axiomatizable theory $F = \mathbb{F}\mathbb{C}(m, s, \sigma)$ of signature σ together with an effective sequence $\theta_n, n \in \mathbb{N}$, of sentences of the signature σ such that the family of extensions of F defined by*

$$F[A] = F \cup \{\theta_i \mid i \in A\} \cup \{\neg \theta_j \mid j \in \mathbb{N} \setminus A\}, \quad A \subseteq \mathbb{N}, \tag{4.2}$$

satisfies the following properties:

- (A) *for any $A \subseteq \mathbb{N}$, the theory $F[A]$ is either complete or contradictory;*
- (B) *the theory $F[A], A \subseteq \mathbb{N}$, is consistent if and only if $A \in \Omega(m)$;*
- (C) *for an arbitrary $A \in \Omega(m)$, theory $F[A]$ has a prime model if and only if the tree \mathcal{D}_s^A is atomic; moreover,*

- a prime model of the theory $F[A]$, if it exists, is strongly constructivizable if and only if the set A is computable and the family of chains $\Pi^{f^n}(\mathcal{D}_s^A)$ is computable,
 - a prime model of the theory $F[A]$, if it exists and is strongly constructivizable, has algorithmic dimension 1 if and only if the tree \mathcal{D}_s^A is computable; otherwise, the model has algorithmic dimension ω .
- (D) for an arbitrary $A \in \Omega(m)$, theory $F[A]$ is ω -stable if and only if the tree \mathcal{D}_s^A is superatomic.

Notice that, Part (A) and Part (B) of Theorem 4.3 state that the collection $F[A]$, $A \in \Omega(m)$, represents exactly the set of all possible complete extensions of the theory $F = \mathbb{F}\mathbb{C}(m, s, \sigma)$.

5 The sentence algebra of the class of prime models with infinite algorithmic dimensions having ω -stable theories

We fix a finite rich signature σ . We denote by $P(\sigma)$ the class of all prime models of signature σ , by $W(\sigma)$ the class of all models of signature σ with ω -stable theories, and by $M_{s.c.}^\omega(\sigma)$, the class of all strongly constructivizable models of signature σ having infinite algorithmic dimensions. Intersection of these classes $PW_{s.c.}^\omega = P(\sigma) \cap W(\sigma) \cap M_{s.c.}^\omega(\sigma)$ is the main object of our further study. We use a fixed Gödel numbering Φ_i , $i \in \mathbb{N}$, for the set of all sentences of signature σ , and $\varphi_i(\bar{x}_i)$, $i \in \mathbb{N}$, for the set of all formulas of signature σ .

Theorem 5.1. *The following assertions hold:*

- (a) $(\mathcal{L}(PW_{s.c.}^\omega), \gamma)$ is a Boolean Σ_1^1 -algebra, where γ is a Gödel numbering of the set of sentences of signature σ ,
- (b) computable ultrafilters of $(\mathcal{L}(PW_{s.c.}^\omega), \gamma)$ form a dense set among arbitrary ultrafilters in the algebra,
- (c) for an arbitrary numerated Boolean Σ_1^1 -algebra (\mathcal{B}, ν) whose computable ultrafilters form a dense set among arbitrary ultrafilters, there is a sentence Θ of signature σ such that $(\mathcal{B}, \nu) \cong (\mathcal{L}(\text{Th}(\text{Mod}(\Theta)) \cap PW_{s.c.}^\omega), \gamma)$.
- (d) for an arbitrary Boolean Σ_1^1 -algebra \mathcal{B} , there is a sentence Θ of signature σ such that $\mathcal{B} \cong \mathcal{L}(\text{Th}(\text{Mod}(\Theta)) \cap PW_{s.c.}^\omega)$.

PROOF. (a) By the Goncharov-Nurtazin-Harrington criterion (Theorem 3.4), a prime model \mathfrak{M} of a complete decidable theory T is strongly constructivizable if and only if the family of principal types realized in \mathfrak{M} is computable. In addition, by a corollary of the Nurtazin criterion for autostability (Theorem 3.5) the strongly constructivizable prime model \mathfrak{M} is autostable with respect to strong constructivizations if and only if the family \mathcal{F} of formulas presenting atoms in the Tarski-Lindenbaum algebras $L_n(T)$, $n \in \mathbb{N}$, is computably enumerable. Otherwise, when the set \mathcal{F} is not computably enumerable, by the Nurtazin criterion for autostability (Theorem 3.5), we have $\dim_{s.c.}(\mathfrak{M}) = \omega$. From this, we obtain that, a sentence Ψ of signature σ has a model in the class $PW_{s.c.}^\omega$ if and only if there are natural parameters m and n such that the following collection of conditions is satisfied

- (a) $W_m \cap W_n = \emptyset$ & $W_m \cup W_n = \mathbb{N}$, \forall & $\forall\exists$
 (b) $T = \{\Phi_i \mid i \in W_m\}$ is a complete theory, $\forall\exists$
 (c) $\Psi \in T$, \exists
 (d) $(\forall\varphi(\bar{x})) [\varphi(\bar{x}) \text{ consistent} \Rightarrow \exists\theta(\bar{x}) [\theta(\bar{x}) \rightarrow \varphi(\bar{x}) \theta(\bar{x}) \text{ is atomic}]]$, $\forall\exists\forall$
 (e) $(\forall q)[(\forall i \in W_q) \varphi_i(\bar{x}_i) \text{ is atomic} \Rightarrow (\exists j \notin W_q)(\varphi_j(\bar{x}_j) \text{ is atomic})]$, $\forall\exists\forall$
 (f) $(\forall \text{countable } \mathfrak{N} \in \text{Mod}(T))(\forall n < \omega)[\mathcal{L}_n(\text{Th}(\mathfrak{N}, |\mathfrak{N}|)) \text{ is superatomic}]$. \forall^1

Thus, we obtain a prefix \forall^1 for the condition as a whole. Finally, sentences Φ and Ψ are equivalent on the class $PW_{s.c.}^\omega$ if and only if $(\Phi \& \neg\Psi) \vee (\Psi \& \neg\Phi)$ does not have a model in this class. This gives the required prefix \exists^1 for (a).

(b) Let T be an arbitrary complete theory extending $\text{Th}(PW_{s.c.}^\omega)$, and Ψ be a sentence provable in T . Obviously, Ψ has a model $\mathfrak{N} \in PW_{s.c.}^\omega$. From this, we have that complete decidable theory $T' = \text{Th}(\mathfrak{N})$ presenting a computable ultrafilter in $\text{St}(\text{Th}(PW_{s.c.}^\omega))$ is found in the neighborhood Ψ of the ultrafilter T in this Stone space. This gives the required density property posed in (b).

The proof of Part (c) is given in the forthcoming text. As for Part (d), it is a simple consequence of Part (c) and Lemma 1.1 with an m -complete in Σ_1^1 set X .

We begin the proof of Part (c) of Theorem 5.1.

We use the following m -complete in the class Π_1^1 set

$$A^1 = \{n \in \mathbb{N} \mid \mathcal{L}_n \text{ is a well ordering}\}, \quad (5.1)$$

where $\mathcal{L}_n = (L_n, \nu_n)$, $n \in \mathbb{N}$, is an effective sequence of constructive linear orderings, cf. Lemma 0.1. Therefore, its complement $E^1 = \mathbb{N} \setminus A^1$ is an m -complete in Σ_1^1 set.

Given a numerated Boolean Σ_1^1 -algebra (\mathcal{B}, ν) that satisfies

$$\text{computable ultrafilters of } (\mathcal{B}, \nu) \text{ form a dense set in } \text{St}(\mathcal{B}). \quad (5.2)$$

It is possible to assume, that \mathcal{B} is a nontrivial algebra. By definition, signature operations \cup, \cap and $-$ in \mathcal{B} are presentable by computable functions on ν -numbers, while the equality is a Σ_1^1 -relation in the numeration ν , i.e., $\nu(x) = \nu(y) \Leftrightarrow H(x, y)$ is satisfied for a binary relation $H \in \Sigma_1^1$. Respectively, the inequality relation $H'(x, y) =_{dfn} (\nu(x) \neq \nu(y))$ is Π_1^1 . Therefore, there is a unary relation H^* in Π_1^1 such that, for any finite tuple of zeros and ones $\alpha = \langle \alpha_0, \alpha_1, \dots, \alpha_n \rangle$, we have

$$\nu(0)^{\alpha_0} \cap \nu(1)^{\alpha_1} \cap \dots \cap \nu(n)^{\alpha_n} \neq \mathbf{0} \Leftrightarrow \langle \alpha_0, \alpha_1, \dots, \alpha_n \rangle \in H^*, \quad H^* \in \Pi_1^1.$$

Since H^* is Π_1^1 , it is m -reducible to A^1 , i.e., there is a general computable function $f(x)$ such that for arbitrary tuples $\alpha \in 2^{<\omega}$, $\alpha = \langle \alpha_0, \dots, \alpha_n \rangle$, we have

$$\nu(0)^{\alpha_0} \cap \nu(1)^{\alpha_1} \cap \dots \cap \nu(n)^{\alpha_n} \neq \mathbf{0} \Leftrightarrow \mathcal{L}_{f(\langle \alpha_0, \alpha_1, \dots, \alpha_n \rangle)} \text{ is well ordering.} \quad (5.3)$$

Let us check that the object we built above represents some kind of a tree.

Lemma 5.2. *The following assertions are true:*

- (a) $f(\emptyset) \in A^1$,

- (b) for any α in $2^{<\omega}$, $f(\alpha) \in A^1 \Leftrightarrow f(\alpha 0) \in A^1$ or $f(\alpha 1) \in A^1$,
(c) for any α in $2^{<\omega}$, $f(\alpha) \in E^1 \Leftrightarrow f(\alpha 0) \in E^1$ and $f(\alpha 1) \in E^1$.

PROOF. Part (a) is a consequence of the fact that algebra \mathcal{B} is nontrivial. Part (b) follows from the definition of the function $f(x)$ and relation $H^*(x)$, representing an ideal in the free Boolean algebra, while (c) is a corollary from (b). \square

Now, our goal is to choose a pair (m, s) of integer parameters to be considered with the canonical construction, cf. Theorem 4.3.

Choice of m . We choose m such that $\Omega(m) = \mathcal{P}(\mathbb{N})$, cf. (4.1). For this, it is enough to get m such that $W_m = \emptyset$.

Choice of s . For this purpose, we describe a computable functional \mathcal{F} from $\mathcal{P}(\mathbb{N})$ to $\mathcal{P}(\mathbb{N})$, actually, yielding compact binary trees. For an arbitrary $A \subseteq \mathbb{N}$, witness of \mathcal{F} with input A is found via the function $f(x)$ in (5.3) by scheme

$$\mathcal{F} : A \mapsto \text{Tree}(L^*, \nu^*) \mapsto \text{Tree}(L^*, \nu^*) \oplus \mathcal{D}^* = \mathcal{F}(A) = \mathcal{D}_s^A, \quad \text{where} \quad (5.4)$$

$$(a) \quad (L^*, \nu^*) = \mathcal{L}_{f(\emptyset)} + \mathcal{L}_{f(\langle \alpha_0 \rangle)} + \dots + \mathcal{L}_{f(\langle \alpha_0, \dots, \alpha_n \rangle)} + \dots,$$

$$(b) \quad \alpha_k = \begin{cases} 1, & \text{if } k \in A, \\ 0, & \text{if } k \notin A, \end{cases} \quad \text{for all } k \in \mathbb{N};$$

(c) s is an index of the algorithm for $A \mapsto \mathcal{F}(A)$ with an oracle A .

Here, \mathcal{D}^* is a fixed superatomic computably enumerable non-computable tree with computable $\Pi^{fin}(\mathcal{D})$, cf. Lemma 4.1, while $(L, \nu) \mapsto \text{Tree}(L, \nu)$ is an effective transformation from linear orderings to binary trees, cf. Lemma 4.2. The transformation $A \mapsto \mathcal{F}(A)$, $A \subseteq \mathbb{N}$, is realized via Turing's computation by an algorithm \mathcal{M} with an oracle A . We put s to be a Gödel number of the algorithm \mathcal{M} , cf. (5.4)(c).

Choice of the pair of parameters (m, s) is finished.

Let us study main properties of the transformation $\mathcal{F} : A \mapsto \mathcal{D}_s^A$.

Lemma 5.3. *The following assertions hold:*

- (a) For any $A \subseteq \mathbb{N}$, \mathcal{D}_s^A is a tree.
(b) For any computable set $A \subseteq \mathbb{N}$, \mathcal{D}_s^A is a computable enumerable non-computable tree.
(c) For any computable set $A \subseteq \mathbb{N}$, the family $\Pi^{fin}(\mathcal{D}_s^A)$ is computable.
(d) For any computable set $A \subseteq \mathbb{N}$, tree \mathcal{D}_s^A is superatomic \Leftrightarrow linear order L^* in (5.4)(a) is well ordered.

PROOF. Given a set $A \subseteq \mathbb{N}$. Let (L^*, ν^*) be a linear order built by rule (5.4)(a) with using function $f(x)$ in (5.3). By construction, (L^*, ν^*) is a constructive linear order whenever A is a computable set.

We turn to prove separate parts of Lemma 5.3.

(a) Use Lemma 4.2 together with Lemma 4.1 describing properties of the tree \mathcal{D}^* in (5.4). Moreover, some elementary properties of the operation \oplus for trees have to be used.

(b) Suppose that A is computable. Then, (L^*, ν^*) is a constructive linear ordering. By Lemma 4.2(a), the tree $\text{Tree}(L^*, \nu^*)$ is computable, however,

by Lemma 4.1, another tree \mathcal{D}^* is a computably enumerable non-computable tree. From this, we obtain that their sum $\text{Tree}(L^*, \nu^*) \oplus \mathcal{D}^*$ is a computably enumerable non-computable tree.

(c) Suppose that A is computable. Then, (L^*, ν^*) is a constructive linear ordering. By Lemma 4.2(a), tree $\text{Tree}(L^*, \nu^*)$ is computable, thus ensuring the family $\Pi^{fin}(\text{Tree}(L^*, \nu^*))$ to be computable. By Lemma 4.1, another family $\Pi^{fin}(\mathcal{D}^*)$ is also computable. Based on elementary properties of the operation \oplus , we obtain that the family $\Pi^{fin}(\text{Tree}(L^*, \nu^*) \oplus \mathcal{D}^*)$ is computable.

(d) Suppose that A is computable. Then, (L^*, ν^*) is a constructive linear ordering. In the case when L^* is well ordering, by Lemma 4.2(b), the tree $\text{Tree}(L^*, \nu^*)$ is superatomic while by Lemma 4.1 the tree \mathcal{D}^* is also superatomic. Therefore, their sum $\text{Tree}(L^*, \nu^*) \oplus \mathcal{D}^*$ is superatomic. In the other case, when the order L^* is not well-ordering, by Lemma 4.2(b), the tree $\text{Tree}(L^*, \nu^*)$ is not superatomic, thus, the whole tree $\text{Tree}(L^*, \nu^*) \oplus \mathcal{D}^*$ is not superatomic as well.

Lemma 5.3 is proved. □

Now we turn to the final part of the proof.

In accordance with demands in Part (c) of Theorem 5.1, we have to point out a sentence Θ of a given finite rich signature σ . For this, we use the canonical construction, cf. Theorem 4.3. Let us apply Theorem 4.3 to the pair (m, s) specifying also signature σ for the target theory. As a result, we obtain an effective sequence $\theta_i, i \in \mathbb{N}$, of sentences of signature σ together with a finitely axiomatizable theory $F = \mathbb{F}\mathbb{C}(m, s, \sigma)$ of signature σ . We put Θ to be the sentence that is a conjunction of axioms of the theory $F = \mathbb{F}\mathbb{C}(m, s, \sigma)$.

Notice that, Part (A) together with Part (B) of Theorem 4.3 ensure that, for an arbitrary set $A \in \Omega(m)$ and corresponding sequence $\alpha \in 2^{<\omega}$ linked with A via the rule (5.4)(b), it is satisfied

$$F[A] \vdash \theta_k^{\alpha_k}, \text{ for all } k \in \mathbb{N}. \tag{5.5}$$

Now, we are going to check that Θ satisfies all requirements in Theorem 5.1(c).

Consider an arbitrary finite tuple of zeros and ones $\alpha = \langle \alpha_0, \dots, \alpha_k \rangle$. Let us construct an intersection of elements in \mathcal{B} by the rule

$$b_\alpha = \nu(0)^{\alpha_0} \cap \nu(1)^{\alpha_1} \cap \dots \cap \nu(k)^{\alpha_k}. \tag{5.6}$$

We also consider a conjunction of corresponding sentences $\theta_i, i \in \mathbb{N}$, by the rule

$$\beta_\alpha = \theta_0^{\alpha_0} \& \theta_1^{\alpha_1} \& \dots \& \theta_k^{\alpha_k}. \tag{5.7}$$

The main idea behind the construction is to provide the following relation:

Lemma 5.4. *For any tuple $\alpha \in 2^{<\omega}$, $b_\alpha \neq \mathbf{0}$ if and only if formula $\Theta \& \beta_\alpha$ is satisfied in a model $\mathfrak{M} \in PW_{s.c.}^\omega$.*

PROOF. First, we assume that $b_\alpha \neq \mathbf{0}$. Since computable ultrafilters form a dense set among arbitrary ultrafilters in the Boolean algebra (\mathcal{B}, ν) , cf.

(5.2), it is possible to find a sequence $\alpha^* = \langle \alpha_i \mid i < \omega \rangle$ extending α such that the set A linked with α^* by rule (5.4)(b) is computable, and

$$\nu(0)^{\alpha_0} \cap \dots \cap \nu(i)^{\alpha_i} \neq \mathbf{0}, \text{ for all } i \in \mathbb{N}. \quad (5.8)$$

By (5.3), we obtain that $\mathcal{L}_{f(\langle \alpha_0, \dots, \alpha_i \rangle)}$ is a well ordered set for all $i \in \mathbb{N}$; thus, the sequence (5.4)(a) consists of well orderings only. Therefore, their sum (L^*, ν^*) is also a well ordering. By Lemma 5.3(d), the tree \mathcal{D}_s^A is superatomic. The more, \mathcal{D}_s^A is an atomic tree. We have, $A \in \Omega(m)$ because $\Omega(m) = \mathcal{P}(\mathbb{N})$ by choice of m . By Parts (A) and (B) of Theorem 4.3, theory $F[A]$ is consistent and complete. Furthermore, by Parts (C) and (D) of Theorem 4.3, theory $F[A]$ has a prime strongly constructivizable model \mathfrak{M} of infinite algorithmic dimension whose theory is ω -stable. By virtue of (5.5), we have $F[A] \vdash \theta_i^{\alpha_i}$ for all $i \in \mathbb{N}$. From this, we obtain $\mathfrak{M} \models \beta_\alpha$. Thereby, the formula $\Theta \& \beta_\alpha$ is satisfied in the model $\mathfrak{M} \in PW_{s.c.}^\omega$.

Now, we assume that sentence $\Theta \& \beta_\alpha$ is satisfied in a model $\mathfrak{M} \in PW_{s.c.}^\omega$. Consider the set of integers

$$A = \{i \in \mathbb{N} \mid \mathfrak{M} \models \theta_i\}, \quad (5.9)$$

which is obviously computable. Build an infinite sequence $\alpha^* = \langle \alpha_i \mid i < \omega \rangle$ linked with A by the rule (5.4)(b). Since $A \in \Omega(m) = \mathcal{P}(\mathbb{N})$, theory $F[A]$ is consistent and complete. By (5.9), all axioms of the theory $F[A]$ are satisfied in the model \mathfrak{M} . By Part (C) of Theorem 4.3, the tree \mathcal{D}_s^A is superatomic, thus the tree $\text{Tree}(L^*, \nu^*)$ in (5.4) is superatomic as well. By Lemma 5.3(c), linear order \mathcal{L}^* in (5.4)(a) is well ordering. In view of (5.3), we obtain that $\mathcal{L}_{f(\langle \alpha_0, \dots, \alpha_s \rangle)}$ is a well ordered set for all $s \in \mathbb{N}$. Applying relation (5.3) to the sequence α , we finally obtain $b_\alpha \neq \mathbf{0}$.

Lemma 5.4 is proved. \square

Let us map elements $\nu(i)$, $i \in \mathbb{N}$, of Boolean algebra \mathcal{B} to sentences θ_i , $i \in \mathbb{N}$, of signature σ by the rule $\lambda^*(\nu(k)) = \theta_k$, $k \in \mathbb{N}$. By Lemma 5.4 together with Theorem 2.1, it is possible to extend this mapping up to a computable isomorphism $\lambda : (\mathcal{B}, \nu) \rightarrow (\mathcal{L}(\text{Th}(\text{Mod}(\Theta) \cap PW_{s.c.}^\omega)), \gamma)$ that is exactly what is required.

Thereby, Part (c) of Theorem 5.1 is completely proved.

Theorem 5.1 is proved. \square

6 Conclusion

Theorem 5.1 characterizes the Tarski-Lindenbaum algebra of the class of all prime strongly constructivizable models of infinite algorithmic dimensions having ω -stable theories. For comparison, the work [11] have been devoted to a similar problem for the class $P_{s.c.}$ of all strongly constructivizable prime models, while another work [13] concerned the same problem for the class $P_{s.c.}^1$ of all prime strongly constructivizable models of algorithmic dimension one.

Let us turn to a few open questions concerning statement of Theorem 5.1. We write $RM(T)$ for Morley Rank of a complete theory T . We use notation

$R^{\leq\alpha}(\sigma)$ for the class of models \mathfrak{M} of signature σ satisfying $RM(\text{Th}(\mathfrak{M})) \leq \alpha$. Similar notation $R^{\geq\alpha}(\sigma)$ is also used.

By α , we denote an arbitrary constructive ordinal.

Question 6.1. *Find a characterization of the Tarski-Lindenbaum algebra of the class of models $PW_{s.c.}^{\omega} \cap R^{\leq\alpha}$.*

Question 6.2. *Find a characterization of the Tarski-Lindenbaum algebra of the class of models $PW_{s.c.}^{\omega} \cap R^{\geq\alpha}$.*

Hypothesis 6.3. *For any constructive ordinal α , the Tarski-Lindenbaum algebra of the class of models $PW_{s.c.}^{\omega} \cap R^{\geq\alpha}$ is characterized by a statement analogous to that presented in Theorem 5.1.*

Notice that, the canonical construction is actually applicable to the cases (involved in the questions we formulated above) with an arbitrary ordinal α satisfying $15 \leq \alpha < \omega_1^{CK}$, cf. [9, Th. 3.1.1].

References

- [1] S.S. Goncharov, Yu.L. Ershov, *Constructive models*, Siberian School of Algebra Logic, Consultants Bureau, New York, 2000. Zbl 0954.03036
- [2] S.S. Goncharov, A.T. Nurtazin, *Constructive models of complete decidable theories*, Algebra Logic, **12**:2 (1973), 67–77. Zbl 0278.02038
- [3] W. Hanf, D. Myers, *Boolean sentence algebras: Isomorphism constructions*, J. Symb. Log., **48**:2 (1983), 329–338. Zbl 0511.03005
- [4] V.S. Harizanov, *Pure computable model theory*, in Ershov, Yu.L. (ed.) et al., *Handbook of recursive mathematics. Vol. 1: Recursive model theory*, Elsevier, Amsterdam, 1998, 3–114. Zbl 0952.03037
- [5] L. Harrington, *Recursively presented prime models*, J. Symb. Log., **39**:2 (1974), 305–309. Zbl 0332.02055
- [6] W. Hodges, *A shorter model theory*, Cambridge University Press, Cambridge, 1997. Zbl 0873.03036
- [7] S. Lempp, M.G. Peretyat'kin, R. Solomon, *The Lindenbaum algebra of the theory of the class of all finite models*, J. Math. Log., **2**:2 (2002), 145–225. Zbl 1033.03035
- [8] A.T. Nurtazin, *Strong and weak constructivizations and computable families*, Algebra Logic, **13**:3 (1974), 177–184. Zbl 0305.02061
- [9] M.G. Peretyat'kin, *Finitely axiomatizable theories*, Consultants Bureau, New York, 1997. Zbl 0884.03031
- [10] M.G. Peretyat'kin, *Finitely axiomatizable theories and Lindenbaum algebras of semantic classes*, in Cholak, Peter A. (ed.) et al., *Computability theory and its applications. Current trends and open problems*, Proceedings of a 1999 AMS-IMS-SIAM joint summer research conference, Boulder, CO, USA, June 13–17, 1999, Contemp. Math., **257**, 2000, 221–239. Zbl 0968.03039
- [11] M.G. Peretyat'kin, *On the Tarski-Lindenbaum algebra of the class of all strongly constructivizable prime models*, in Cooper, S. Barry (ed.) et al., *How the world computes*, Proceedings of the Turing Centenary Conference, CiE2012, Lecture Notes in Computer Science, **7318**, Springer, Berlin, 2012, 589–598. Zbl 1358.03045
- [12] M.G. Peretyat'kin, *The Tarski-Lindenbaum algebra of the class of all strongly constructivizable countable saturated models*, in Bonizzoni, Paola (ed.) et al., *The nature of computation. Logic, algorithms, applications*, Proceedings CiE 2013, Lecture Notes in Computer Science, **7921**, Springer, Berlin, 2013, 342–352. Zbl 1433.03086

- [13] M.G. Peretyat'kin, *The Tarski-Lindenbaum algebra of the class of all prime strongly constructivizable models of algorithmic dimension one*, Sib. Elektron. Mat. Izv., **17** (2020), 913–922. Zbl 1437.03128
- [14] H.J. Rogers, *Theory of recursive functions and effective computability*. McGraw-Hill Publishing Company, New York, 1967. Zbl 0183.01401
- [15] R. Sikorski, *Boolean algebras*. 3rd edition, Springer-Verlag, New York, 1969. MR0242724
- [16] R.L. Vaught, *Denumerable models of complete theories*, Infinitistic methods, Proc. Symp. Foundations Math., Warsaw, 1959, Pergamon Press, Oxford, London, New York, Paris, 1961, 303–321. Zbl 0113.24302

MIKHAIL G. PERETYATKIN
INSTITUTE OF MATHEMATICS AND MATHEMATICAL MODELING,
SHEVCHENKO 28,
050010, ALMATY, KAZAKHSTAN
Email address: peretyatkin@math.kz