

ON EXTENSIONS OF MINIMAL LOGIC WITH
LINEARITY AXIOMD.M. ANISHCHENKO, S.P. ODINTSOV *Communicated by S.V. SUDOPLATOV*

Abstract: The Dummett logic is a superintuitionistic logic obtained by adding the linearity axiom to intuitionistic logic. This is one of the first non-classical logics, whose lattice of axiomatic extensions was completely described. In this paper we investigate the logic JC obtained via adding the linearity axiom to minimal logic of Johansson. So JC is a natural paraconsistent analog of the Dummett logic. We describe the lattice of JC -extensions, prove that every element of this lattice is finitely axiomatizable, has the finite model property, and is decidable. Finally, we prove that JC has exactly two pretabular extensions.

Keywords: Dummett's logic, minimal logic, linearity axiom, lattice of extensions, algebraic semantics, j -algebra, opremum, decidability, pretabularity.

The *Dummett logic* (or *Gödel-Dummett logic*, or *logic of chains*) LC is an important superintuitionistic logic obtained by adding to intuitionistic logic Int the linearity axiom $(p \rightarrow q) \vee (q \rightarrow p)$.

In 2003 J. von Plato [20] opened for the logical community that a logical system equivalent to LC was introduced in the work by T. Skolem written in 1913 [17]. It is remarkable that this work was written in the setting of

ANISHCHENKO D.M., ODINTSOV S.P., ON EXTENSIONS OF MINIMAL LOGIC WITH LINEARITY AXIOM.

© 2024 ANISHCHENKO D.M., ODINTSOV S.P..

The work is supported by State Contracts of the Sobolev Institute of Mathematics (Project FWNF-2022-0012).

Received June, 23, 2024, Published October, ??, 2024.

Schröder’s algebraic approach, which differs essentially from the dominating Frege-Russel-Hilbert tradition. This difference von Plato describes [20, p. 153] as follows: ‘Logic is not tied to a strict formal syntax, but studies the relations between concepts in a direct way. The situation is similar to mathematics in general, where one usually studies the structures of interest, say arithmetic, without the mediation of a concrete formal syntax’. The main technical result of Skolem’s work is the discovery of a normal form of formulas. In LC every formula is equivalent to a disjunction of conjunctions or a conjunction of disjunctions of formulas of the form $(p \rightarrow q)$ or $\neg(p \rightarrow q)$, where p and q are propositional variables.

The rediscovery of LC by M. Dummett was inspired by K. Gödel [6], who constructed a sequence of matrices to prove the non-tabularity of intuitionistic logic. In 1959 M. Dummett [2] introduced LC as the extension of the intuitionistic logic via the linearity axiom and proved that LC is complete w.r.t. the family of matrices introduced in [6].

Later D. Ulrich [19] proved that every LC extension has the finite model property. Finally, M. Dunn and R. Meyer [4] have found out all proper extensions of LC . It turns out that all such extensions are tabular, which strengthens the results of Ulrich and implies the pretabularity of LC . Notice that one year later L. Maksimova [9] proved that there are exactly three pretabular superintuitionistic logics. Naturally, LC is one of such logics.

Recall that a logic is called *paraconsistent*, if it admits theories that are contradictory, but however non-trivial. Logics with trivial contradictory theories are called *explosive*. Minimal logic J of Johansson [7] can be obtained from Int via omitting the axiom *ex contradictione quodlibet* $ECQ := \neg p \rightarrow (p \rightarrow q)$, more exactly, $Int = J + ECQ$. This is a natural paraconsistent analog of intuitionistic logic.




Figure 1. The lattice \mathcal{ELC} .

The main object of our investigations is the logic JC obtained by adding the linearity axiom to J . So JC can be considered as a paraconsistent analog of the Dummett logic LC . We transfer the results by M. Dunn and R. Meyer to the logic JC . Except for description of the lattice of JC -extensions we prove that every such extension is finitely axiomatizable modulo JC , has

the finite model property, and is decidable as a consequence. Finally, we prove that the logic JC has exactly two pretabular extensions, one of which coincides with LC .

It should be noted that LC is one of the first non-classical logics, whose lattice of extensions was completely described. Earlier such description was obtained for the modal logic $S5$ [16] and for the logic RM [3], an extension of the relevant logic R with so called mingle-axiom. The lattice \mathcal{ELC} of all LC -extensions is an infinite descending chain with the least element as presented at Figure 1. The greatest element Form is the trivial logic consisting of all formulas, the second element is the classical logic CL , the logic L_n is determined by the n -element linearly ordered Heyting algebra.

M. Kracht [8] studied extensions of the constructive Nelson's logic $N3$ with strong negation. In particular, he described the lattice of extensions of the logic $N3C = N3 + \{(p \rightarrow q) \vee (q \rightarrow p)\}$, a conservative extension of LC . As we can see from Figure 2 this lattice is essentially more complicated than \mathcal{ELC} , but still admits an explicit presentation.




Figure 2. The lattice $\mathcal{EN3C}$.

Later S. Odintsov [12] described the lattice of extensions of the logic $N4C = N4 + \{(p \rightarrow q) \vee (q \rightarrow p)\}$, where $N4$ is the paraconsistent version of the constructive Nelson's logic, $N3 = N4 + \{\sim p \rightarrow (p \rightarrow q)\}$. $N4C$ also is a conservative extension of LC . It turns out that the lattice $\mathcal{EN4C}$ has much more complicated structure as compared to $\mathcal{EN3C}$ and we cannot draw a diagram presenting explicitly the structure of this lattice. As we will see the same holds for the lattice of JC -extensions. To explain the nature of such complications we say a few words on methods that are used to describe the lattice of extensions of one or another logic. Usually we deal with logics which can be characterized via varieties of algebras. Every variety is generated

by its finitely generated subdirectly irreducible algebras (**fsi**-algebras). So first we describe the **fsi**-algebras modelling the logic L we are interested in. Temporarily we denote this class of algebras as $\text{Mod}_{\text{fsi}}(L)$. Further, we consider the preorder \sqsubseteq on $\text{Mod}_{\text{fsi}}(L)$ defined as follows:

$$\mathbf{A} \sqsubseteq \mathbf{B} \Leftrightarrow LA \subseteq LB,$$

where LA denotes the logic of \mathbf{A} , i.e. the set of all formulas valid on the algebra \mathbf{A} . Naturally, this preorder is stable under isomorphisms, and factoring out the preorder $\langle \text{Mod}_{\text{fsi}}(L), \sqsubseteq \rangle$ w.r.t. the isomorphism relation we obtain the partial order $\mathcal{P}_{\text{fsi}}L$ of isomorphism types of **fsi**-algebras modelling L . It is clear that for a logic L' extending L isomorphism types of algebras from $\text{Mod}_{\text{fsi}}(L')$ form a cone (upward closed subset) of $\mathcal{P}_{\text{fsi}}L$, and that different cones correspond to different L -extensions. Usually it is possible to prove that just defined mapping from $\mathcal{E}(L)$ to the cone lattice of $\mathcal{P}_{\text{fsi}}L$ is onto, and that this is a dual isomorphism of lattices.

The simple structure of \mathcal{ELC} is explained by the fact that its partial order $\mathcal{P}_{\text{fsi}}LC$ is linear. Due to this reason the lattice of cones of this order is linear too. The order $\mathcal{P}_{\text{fsi}}N3C$ consists of two infinite chains connected one with the other. In case of $N4C^\perp$ the order $\mathcal{P}_{\text{fsi}}N4C^\perp$ has an explicit description, but it is much more complicated, so one can describe $\mathcal{EN}4C^\perp$ only as the lattice dually isomorphic to the cone lattice of $\mathcal{P}_{\text{fsi}}N4C^\perp$. The same holds for the logic JC investigated in this article.

Our paper is structured as follows. Section 1 is devoted to preliminary remarks. In Section 2 we describe the partial order $\mathcal{P}_{\text{fsi}}JC$ and cones of this order. Finally, in Section 3 we establish that \mathcal{EJC} and the cone lattice of $\mathcal{P}_{\text{fsi}}JC$ are dually isomorphic, prove that all elements of \mathcal{EJC} are finitely axiomatizable and decidable, and find out pretabular JC -extensions.

1 Preliminaries

We consider logics defined in the propositional language $\mathcal{L} = \{\vee, \wedge, \rightarrow, \perp\}$ containing the binary connectives of disjunction, conjunction, implication, and the absurdity constant. Formulas of this language (\mathcal{L} -formulas) are constructed in a standard way from \perp and propositional variables occurring to a fixed countable set Prop with the help of logical connectives. The negation connective we consider as an abbreviation, $\neg\varphi := \varphi \rightarrow \perp$. The set of all \mathcal{L} -formulas is denoted as $\text{Form}_{\mathcal{L}}$. We write $\varphi(p_1, \dots, p_n)$ for $\varphi \in \text{Form}_{\mathcal{L}}$ in case all propositional variables occurring φ are among p_1, \dots, p_n .

By a *logic* L we mean a set of formulas ($L \subseteq \text{Form}_{\mathcal{L}}$) closed under the rules of substitution (SUB) and *modus ponens* (MP):

$$\text{(SUB)} \frac{\varphi(p_1, \dots, p_n)}{\varphi(\psi_1, \dots, \psi_n)}, \quad \text{(MP)} \frac{\varphi \quad \varphi \rightarrow \psi}{\psi},$$

where $\varphi, \psi, \psi_1, \dots, \psi_n \in \text{Form}_{\mathcal{L}}$.

For a logic L , we denote by \mathcal{EL} the set of all logics containing L . Set-theoretical intersection \cap and the operation $+$, where $L_1 + L_2$ is the least

logic containing $L_1 \cup L_2$, turn $\mathcal{E}L$ to a complete lattice. This follows from the fact that the intersection of any family of logics is a logic too. We call $\mathcal{E}L$ the *lattice of L-extensions*. If $X \subseteq \text{Form}_{\mathcal{L}}$, then $L + X$ denotes the least logic in $\mathcal{E}L$ containing X .

Minimal logic or *Johansson's logic* J is the least logic containing the following formulas:

$$\begin{array}{ll} J1. p \rightarrow (q \rightarrow p); & J2. (p \wedge q) \rightarrow p; \\ J3. (p \rightarrow (q \rightarrow r)) \rightarrow ((p \rightarrow q) \rightarrow (p \rightarrow r)); & J4. (p \wedge q) \rightarrow q; \\ J5. (p \rightarrow q) \rightarrow ((p \rightarrow r) \rightarrow (p \rightarrow (q \wedge r))); & J6. p \rightarrow (p \vee q); \\ J7. (p \rightarrow r) \rightarrow ((q \rightarrow r) \rightarrow ((p \vee q) \rightarrow r)); & J8. q \rightarrow (p \vee q). \end{array}$$

Now *intuitionistic logic* Int can be defined as $Int = J + \{\perp \rightarrow p\}$, and *classical logic* as $CL = Int + \{p \vee \neg p\}$.

Put:

$$JC = J + \{(p \rightarrow q) \vee (q \rightarrow p)\}, \quad LC = Int + \{(p \rightarrow q) \vee (q \rightarrow p)\}.$$

Logic LC is the *logic of chains* or the *Dummett logic*. The main object of our investigations is logic JC , a natural paraconsistent analog of the Dummett logic.

Now we consider an algebraic semantics for logics from $\mathcal{E}J$. Algebras we denote by boldface uppercase letters, the universes of algebras by corresponding italic uppercase letters.

An *implicative lattice* is an algebra $\mathbf{A} = \langle A, \wedge, \vee, \rightarrow \rangle$ such that its reduct $\langle A, \wedge, \vee \rangle$ is a lattice and the operation \rightarrow satisfies the following equivalence for all $a, b, c \in A$:

$$c \leq_{\mathbf{A}} a \rightarrow b \Leftrightarrow c \wedge a \leq_{\mathbf{A}} b,$$

where $\leq_{\mathbf{A}}$ denotes the lattice ordering of $\langle A, \wedge, \vee \rangle$. Recall that $a \leq_{\mathbf{A}} b$ iff $a \wedge b = a$.

It is easy to see that for every $a \in A$, the implication $a \rightarrow a$ is the greatest element of \mathbf{A} w.r.t. $\leq_{\mathbf{A}}$. In what follows the greatest element of \mathbf{A} we denote as $1_{\mathbf{A}}$. The next property of implication readily follows from its definition:

$$a \rightarrow b = 1_{\mathbf{A}} \quad \text{iff} \quad a \leq_{\mathbf{A}} b \quad \text{for any } a, b \in A \quad (1)$$

It is well known that every implicative lattice is distributive and that the class \mathcal{V}_{imp} of all implicative lattices is a variety, i.e. this class can be axiomatized by identities (see, e.g., [14])

An algebra $\mathbf{A} = \langle A, \wedge, \vee, \rightarrow, \perp \rangle$ is called a *j-algebra* if its \perp -free reduct $\langle A, \wedge, \vee, \rightarrow \rangle$ is an implicative lattice.

A *j-algebra* $\mathbf{A} = \langle A, \wedge, \vee, \rightarrow, \perp \rangle$ is called a *Heyting algebra* (*negative algebra*) if \perp is the least (greatest) element of \mathbf{A} with respect to $\leq_{\mathbf{A}}$.

Obviously, the class \mathcal{V}_j of all *j-algebras*, the class \mathcal{V}_H of all Heyting algebras, and the class \mathcal{V}_{neg} of all negative algebras are varieties too.

Let \mathbf{A} be a *j-algebra*. Every homomorphism $v : \text{Form}_{\mathcal{L}} \rightarrow \mathbf{A}$ from the algebra of formulas into \mathbf{A} is called an *A-valuation*. For every formula φ , we say that φ is *true at an A-valuation* v and write $\mathbf{A}, v \models \varphi$ if $v(\varphi) = 1_{\mathbf{A}}$.

We say that φ is *true on \mathbf{A}* and write $\mathbf{A} \models \varphi$ if $\mathbf{A}, v \models \varphi$ holds for every \mathbf{A} -valuation v .

For a set Γ of formulas, we write $\mathbf{A} \models \Gamma$ and say that \mathbf{A} is a *model for Γ* if $\mathbf{A} \models \varphi$ for all $\varphi \in \Gamma$.

Let \mathcal{K} be a class of j -algebras. We write $\mathcal{K} \models \varphi$ if $\mathbf{A} \models \varphi$ for all $\mathbf{A} \in \mathcal{K}$, and $\mathcal{K} \models \Gamma$ if $\mathcal{K} \models \varphi$ for all $\varphi \in \Gamma$.

For an algebra \mathbf{A} , the lattice of its congruences is denoted by $\text{Con}(\mathbf{A})$. For a variety \mathcal{V} , we denote by $\text{Sub}(\mathcal{V})$ the lattice of its subvarieties. In both cases, the lattice ordering coincides with the inclusion relation, see [1] for details. Moreover, it is known that the variety \mathcal{V}_j of all j -algebras is congruence-distributive, i.e. the lattice $\text{Con}(\mathbf{A})$ is distributive for every $\mathbf{A} \in \mathcal{V}_j$.

Congruences of a j -algebra \mathbf{A} are in one-to-one correspondence with filters on \mathbf{A} . Recall that a non-empty subset $F \subseteq A$ is a *filter on a j -algebra \mathbf{A}* if for every $a, b \in A$: (i) $a \in F$ and $b \in F$ imply $a \wedge b \in F$; (ii) $a \leq_{\mathbf{A}} b$ and $a \in F$ imply $b \in F$. The lattice of all filters on \mathbf{A} we denote $\mathcal{F}(\mathbf{A})$. For every $\theta \in \text{Con}(\mathbf{A})$ and every $F \in \mathcal{F}(\mathbf{A})$ we put:

$$F_\theta := \{a \in A \mid (a, 1_{\mathbf{A}}) \in \theta\}, \quad \theta_F := \{(a, b) \mid a \rightarrow b, b \rightarrow a \in F\}.$$

The mappings $\theta \mapsto F_\theta$ and $F \mapsto \theta_F$ are mutually inverse isomorphisms between $\text{Con}(\mathbf{A})$ and $\mathcal{F}(\mathbf{A})$.

For a logic $L \in \mathcal{EJ}$ and for a variety $\mathcal{V} \in \text{Sub}(\mathcal{V}_j)$, we put

$$V(L) := \{\mathbf{A} \mid \mathbf{A} \in \mathcal{V}_j \text{ and } \mathbf{A} \models L\}; \quad L\mathcal{V} := \{\varphi \mid \mathcal{V} \models \varphi\}.$$

For an arbitrary class of j -algebras \mathcal{K} , we also put $L\mathcal{K} = \{\varphi \mid \mathcal{K} \models \varphi\}$. If a logic $L \in \mathcal{EJ}$ is such that $L = L\mathcal{K}$ we say that L is *complete w.r.t. \mathcal{K}* . For a j -algebra \mathbf{A} , we write $L\mathbf{A}$ instead of $L\{\mathbf{A}\}$ and call $L\mathbf{A}$ the *logic of \mathbf{A}* .

Since the fact that a formula φ is true on a j -algebra \mathbf{A} is equivalent to the validity of the identity $\varphi = p \rightarrow p$ on \mathbf{A} , we have $V(L) \in \text{Sub}(\mathcal{V}_j)$ for every $L \in \mathcal{EJ}$. It is known also that $L\mathcal{V} \in \mathcal{EJ}$ for every $\mathcal{V} \in \text{Sub}(\mathcal{V}_j)$. Moreover, the following statement holds, see, e.g. [13].

Theorem 1. *The mappings $L \mapsto V(L)$ and $\mathcal{V} \mapsto L\mathcal{V}$ are mutually inverse dual isomorphisms¹ between lattices \mathcal{EJ} and $\text{Sub}(\mathcal{V}_j)$. In particular, for every $L_0 \in \mathcal{EJ}$ and $\mathcal{V}_0 \in \mathcal{V}_j$ we have*

$$L(V(L_0)) = L_0 \quad \text{and} \quad V(L(\mathcal{V}_0)) = \mathcal{V}_0.$$

This statement implies, in particular, that $\varphi \in J$ iff φ is true on every j -algebra. Since $\perp \rightarrow p$ is true on \mathbf{A} iff \perp is the least element of \mathbf{A} w.r.t. $\leq_{\mathbf{A}}$ we obtain that $V(\text{Int}) = \mathcal{V}_H$ and $L(\mathcal{V}_H) = \text{Int}$. Naturally, the restriction of $L \mapsto V(L)$ to \mathcal{EInt} is a dual isomorphism between \mathcal{EInt} and $\text{Sub}(\mathcal{V}_H)$. Similarly, the restriction of $L \mapsto V(L)$ to \mathcal{EL}_0 for every $L_0 \in \mathcal{EJ}$ is a dual isomorphism between \mathcal{EL}_0 and $\text{Sub}(V(L_0))$.

Let us recall an important fact from universal algebra. Every variety \mathcal{V} of algebras is generated by the family \mathcal{V}_{fsi} of its finitely generated subdirectly

¹By a dual isomorphism between lattices \mathbf{A} and \mathbf{B} we mean an isomorphism between \mathbf{A} and \mathbf{B}^{op} , where \mathbf{B}^{op} is the lattice with the same universe as \mathbf{B} and reversed order.

irreducible algebras (fsi-algebras). For a definition of subdirectly irreducible algebra the reader may consult [1]. For our goals it will be enough to give a characterization of subdirectly irreducible j -algebras.

An element $*_{\mathbf{A}}$ of a j -algebra \mathbf{A} is called an *opremum* of \mathbf{A} if $*_{\mathbf{A}} \neq 1_{\mathbf{A}}$ and $b \leq *_{\mathbf{A}}$ for every $b \in \mathbf{A}$ different from $1_{\mathbf{A}}$.

Theorem 2. (see, e.g. [15]) *A j -algebra \mathbf{A} is subdirectly irreducible if and only if there is an opremum $*_{\mathbf{A}}$ in \mathbf{A} .*

We define the preorder on $(\mathcal{V}_j)_{\text{fsi}}$ as follows: $\mathbf{A} \sqsubseteq \mathbf{B}$ iff $L\mathbf{A} \subseteq L\mathbf{B}$ for $\mathbf{A}, \mathbf{B} \in (\mathcal{V}_j)_{\text{fsi}}$. The well known Jonsson Theorem (see [1]) provides a useful characterization of this preorder. This theorem says that in a congruence distributive variety, all subdirectly irreducible elements of the subvariety generated by the class \mathcal{K} of algebras belong to $\text{HSUp}(\mathcal{K})$.² If an algebra \mathbf{A} is finite, then $\text{HSUp}(\{\mathbf{A}\}) = \text{HS}(\{\mathbf{A}\})$. The variety of j -algebras is congruence distributive, therefore, for any finite $\mathbf{A}, \mathbf{B} \in (\mathcal{V}_j)_{\text{fsi}}$ we have

$$\mathbf{A} \sqsubseteq \mathbf{B} \Leftrightarrow L\mathbf{A} \subseteq L\mathbf{B} \Leftrightarrow \mathbf{B} \in \text{HS}(\{\mathbf{A}\}).$$

2 Isomorphism types of algebras from $(V(JC))_{\text{fsi}}$

As it follows from Theorem 1 the lattice $\mathcal{E}JC$ is dually isomorphic to the lattice $\text{Sub}(V(JC))$. So it would be enough to describe the lattice of subvarieties of the variety $V(JC)$ consisting of all JC -models.

First we describe fsi-algebras from the variety $V(JC)$. The next statement is an immediate generalization of the characterization of fsi-algebras from $V(LC)$ obtained in [4].

Proposition 1. *Let \mathbf{A} be a j -algebra. Then $\mathbf{A} \in (V(JC))_{\text{fsi}}$ if and only if \mathbf{A} is finite and linearly ordered.*

Proof. First we assume that \mathbf{A} is finite and linearly ordered j -algebra. Since \mathbf{A} is finite it is finitely generated. Finiteness of \mathbf{A} implies that every non-unit element lies under some maximal non-unit element. The linearity of \mathbf{A} implies that there is only one maximal non-unit element, which is an opremum of \mathbf{A} . Thus, \mathbf{A} is an fsi-algebra. It follows immediately from (1) that $(p \rightarrow q) \vee (q \rightarrow p)$ is true on a linearly ordered j -algebra. Consequently, $\mathbf{A} \in (V(JC))_{\text{fsi}}$.

Now we assume that $\mathbf{A} \in (V(JC))_{\text{fsi}}$. If $a \not\leq_{\mathbf{A}} b$ and $b \not\leq_{\mathbf{A}} a$ for some $a, b \in \mathbf{A}$, then we have $(a \rightarrow b) \vee (b \rightarrow a) \leq *_{\mathbf{A}}$ by (1), which conflicts with $\mathbf{A} \models (p \rightarrow q) \vee (q \rightarrow p)$. Thus, \mathbf{A} is linearly ordered.

Let X be a finite set generating \mathbf{A} . We just proved that \mathbf{A} is linearly ordered. Consequently, $a \vee b, a \wedge b \in \{a, b\}$ for $a, b \in \mathbf{A}$. Further, notice that $a \rightarrow b \in \{1_{\mathbf{A}}, b\}$. If $a \leq_{\mathbf{A}} b$, then $a \rightarrow b = 1_{\mathbf{A}}$ by (1). If $b \not\leq_{\mathbf{A}} a$, then

²For a class of algebras \mathcal{K} , the class of all homomorphic images of algebras from \mathcal{K} is denoted as $\text{H}(\mathcal{K})$, the class of all subalgebras of algebras from \mathcal{K} as $\text{S}(\mathcal{K})$, finally, the class of all ultraproducts of algebras from \mathcal{K} is denoted as $\text{Up}(\mathcal{K})$.

$a \rightarrow b = b$. Indeed, $a \wedge b = b$ and $a \wedge c \not\geq b$ for $c \not\geq b$. Thus, except for generators \mathbf{A} may contain only \perp and $1_{\mathbf{A}}$. Consequently, \mathbf{A} is finite. □

In the last proof we established the following.

Corollary 1. *Let \mathbf{A} be a linearly ordered j -algebra. The implication act on \mathbf{A} as follows:*

$$a \rightarrow b = \begin{cases} 1_{\mathbf{A}}, & a \leq_{\mathbf{A}} b; \\ b, & \text{otherwise.} \end{cases} \tag{2}$$

Moreover, if $\{\perp_{\mathbf{A}}, 1_{\mathbf{A}}\} \subseteq X \subseteq |\mathbf{A}|$, then X is the universe of a subalgebra of \mathbf{A} .

From Proposition 1 we conclude that up to isomorphism every element of $(V(JC))_{\text{fsi}}$ is determined by the number of its elements and by the interpretation of \perp in this finite linear order. Thus, we may code isomorphism types of algebras from $(V(JC))_{\text{fsi}}$ by pairs of positive natural numbers. We say that $\mathbf{A} \in (V(JC))_{\text{fsi}}$ is of type (m, n) , and write $\text{tp}(\mathbf{A}) = (m, n)$, if m is the number of elements greater or equal to \perp , and n is the number of elements of \mathbf{A} , which are smaller or equal to \perp .

Now we are ready to describe the preordering \sqsubseteq on $(V(JC))_{\text{fsi}}$. Recall that $\mathbf{A} \sqsubseteq \mathbf{B}$ iff $L_{\mathbf{A}} \subseteq L_{\mathbf{B}}$ iff $\mathbf{B} \in \text{HS}(\{\mathbf{A}\})$ for fsi-algebras \mathbf{A} and \mathbf{B} .

Proposition 2. *Let $\mathbf{A}, \mathbf{B} \in (V(JC))_{\text{fsi}}$, $\text{tp}(\mathbf{A}) = (m, n)$, and $\text{tp}(\mathbf{B}) = (k, l)$. The following equivalence holds:*

$$\mathbf{A} \sqsubseteq \mathbf{B} \iff k \leq m \text{ and } l \leq n.$$

Proof. If $\mathbf{B} \in \text{HS}(\{\mathbf{A}\})$, then obviously $k \leq m$ and $l \leq n$.

Assume that $k \leq m$ and $l \leq n$ and prove the inverse implication. Assume additionally that $1_{\mathbf{A}} \neq \perp_{\mathbf{A}}$ and $1_{\mathbf{B}} \neq \perp_{\mathbf{B}}$. By Corollary 1 arbitrary subset X of $|\mathbf{A}|$ containing $1_{\mathbf{A}}$ and $\perp_{\mathbf{A}}$ is the universe of a subalgebra of \mathbf{A} . This immediately implies that \mathbf{A} has a subalgebra isomorphic to \mathbf{B} , i.e., $\mathbf{B} \in \text{S}(\{\mathbf{A}\})$. The same holds if $1_{\mathbf{A}} = \perp_{\mathbf{A}}$ and $1_{\mathbf{B}} = \perp_{\mathbf{B}}$.

It remains to consider the case $1_{\mathbf{A}} \neq \perp_{\mathbf{A}}$ and $1_{\mathbf{B}} = \perp_{\mathbf{B}}$, when $\mathbf{B} \notin \text{S}(\{\mathbf{A}\})$. Put $F = \{a \mid \perp_{\mathbf{A}} \leq_{\mathbf{A}} a\}$ and $\mathbf{A}' := \mathbf{A}/F$. It follows easily from (2) that $[a]_F = F$ for $a \in F$ and $[a]_F = \{a\}$ for $a \notin F$. Thus, $\text{tp}(\mathbf{A}') = (1, n)$ and by the above argument $\mathbf{B} \in \text{S}(\mathbf{A}')$. From $\mathbf{A}' \in \text{H}(\mathbf{A})$ we conclude $\mathbf{A} \sqsubseteq \mathbf{B}$. □

Put $\mathbb{N}^+ = \{1, 2, 3, \dots\}$ and $\mathcal{P} = \langle (\mathbb{N}^+)^2, \preceq \rangle$, where $(m, n) \preceq (k, l)$ iff $k \leq m$ and $l \leq n$. It is clear that \preceq is a partial order on the set of pairs of positive natural numbers. The structure of this ordering is presented on the following diagramm.




Figure 3.

Recall that a subset $X \subseteq (\mathbb{N}^+)^2$ is a *cone* (*upward closed set*) w.r.t. \preceq if for every $a \in X$ and $b \in (\mathbb{N}^+)^2$ such that $a \preceq b$ we have $b \in X$. The set of all cones w.r.t. \preceq form a lattice with the inclusion relation as its lattice order. We denote this lattice $\text{Up}(\mathcal{P})$.

For $L \in \mathcal{EJC}$ put $\mathbf{tp}(L) := \{\mathbf{tp}(\mathbf{A}) \mid \mathbf{A} \in (V(L))_{\mathbf{fsi}}\}$. From the just obtained characterization of \sqsubseteq we infer the next fact.

Corollary 2. *For every $L \in \mathcal{EJC}$ we have $\mathbf{tp}(L) \in \text{Up}(\mathcal{P})$. Moreover, the mapping $\mathbf{tp} : \mathcal{EJC} \rightarrow \text{Up}(\mathcal{P})$ is one-to-one and order reversing.*

Proof. From Proposition 2 we obtain $\mathbf{tp}(L) \in \text{Up}(\mathcal{P})$ for $L \in \mathcal{EJC}$. According to Theorem 1 the mapping $L \mapsto V(L)$ is order reversing, consequently, \mathbf{tp} is order reversing too. Finally, if $L_1 \neq L_2$, then $(V(L_1))_{\mathbf{fsi}} \neq (V(L_2))_{\mathbf{fsi}}$. This fact implies that \mathbf{tp} is one-to-one. \square

In other words, for every JC -extension L the set of types of \mathbf{fsi} -algebras modelling L form a cone w.r.t. \preceq . In what follows we prove that every cone from $\text{Up}(\mathcal{P})$ can be realized as $\mathbf{tp}(L)$ for some JC -extension L . But first we obtain more explicit description of cones from $\text{Up}(\mathcal{P})$.

For $(m, n) \in (\mathbb{N}^+)^2$ we denote by $(m, n) \uparrow$ the principal cone generated by (m, n) , i.e.,

$$(m, n) \uparrow := \{(k, l) \mid (m, n) \preceq (k, l)\} = \{(k, l) \mid k \leq m, l \leq n\}.$$

Every principal cone $(m, n) \uparrow$ is finite as it follows from its definition. For a proper cone $X \in \text{Up}(\mathcal{P})$, i.e., such cone that $\emptyset \neq X \neq (\mathbb{N}^+)^2$, we define two natural numbers $r(X)$ and $l(X)$ as follows: $r(X)$ is the greatest natural number such that $(m, r(X)) \in X$ for all $m \in \mathbb{N}^+$, and there is $n \in \mathbb{N}^+$ such that $(n, r(X) + 1) \notin X$. The number $l(X)$ is defined is a dual way, this is the greatest natural number such that $(l(X), m) \in X$ for all $m \in \mathbb{N}^+$, and there is $n \in \mathbb{N}^+$ such that $(l(X) + 1, n) \notin X$.

It can be easily seen that natural numbers $l(X)$ and $r(X)$ are well defined for every proper cone X . Moreover, $r(X) = 0$ iff $(m, 1) \notin X$ for some m , and $l(X) = 0$ iff $(1, n) \notin X$ for some n .

Now we associate with every proper cone $X \in \text{Up}(\mathcal{P})$ two subcones $W^r(X)$ and $W^l(X)$:

$$W^r(X) := \{(m, n) \mid n \leq r(X)\} \quad \text{and} \quad W^l(X) := \{(m, n) \mid m \leq l(X)\}.$$

Obviously, $W^r(X) = \emptyset$ iff $r(X) = 0$, and $W^l(X) = \emptyset$ iff $l(X) = 0$. Moreover, if one of these subcones is not empty, then it is infinite.

Now we are ready to describe cones from $\text{Up}(\mathcal{P})$.

Proposition 3. *Every proper cone $X \in \text{Up}(\mathcal{P})$ is the union of cones $W^r(X)$, $W^l(X)$, and of finite number of principal cones.*

Proof. By definition of $r(X)$ and $l(X)$ there are m and n such that

$$\{(m, r(X) + 1), (l(X) + 1, n)\} \cap X = \emptyset.$$

Notice that in this case

$$X \setminus (W^l(X) \cup W^r(X)) \subseteq (m, n) \uparrow.$$

Indeed, assume that (s, t) is such that $l(X) + 1 \leq s$, $r(X) + 1 \leq t$, and $(s, t) \notin (m, n) \uparrow$. In this case either $m < s$, or $n < t$. If $m < s$, then $(s, t) \preceq (m, r(X) + 1)$, and so $(s, t) \notin X$. If $n < t$, then again $(s, t) \notin X$ in view of $(s, t) \preceq (l(X) + 1, n)$.

Let $(u_1, v_1), \dots, (u_n, v_n)$ be all elements of $X \cap (m, n) \uparrow$ minimal w.r.t. \preceq . Then

$$X = W^l(X) \cup W^r(X) \cup (u_1, v_1) \uparrow \cup \dots \cup (u_n, v_n) \uparrow.$$

□

3 The dual isomorphism of lattices \mathcal{EJC} and $\text{Up}(\mathcal{P})$

We just noticed that the mapping tp assigning to a JC -extension L the cone of isomorphism types of algebras from $(V(L))_{\text{fsi}}$ is one-to-one and order reversing. Now we prove that tp is a dual lattice isomorphism between \mathcal{EJC} and $\text{Up}(\mathcal{P})$. To this end given a cone $X \in \text{Up}(\mathcal{P})$ we provide an axiomatization for a logic $L \in \mathcal{EJC}$ such that $\text{tp}(L) = X$.

For $n \in \mathbb{N}^+$, we define the following formula with propositional variables from the list p_1, \dots, p_n :

$$\tau_n := p_1 \vee (p_1 \rightarrow p_2) \vee \dots \vee (p_{n-1} \rightarrow p_n).$$

Lemma 1. *Let \mathbf{A} be a finite linearly ordered j -algebra. Then the following equivalence holds:*

$$\mathbf{A} \models \tau_n \Leftrightarrow |A| \leq n.$$

Proof. Let v be an arbitrary \mathbf{A} -valuation. It follows easily from (2) that $\mathbf{A}, v \not\models \tau_n$ iff

$$1_{\mathbf{A}} \not\geq v(p_1) \not\geq v(p_2) \not\geq \dots \not\geq v(p_n).$$

Thus, $\mathbf{A} \not\models \tau_n$ iff A contains at least $n + 1$ different elements. □

Let \mathbf{A} be a j -algebra. Put

$$A^\perp := \{b \in A \mid b \geq_{\mathbf{A}} \perp_{\mathbf{A}}\} \quad \text{and} \quad A_\perp := \{b \in A \mid b \leq_{\mathbf{A}} \perp_{\mathbf{A}}\}.$$

For $a, b \in A_\perp$, we put $a \rightarrow_\perp b = (a \rightarrow b) \wedge \perp_{\mathbf{A}}$.

It is obvious that the set A^\perp is closed under the operations of \mathbf{A} and that $\mathbf{A}^\perp = \langle A^\perp, \wedge, \vee, \rightarrow, \perp \rangle$ is a Heyting algebra. The set A_\perp is closed under $\vee, \wedge, \rightarrow_\perp$, and $\mathbf{A}_\perp = \langle A_\perp, \wedge, \vee, \rightarrow_\perp, \perp \rangle$ is a negative algebra.

Moreover, it is known [13, Prop. 4.2.3] that for every formula $\phi(p_1, \dots, p_n)$ the following holds:

$$\mathbf{A} \models \phi(p_1 \vee \perp, \dots, p_n \vee \perp) \Leftrightarrow \mathbf{A}^\perp \models \phi(p_1, \dots, p_n);$$

$$\mathbf{A} \models \perp \rightarrow \phi(p_1, \dots, p_n) \Leftrightarrow \mathbf{A}_\perp \models \phi(p_1, \dots, p_n).$$

We will use these equivalences and Lemma 1 to establish the following facts.

Proposition 4. *Let $m, n \in \mathbb{N}^+$, $\mathbf{A} \in (V(JC))_{\text{fsi}}$, and $X \in \text{Up}(\mathcal{P})$. Then the following equivalences hold:*

- (1) $\text{tp}(\mathbf{A}) \in (m, n) \uparrow$ iff $\mathbf{A} \models \tau_m(p_1 \vee \perp, \dots, p_m \vee \perp)$ and $\mathbf{A} \models \perp \rightarrow \tau_n$;
- (2) $\text{tp}(\mathbf{A}) \in W^r(X)$ iff $\mathbf{A} \models \perp \rightarrow \tau_{r(X)}$;
- (3) $\text{tp}(\mathbf{A}) \in W^l(X)$ iff $\mathbf{A} \models \tau_{l(X)}(p_1 \vee \perp, \dots, p_{l(X)} \vee \perp)$.

Proof. To prove Item 1 recall that $\text{tp}(\mathbf{A}) \in (m, n) \uparrow$ means exactly that $|A^\perp| \leq m$ and $|A_\perp| \leq n$. The latter is equivalent by Lemma 1 to $\mathbf{A}^\perp \models \tau_m$ and $\mathbf{A}_\perp \models \tau_n$, i.e., $\mathbf{A} \models \tau_m(p_1 \vee \perp, \dots, p_m \vee \perp)$ and $\mathbf{A} \models \perp \rightarrow \tau_n$.

Items 2 and 3 follow in a similar way from the observations that $\text{tp}(\mathbf{A}) \in W^r(X)$ iff $|A_\perp| \leq r(X)$, and that $\text{tp}(\mathbf{A}) \in W^l(X)$ iff $|A^\perp| \leq l(X)$. \square

Theorem 3. *The mapping $\text{tp} : \mathcal{EJC} \rightarrow \text{Up}(\mathcal{P})$ is a dual lattice isomorphism between \mathcal{EJC} and $\text{Up}(\mathcal{P})$.*

Proof. Since an order isomorphism of two lattices is also a lattice isomorphism and we know that $\text{tp} : \mathcal{EJC} \rightarrow \text{Up}(\mathcal{P})$ is a one-to-one and order reversing mapping, it will be enough to prove that tp maps \mathcal{EJC} onto $\text{Up}(\mathcal{P})$.

Take an arbitrary cone $X \in \text{Up}(\mathcal{P})$. According to Proposition 3 it can be represented as

$$X = W^l(X) \cup W^r(X) \cup (u_1, v_1) \uparrow \cup \dots \cup (u_n, v_n) \uparrow,$$

for some $u_1, v_1, \dots, u_n, v_n \in \mathbb{N}^+$. Put

$$L^{l(X)} = JC + \{\tau_{l(X)}(p_1 \vee \perp, \dots, p_{l(X)} \vee \perp)\}, \quad L^{r(X)} = JC + \{\perp \rightarrow \tau_{r(X)}\},$$

$$L^{u_i, v_i} = JC + \{\tau_{u_i}(p_1 \vee \perp, \dots, p_{u_i} \vee \perp), \perp \rightarrow \tau_{v_i}\}, \quad i = 1, \dots, n.$$

Proposition 4 implies that

$$\text{tp}(L^{l(X)}) = W^l(X), \quad \text{tp}(L^{r(X)}) = W^r(X), \quad \text{tp}(L^{u_i, v_i}) = (u_i, v_i) \uparrow, \quad i = 1, \dots, n.$$

Since $L \mapsto V(L)$, $L \in \mathcal{EJC}$, is a dual lattice isomorphism, we have $V(L_1 \cap L_2) = V(L_1) + V(L_2)$ and

$$(V(L_1))_{\text{fsi}} \cup (V(L_2))_{\text{fsi}} \subseteq (V(L_1 \cap L_2))_{\text{fsi}}.$$

To prove the inverse inclusion, and so the equality

$$(V(L_1 \cap L_2))_{\text{fsi}} = (V(L_1))_{\text{fsi}} \cup (V(L_2))_{\text{fsi}} \tag{3}$$

for logics finitely axiomatizable modulo JC we need the following results.

S. Miura [10] noticed that an intersection of two superintuitionistic logics can be axiomatized by repeatedless disjunctions of their axioms. This result can be easily generalized to extensions of minimal logic [13, Prop. 2.1.5]. More exactly, a *repeatedless disjunction* of formulas $\varphi(p_1, \dots, p_n)$ and $\psi(p_1, \dots, p_m)$ is a formula $\varphi \underline{\vee} \psi := \varphi(p_1, \dots, p_n) \vee \psi(p_{n+1}, \dots, p_{n+m})$. Further, if $L \in \mathcal{EJ}$, $L_1 = L + \{\varphi_i \mid i \in I_1\}$, and $L_2 = L + \{\psi_j \mid j \in I_2\}$, then

$$L_1 \cap L_2 = L + \{\varphi_i \underline{\vee} \psi_j \mid i \in I_1, j \in I_2\}.$$

In particular, if $L_1 = JC + \{\varphi(p_1, \dots, p_n)\}$ and $L_2 = JC + \{\psi(p_1, \dots, p_m)\}$, then $L_1 \cap L_2 = JC + \{\varphi \underline{\vee} \psi\}$. Take a linearly ordered fsi-algebra \mathbf{A} such that $\mathbf{A} \notin (V(L_1))_{\text{fsi}} \cup (V(L_2))_{\text{fsi}}$, i.e.

$$v_1(\varphi(p_1, \dots, p_n)) \leq *_{\mathbf{A}} \text{ and } v_2(\psi(p_1, \dots, p_m)) \leq *_{\mathbf{A}}$$

for some \mathbf{A} -valuations v_1 and v_2 . Let $v(p_i) = v_1(p_i)$ for $i = 1, \dots, n$ and $v(p_{n+i}) = v_2(p_i)$ for $i = 1, \dots, m$. Then we have $v(\varphi \underline{\vee} \psi) \leq *_{\mathbf{A}}$, and so $\mathbf{A} \notin (V(L_1 \cap L_2))_{\text{fsi}}$. We have thus proved (3) for finitely axiomatizable JC -extensions.

The just obtained result implies that for an arbitrary cone $X \in \text{Up}(\mathcal{P})$ we have $\text{tp}(L) = X$, where

$$L = L^{l(X)} \cap L^{r(X)} \cap L^{u_1, v_1} \cap \dots \cap L^{u_n, v_n}. \tag{4}$$

□

Proposition 5. *Every logic from \mathcal{EJC} is finitely axiomatizable, has the finite model property, and is decidable.*

Proof. Every logic $L \in \mathcal{EJC}$ is finitely axiomatizable modulo JC according to (4).

Every $L \in \mathcal{EJC}$ has the finite model property due to the fact that L is complete w.r.t. the family of its fsi-models, which are all finite according to Proposition 1.

Finally, for every $L \in \mathcal{EJC}$ its set of fsi-models is decidable. Indeed, we noticed that up to isomorphism fsi-models of JC can be coded by pairs of natural numbers. According to Corollary 2 and Proposition 3 for every $L \in \mathcal{EJC}$, the family $(V(L))_{\text{fsi}}$ is coded by a finite union of sets having one of the following forms

$$\{(k, l) \mid k \leq n\}, \quad \{(k, l) \mid l \leq m\}, \quad \{(k, l) \mid k \leq r, l \leq s\}.$$

Thus, every $L \in \mathcal{EJC}$ is complete w.r.t. to a decidable set of finite algebras. Consequently, L is decidable according to the well known Harrop Theorem saying that a logic with enumerable set of theorems, which is complete w.r.t. to an enumerable set of finite models, is decidable.

□

Finally, we describe pretabular JC -extensions. Recall that a logic $L \in \mathcal{E}J$ is *tabular* if $L = L\mathbf{A}$ for some finite j -algebra \mathbf{A} . We call a logic $L \in \mathcal{E}J$ *pretabular* if L is not tabular, but every its proper extension is.

Proposition 6. *The lattice $\mathcal{E}JC$ contains exactly two pretabular logics: the Dummett logic LC and $NegC = JC + \{\perp\}$.*

Proof. If $L = L\mathbf{A}$, then $V(L) = V(\{\mathbf{A}\})$. If additionally we assume that \mathbf{A} is finite, then $(V(L))_{\text{fsi}} \subseteq \text{HS}(\{\mathbf{A}\})$ by Jonsson's lemma. The finiteness of \mathbf{A} implies that $(V(L))_{\text{fsi}}$ is finite up to isomorphism. Thus, tabular logics from $\mathcal{E}JC$ are such logics L that the cone $\text{tp}(L)$ is finite. Respectively, pretabular logics from $\mathcal{E}JC$ are such logics L that the cone $\text{tp}(L)$ is infinite, but every proper subcone of $\text{tp}(L)$ is finite. It is enough to have a look at Figure 3 to understand that there are only two such cones in $\text{Up}(\mathcal{P})$:

$$C_1 = \{(n, 1) \mid n \in \mathbb{N}^+\} \quad \text{and} \quad C_2 = \{(1, n) \mid n \in \mathbb{N}^+\}.$$

The cone C_1 contains exactly isomorphism types of algebras that have no elements lying strictly under \perp , i.e. $\text{tp}(\mathbf{A}) \in C_1$ if and only if \mathbf{A} is a finite linear Heyting algebra. Thus, $\text{tp}(LC) = C_1$. On the other hand, $\text{tp}(\mathbf{A}) \in C_2$ iff \perp is the greatest element of \mathbf{A} . Clearly, such algebras are distinguished in $(V(JC))_{\text{fsi}}$ with the help of formula \perp , and we have $\text{tp}(NegC) = C_2$. \square

References

- [1] S. Burris, H.P. Sankappanavar, *A course in universal algebra*, Graduate Texts in Math., **78**, Springer, New York etc., 1981. Zbl 0478.08001
- [2] M. Dummett, *A propositional calculus with a denumerable matrix*, J. Symb. Log., **24** (1959), 97–106. Zbl 0089.24307
- [3] J.M. Dunn, *Algebraic completeness results for R-mingle and its extensions*, J. Symb. Log., **35** (1970), 1–13. Zbl 0231.02024
- [4] J.M. Dunn, R.K. Meyer, *Algebraic completeness results for Dummett's LC and its extensions*, Z. Math. Logik Grundlagen Math., **17:2** (1971), 225–230. Zbl 0252.02018
- [5] J.K. Feibleman, R.M. Smullyan, R.L. Vaught, *Meeting of the association for symbolic logic*, J. Symb. Logic, **35:2** (1970), 352–363. ‘
- [6] K. Gödel, *Zum intuitionistischen Aussagenkalkül*, S. Feferman et al. (eds.) Kurt Gödel Collected Works, Vol.I: Publications 1929–1936, Oxford University Press, Oxford, 2001. (Zbl 59.0866.02 1933)
- [7] I. Johansson, *Der Minimalkalkül, ein reduzierter intuitionistischer Formalismus*, Compos. Math., **4** (1936), 119–136. Zbl 0015.24102
- [8] M. Kracht, *On extensions of intermediate logics by strong negation*. J. Philos. Log., **27:1** (1998), 49–73. Zbl 0929.03035
- [9] L.L. Maksimova, *Pretabular superintuitionistic logics*, Algebra Logic, **11:5** (1972), 558–570. Zbl 0265.02018
- [10] S. Miura, *A remark on the intersection of two logics*, Nagoja Math. J., **26** (1966), 167–171. Zbl 0202.29502
- [11] S.P. Odintsov, *The lattice of extensions of minimal logic*, Sib. Adv. Math., **17:2** (2007), 112–143. Zbl 1249.03033

- [12] S.P. Odintsov, *On extensions of Nelson's logic satisfying Dummett's axiom*, Sib. Math. J., **48**:1 (2007), 112–125. Zbl 1164.03313
- [13] S. Odintsov, *Constructive negations and paraconsistency*, Springer, Dordrecht, 2008. Zbl 1161.03014
- [14] H. Rasiowa, *An algebraic approach to non-classical logics*, Studies in Logic and the Foundations of Mathematics, **78**, North-Holland Publishing Company, Amsterdam-London, PWN, Warszawa, 1974. Zbl 0299.02069
- [15] W. Rautenberg, *Klassische und nichtclassische Aussagenlogik*, Friedr. Vieweg & Sohn, Braunschweig/Wiesbaden, 1979. Zbl 0424.03007
- [16] S.J. Scroggs, *Extensions of the Lewis system S5*, J. Symb. Log., **16** (1951), 112–120. Zbl 0043.00804
- [17] T. Skolem, *Om konstitutionen av den identiske kalkuls grupper*, Proceedings of the 3rd Scandinavian Mathematical Congress, Kristiania, 1913, 149–163. English translation in [18].
- [18] T. Skolem, *Selected Works in Logic*, ed. by J.E. Fenstad. Universitetsforlaget, Oslo etc., 1970. Zbl 0228.02001
- [19] D. Ulrich, *Decidability results for some classes of propositional calculi*, J. Symb. Log., **35** (1970), 353–354, as a part of [5].
- [20] J. von Plato, *Skolem's discovery of Gödel-Dummett logic*, Stud. Log., **73**:1 (2003), 153–157. Zbl 1015.03005

DMITRIY MIKHAILOVICH ANISHCHENKO
NOVOSIBIRSK STATE UNIVERSITY,
UL. PIROGOVA, 1,
630090, NOVOSIBIRSK, RUSSIA
Email address: d.anishchenko@ng.nsu.ru

SERGEI PAVLOVICH ODINTSOV
SOBOLEV INSTITUTE OF MATHEMATICS,
PR. KOPTYUGA, 4,
630090, NOVOSIBIRSK, RUSSIA
Email address: odintsov@math.nsc.ru