

THE CRITICAL GROUP OF THE CONE OVER A SANDWICH GRAPH

L.A. GRUNVALD 

Communicated by ???

Abstract: We study the critical group of the cone over a sandwich graph. It is shown that this group can be described as the cokernel of a discrete Laplace operator. Furthermore, we prove that its exact structure can be determined by making use of a recurrence relation derived from the Laplace operator. This relation involves symmetric Laurent polynomials and provides a more efficient way to compute the group. The proposed method offers a powerful tool for analyzing invariants of graphs with complex combinatorial structure.

Keywords: critical group, circulant graph, discrete Laplacian, Smith normal form.

1 Introduction

An important problem in algebraic graph theory is describing the structure of the *critical group* of a graph. This notion is closely related to discrete analogues of classical Laplace operator. The discrete Laplacian naturally

arises as a combinatorial version of its continuous counterpart, providing a useful tool for studying various graph invariants.

The significance of the critical group is not limited to the classical fact that its order equals the number of spanning trees in a graph [1]. Often, it also captures deeper structural properties, reflecting analogies with discrete versions of classical concepts from the theory of Riemann surfaces [1, 2], as well as connections to statistical physics [3, 4].

Although determining the exact structure of critical groups remains an open problem for many classes of graphs, certain families possess specific features that make them easier to analyze. Such special cases allow step-by-step progress toward understanding the general problem.

In Section 2, we define the main object of the study, namely the *cone graph* \hat{G} constructed over a given sandwich graph G . The cone graph \hat{G} is obtained by adding one new vertex connected to every vertex of G , thus forming a discrete analogue of a geometric cone with base G .

Several well-known graphs fit naturally into this sandwich graph construction. Important examples include the I -graph $I(n, k, l)$, the generalized Petersen graph $GP(n, k)$ (see for example [5, 6]), and the generalized prism $Pr(n)$, which is also known as the cobordism of two circulant graphs [7].

Sandwich graphs form a special case of a more general construction known as *circulant foliation* [8, 9], a structure relevant to applications such as network design [10], quantum computing [11], chemistry [12], and number theory [13].

Section 2 also summarizes key known results. In particular, for cone graphs \hat{G} , there exists another naturally related group whose order equals the number of rooted spanning forests in the base graph G [14]. Understanding the structure of the discrete Laplacian for sandwich graphs thus gives essential insights into the critical group of \hat{G} and allows us to develop computational methods that improve numerical efficiency, as shown in Section 3.

We believe the results in this paper are new and may help to develop a more general understanding of circulant foliations graphs [8, 9].

2 Cone over a sandwich graph

Let us consider a classical path graph with vertices v_1, v_2, \dots, v_m , characterized by the conditions $\deg(v_1) = \deg(v_m) = 1$ and $\deg(v_i) = 2$ for $i = 2, \dots, m-1$, where $\deg(\cdot)$ denotes the vertex degree.

For given integers n and m we introduce the concept of a *sandwich graph* G , whose vertex set is defined as

$$V(G) = \{(k, v_i) \mid k = 0, 1, \dots, n-1, i = 1, 2, \dots, m\}.$$

In this graph, for each fixed i , vertices (k, v_i) together with integer parameters (called *jumps*) $s_{i,1}, s_{i,2}, \dots, s_{i,k_i} \in \mathbb{Z}$ satisfying

$$1 \leq s_{i,1} < s_{i,2} < \dots < s_{i,k_i} < \frac{n}{2},$$

form the circulant graph

$$G_i = C_n(s_{i,1}, s_{i,2}, \dots, s_{i,k_i}), \quad (1)$$

where each vertex (k, v_i) is adjacent to

$$(k \pm s_{i,1}, v_i), (k \pm s_{i,2}, v_i), \dots, (k \pm s_{i,k_i}, v_i) \pmod n.$$

From this point of view, the circulant graphs play the role of *layers* in the sandwich graph. Furthermore, if we fix k , then there is an edge connecting the vertices (k, v_i) and (k, v_{i+1}) . Thus, for the sandwich graph G defined above, we will say that G has m layers G_1, \dots, G_m , each on n vertices.




FIG. 1. Sandwich graph G with $m = 3$ layers. From top to bottom, the layers are circulant graphs: $G_1 = C_6(1)$, $G_2 = C_6(2)$ and $G_3 = C_6(1, 2)$.

An important property of circulant graphs relevant to our study is their natural connection with Laurent polynomials [15]. Specifically, each circulant layer corresponds to a Laurent polynomial

$$P_i(z) = 2k_i - \sum_{j=1}^{k_i} (z^{s_{i,j}} + z^{-s_{i,j}}), \quad i = 1, \dots, m, \quad j = 0, \dots, n-1, \quad (2)$$

which is *symmetric*, i. e. $P_i(z) = P_i(\frac{1}{z})$, and *bimonic*, meaning that the leading and trailing coefficients are equal to ± 1 .

Finally, we construct the cone graph \hat{G} over the sandwich graph G by adding one new vertex adjacent to all vertices of G . This provides a discrete analogue of a geometric cone with the graph G as its *base*.

Critical group of cone over a graph and Laplacian. Formally, for a connected graph G on n vertices, the critical group $K(G)$ is the torsion part of the abelian group $\text{coker}(L(G))$ (see for example [1]), where $L(G)$ is the discrete Laplacian operator viewed as a homomorphism $\mathbb{Z}^n \rightarrow \mathbb{Z}^n$.

The discrete Laplacian is a discrete analogue of the classical Laplace operator, and it is represented by $n \times n$ matrix

$$L(G) = D(G) - A(G),$$

where $D(G)$ is the diagonal degree matrix (whose diagonal entries are the degrees of vertices), and $A(G)$ is the adjacency matrix of G .

A fundamental property of the critical group is that its order equals the number of spanning trees of the graph [1]. The following Theorem 1 in [14] generalizes this classical fact, introducing a new group whose order equals the number of rooted spanning forests.

Theorem 1. *Let G be a connected graph on n vertices. Then the critical group of the cone \hat{G} over the graph G is isomorphic to the cokernel of the linear operator $I_n + L(G)$, where $L(G)$ is the discrete Laplacian of G , and I_n is the $n \times n$ identity matrix. More precisely,*

$$\text{coker}(L(\hat{G})) \cong \text{coker}(I_n + L(G)).$$

We refer to this group as the *forest group of a graph* $F(G)$.

This result connects the critical group of the cone graph \hat{G} with the operator $I_n + L(G)$, also known as the discrete Helmholtz operator [16]. It provides a discrete analogue of classical analytic concepts, thus extending continuous analytical methods into the discrete setting of combinatorial graph theory.

Theorem 1 shows that the cokernel of the Laplacian of the cone graph is a torsion group, as the operator $I_n + L(G)$ is nonsingular (has trivial kernel). Therefore, we have a decomposition

$$\text{coker}(L(\hat{G})) \cong \mathbb{Z}_{d_1} \oplus \mathbb{Z}_{d_2} \oplus \cdots \oplus \mathbb{Z}_{d_n},$$

where (d_1, d_2, \dots, d_n) are *invariant factors* obtained from the Smith normal form of $L(\hat{G})$, satisfying $d_i \mid d_{i+1}$ for all $1 \leq i \leq n - 1$. Consequently, we obtain the following representation of the critical group

$$K(\hat{G}) \cong \mathbb{Z}_{d_1} \oplus \mathbb{Z}_{d_2} \oplus \cdots \oplus \mathbb{Z}_{d_n}. \quad (3)$$

From a computational perspective, determining the Smith normal form of the discrete Laplacian $L(\hat{G})$ becomes challenging as the size of the graph grows, since the dimension of $L(\hat{G})$ increases with the number of vertices. However, the special algebraic structure of circulant graphs, encoded in their associated Laurent polynomials (2), provides a useful way to simplify the problem. The following Theorem 2 in [14], originally formulated for arbitrary connected graphs, offers a significant computational simplification.

Theorem 2. *Let $P(z)$ be a Laurent polynomial with integer coefficients, and let \mathcal{A} be the companion matrix associated with $P(z)$. Then*

$$\text{coker}(L(G)) \cong \text{coker}(\mathcal{A}^n - I).$$

This result relies on a classical construction from the theory of polynomials, namely the *companion matrix*; see [17] for further details.

Theorem 2 allows us to replace the computation of the Smith normal form of the potentially large Laplacian matrix with the simpler computation of a fixed-size matrix $\mathcal{A}^n - I$, thus significantly reducing computational complexity.

3 Critical group of the cone over a sandwich graph

Let $A(G)$ denote the adjacency matrix of a sandwich graph G , and $D(X)$ denote the diagonal matrix with given independent variables $X = (x_v)$, indexed by the vertices. We define the matrix

$$L(G, X) = D(X) - A(G)$$

as the *generalized Laplacian matrix* of the graph G .

For convenience, let us denote the Laurent polynomials $L_i(z)$ defined for graphs (1) by

$$L_i = L_i(z) = P(z) + d_i + 1, \quad i = 1, \dots, m, \tag{4}$$

where $d_i = \deg(v_i)$ is the degree of vertex v_i in the path graph H .

Theorem 3. *Let \hat{G} be the cone over the sandwich graph G with m circulant layers G_1, \dots, G_m , each on n vertices. Then the forest group $F(\hat{G})$ is isomorphic to the cokernel of the linear operator $\mathcal{A}^n - I$, where \mathcal{A} is the companion matrix of the bimonic and symmetric Laurent polynomial $D_m(z)$, which satisfies the recurrence relation*

$$D_m(z) = L_m(z)D_{m-1}(z) - D_{m-2}(z), \tag{5}$$

with initial conditions

$$D_{-1}(z) = 0, \quad D_0(z) = 1, \quad D_1(z) = L_1(z), \quad D_2(z) = L_1(z)L_2(z) - 1. \tag{6}$$

Proof. Let G be a sandwich graph. Consider its generalized Laplacian matrix

$$L(G, X) = \begin{pmatrix} L_1 & -1 & 0 & \cdots & 0 & 0 \\ -1 & L_2 & -1 & \cdots & 0 & 0 \\ 0 & -1 & L_3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & L_{m-1} & -1 \\ 0 & 0 & 0 & \cdots & -1 & L_m \end{pmatrix}, \tag{7}$$

where the entries $X = (L_i)$ are symmetric and bimonic Laurent polynomials with integer coefficients, satisfying $L_i(z) = L_i(\frac{1}{z})$.

As in the Introduction, let \hat{G} denote the cone over G , and let $L(G)$ denote the discrete Laplacian of its base graph. According to Theorem 1, the forest group of G is given by $F(G) = \text{coker}(I + L(G))$. Using the standard

description of the cokernel as an abelian group of relations among n -periodic sequences, we obtain

$$\langle x_i^1, x_i^2, \dots, x_i^m \mid i \in \mathbb{Z} \mid \\ L_1 x_j^1 - x_j^2 = 0, \quad L_2 x_j^2 - x_j^1 - x_j^3 = 0, \quad \dots, \quad L_m x_j^m - x_j^{m-1} = 0, \\ x_{j+n}^1 - x_j^1 = 0, \quad x_{j+n}^2 - x_j^2 = 0, \quad \dots, \quad x_{j+n}^m - x_j^m = 0, \quad j \in \mathbb{Z} \rangle \quad (8)$$

This is an abelian group generated by m infinite families of generators with $2m$ relations: m linear relations and m periodicity conditions.

By sequentially eliminating variables x_j^i for $i = 2, \dots, m$, the system reduces to a single linear relation involving only the variables x_j^1 . In particular, for $m = 2$ we have

$$(L_1 L_2 - 1)x_j^1 = 0,$$

and for $m = 3$, we obtain

$$(L_3 L_2 L_1 - L_3 - L_1)x_j^1 = 0.$$

In general, since each x_j^i is expressed linearly through x_j^{i-1} and x_j^{i+1} , the system collapses to a single linear recurrence relation

$$D_m x_j^1 = 0,$$

where the polynomial D_m satisfies the recurrence

$$D_m = L_m D_{m-1} - D_{m-2},$$

with with initial conditions

$$D_{-1} = 0, \quad D_0 = 1, \quad D_1 = L_1, \quad D_2 = L_1 L_2 - 1.$$

Using the definition of $L_i(z)$ given in (4), it is straightforward to verify that the Laurent polynomial $D_m(z)$ satisfies the symmetry condition and is bimonic.

Since every x_j^i can be expressed through x_j^1 , the periodicity condition for all variables is equivalent to the periodicity condition on x_j^1 . Thus, the group (8) has the following presentation

$$\text{coker}(I + L(H_n)) \cong \langle x_j^1 \mid D_m(z) x_j^1 = 0, \quad x_{j+n}^1 - x_j^1 = 0, \quad j \in \mathbb{Z} \rangle.$$

Applying Theorem 2, we immediately obtain the statement of the theorem. \square

One can recognize that the recurrence relation (5) with initial conditions (6), coincides with the classical formula for the determinant of the tridiagonal matrix (7).

If the sandwich graph has at least one layer with zero jumps, i.e., if for some $i \in \{1, \dots, m\}$ we have $s_{i,1} = s_{i,2} = \dots = s_{i,k_i} = 0$, then the polynomial $L_i(z)$ reduces to $L_i(z) = d_i + 1$. In this case, the Laurent polynomial $D_m(z)$ is not bimonic, and the statement of Theorem 3 no longer holds.

4 Examples

For clarity and simplicity, we consider the special case of a sandwich graph whose layers are identical and represented by simple cycles, i.e., $G_i = C_n(s_{i,1})$ with $s_{i,1} = 1$ for all $i = 1, \dots, m$. Here, m denotes the number of layers, and n is the number of vertices in each layer.

For example, if $m = 2$, then, by Theorem 3, the forest group $F(\hat{G})$ is isomorphic to the cokernel of the linear operator $\mathcal{A}^n - I$, where \mathcal{A} is the companion matrix corresponding to the Laurent polynomial $D_2(z) = 17 + \frac{1}{z^2} - \frac{8}{z} - 8z + z^2$. Explicitly, the matrix \mathcal{A} has the following form

$$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 8 & -17 & 8 \end{pmatrix}. \tag{9}$$

For $m = 2$, the invariant factors were computed via Theorem 3 from the Smith normal form of $\mathcal{A}^n - I$, where \mathcal{A} is defined in (9); see Table 1 for $n = 3, \dots, 10$.

Note for Table 1. Each cell in Table 1 lists the invariant factors of the corresponding forest group (3) vertically. Each factor is shown explicitly in prime factorized form for clarity. For example, if $n = 3$ we have $F(G) \cong \mathbb{Z}_{24} \oplus \mathbb{Z}_{72}$.

n	3	4	5	6	7	8	9	10
$m = 2$	$2^3 \cdot 3$ $2^3 \cdot 3^2$	$3 \cdot 5$ $3 \cdot 5$ $3 \cdot 5 \cdot 7$	$11 \cdot 29$ $3 \cdot 11 \cdot 29$	2^3 2^3 $2^3 \cdot 3$ $2^3 \cdot 3^2 \cdot 5 \cdot 7$	$29 \cdot 139$ $3 \cdot 29 \cdot 139$	$3 \cdot 5 \cdot 7$ $3 \cdot 5 \cdot 7 \cdot 23$ $3 \cdot 5 \cdot 7 \cdot 23 \cdot 47$	$2^3 \cdot 3^2 \cdot 19 \cdot 37$ $2^3 \cdot 3^3 \cdot 19 \cdot 37$	$5 \cdot 11 \cdot 19 \cdot 29$ $3 \cdot 5^2 \cdot 7 \cdot 11 \cdot 19 \cdot 29$

TABLE 1. Invariant factors of the forest group for a sandwich graph with two layers.

Additional numerical results are shown in Table 2. This data was collected in an effort to identify patterns by varying both the number of vertices per layer (n) and the total number of layers (m). However, the general pattern remains an open question.

Note for Table 2. The structure of Table 2 is similar to Table 1, except that n now indexes rows and m indexes columns.

$m \mapsto$ $n \downarrow$	3	4	5	6
3	2^2 $2^2 \cdot 5 \cdot 7$ $2^3 \cdot 5 \cdot 7$	$2^4 \cdot 3 \cdot 17$ $2^4 \cdot 3^2 \cdot 7 \cdot 17$	$2^2 \cdot 29 \cdot 41$ $2^2 \cdot 5 \cdot 11 \cdot 29 \cdot 41$	$2^3 \cdot 3$ $2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11$ $2^4 \cdot 3^3 \cdot 5 \cdot 7 \cdot 11$
4	2^2 $2^3 \cdot 3$ $2^3 \cdot 3^2$ $2^6 \cdot 3^2 \cdot 5$	$3 \cdot 5$ $3 \cdot 7 \cdot 23$ $3 \cdot 5 \cdot 7 \cdot 23 \cdot 47$	$3 \cdot 5 \cdot 11 \cdot 19 \cdot 29$ $3 \cdot 5^2 \cdot 11 \cdot 19 \cdot 29 \cdot 41$	$2^2 \cdot 3$ $2^4 \cdot 3^2 \cdot 5$ $2^4 \cdot 3^2 \cdot 5 \cdot 11$ $2^7 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 23$
5	$11 \cdot 19 \cdot 41$ $2^3 \cdot 11 \cdot 19 \cdot 41$	$11 \cdot 29 \cdot 719$ $3 \cdot 7 \cdot 11 \cdot 29 \cdot 719$	$2^2 \cdot 3 \cdot 5$ $2^2 \cdot 3 \cdot 5^2 \cdot 11$ $2^2 \cdot 3 \cdot 5^2 \cdot 11 \cdot 31$ $2^2 \cdot 3 \cdot 5^2 \cdot 11 \cdot 31$	$11 \cdot 19 \cdot 29 \cdot 41 \cdot 661$ $2^4 \cdot 3^2 \cdot 11 \cdot 19 \cdot 29 \cdot 41 \cdot 661$
6	5 $2^2 \cdot 5$ $2^3 \cdot 5$ $2^3 \cdot 5 \cdot 7$ $2^3 \cdot 3 \cdot 5 \cdot 7$ $2^3 \cdot 3^2 \cdot 5 \cdot 7$	$2^4 \cdot 7$ $2^4 \cdot 7$ $2^4 \cdot 3 \cdot 7 \cdot 17$ $2^4 \cdot 3^2 \cdot 5 \cdot 7 \cdot 17 \cdot 47$	11 $11 \cdot 41$ $2^3 \cdot 5 \cdot 11 \cdot 19 \cdot 29 \cdot 41$ $2^3 \cdot 5^2 \cdot 11 \cdot 19 \cdot 29 \cdot 41$	2^3 $2^3 \cdot 3 \cdot 5$ $2^3 \cdot 3 \cdot 5$ $2^3 \cdot 3^2 \cdot 5 \cdot 7$ $2^3 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 13$ $2^6 \cdot 3^3 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 23$
7	$29 \cdot 71 \cdot 239$ $2^3 \cdot 29 \cdot 71 \cdot 239$	$7 \cdot 29 \cdot 139 \cdot 2113$ $3 \cdot 7^2 \cdot 29 \cdot 139 \cdot 2113$	$29 \cdot 71 \cdot 421 \cdot 8329$ $5 \cdot 11 \cdot 29 \cdot 71 \cdot 421 \cdot 8329$	$29 \cdot 71 \cdot 139 \cdot 239 \cdot 12781$ $2^4 \cdot 3^2 \cdot 29 \cdot 71 \cdot 139 \cdot 239 \cdot 12781$ $2^4 \cdot 3^3 \cdot 5 \cdot 7 \cdot 11$
8	$2^2 \cdot 7$ $2^4 \cdot 3 \cdot 7$ $2^4 \cdot 3^2 \cdot 7 \cdot 17$ $2^7 \cdot 3^2 \cdot 5 \cdot 7 \cdot 17$	5 5 5 $5 \cdot 7 \cdot 23$ $3 \cdot 5 \cdot 7 \cdot 23$ $3 \cdot 5 \cdot 7 \cdot 23 \cdot 47$	3 3 3 3 $3 \cdot 5 \cdot 7 \cdot 11 \cdot 19 \cdot 29 \cdot 31 \cdot 719$ $3 \cdot 5^2 \cdot 7 \cdot 11 \cdot 19 \cdot 29 \cdot 31 \cdot 41 \cdot 719$	2 2 2^2 $2^2 \cdot 7$ $2^3 \cdot 3 \cdot 7$ $2^5 \cdot 3^2 \cdot 5 \cdot 7 \cdot 23$ $2^5 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 23 \cdot 47$ $2^8 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 17 \cdot 23 \cdot 47$
9	2^2 $2^2 \cdot 5 \cdot 7 \cdot 19 \cdot 53 \cdot 199$ $2^3 \cdot 5 \cdot 7 \cdot 19 \cdot 53 \cdot 199$	$2^4 \cdot 3^2 \cdot 17 \cdot 19 \cdot 37 \cdot 8929$ $2^4 \cdot 3^3 \cdot 7 \cdot 17 \cdot 19 \cdot 37 \cdot 8929$	19 19 $2^2 \cdot 19 \cdot 29 \cdot 41 \cdot 109 \cdot 179 \cdot 251$ $2^2 \cdot 5 \cdot 11 \cdot 19 \cdot 29 \cdot 41 \cdot 109 \cdot 179 \cdot 251$	3 3 3 $2^3 \cdot 3^2$ $2^3 \cdot 3^3 \cdot 5 \cdot 7 \cdot 11 \cdot 19 \cdot 37 \cdot 53 \cdot 199 \cdot 829$ $2^4 \cdot 3^4 \cdot 5 \cdot 7 \cdot 11 \cdot 19 \cdot 37 \cdot 53 \cdot 199 \cdot 829$
10	11 11 $2^2 \cdot 5 \cdot 11 \cdot 19 \cdot 29 \cdot 41$ $2^5 \cdot 3 \cdot 5^2 \cdot 11 \cdot 19 \cdot 29 \cdot 41$	3 3 3 $3 \cdot 5 \cdot 7 \cdot 11 \cdot 19 \cdot 29 \cdot 31 \cdot 719$ $3 \cdot 5^2 \cdot 7 \cdot 11 \cdot 19 \cdot 29 \cdot 31 \cdot 41 \cdot 719$	5 $2 \cdot 5$ $2 \cdot 5^2$ $2^1 \cdot 5^2 \cdot 11$ $2^4 \cdot 3 \cdot 5^2 \cdot 11$ $2^4 \cdot 3 \cdot 5^2 \cdot 11$ $2^4 \cdot 3 \cdot 5^2 \cdot 11 \cdot 31$ $2^4 \cdot 3 \cdot 5^3 \cdot 11 \cdot 31 \cdot 41$	$11 \cdot 19 \cdot 29$ $11 \cdot 19 \cdot 29$ $2^3 \cdot 3 \cdot 5 \cdot 11 \cdot 19 \cdot 29 \cdot 41 \cdot 241 \cdot 661$ $2^6 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 19 \cdot 29 \cdot 241 \cdot 661$

TABLE 2. Invariant factors of the forest group for a sandwich graph with $m = 3, 4, 5, 6$ layers.

References

- [1] R. Bacher, P. de la Harpe, T. Nagnibeda, *The lattice of integral flows and the lattice of integral cuts on a finite graph*, Bull. Soc. Math. France., **125**:2 (1997), 167–198.
- [2] M. Baker, S. Norine, *Riemann-Roch and Abel-Jacobi theory on a finite graph*, Adv. Math., **215**:2 (2007), 766–788.
- [3] P. Bak, C. Tang, K. Wiesenfeld, *Self-organized criticality*, Physical Review A., **38** (1988), 364–374.
- [4] D. Dhar, *Self-organized critical state of sandpile automaton models*, Phys. Rev. Lett., **64** (1990), 1613–1616.
- [5] N. L. Biggs, *Three remarkable graphs*, Canad. J. Math., **15**:2 (1973), 397–411.
- [6] M. Boben, T. Pisanski, A. Zitnik, *I-graphs and the corresponding configurations*, J. Comb. Designs., **13**:6 (2005), 406–424.
- [7] N. V. Abrosimov, G. A. Baigonakova, L. A. Grunwald, I. A. Mednykh, *Counting rooted spanning forests in cobordism of two circulant graphs*, Sib. Elektron. Mat. Izv., **17** (2020), 814–823.
- [8] L. A. Grunwald, Young Soo Kwon, I. A. Mednykh, *Counting rooted spanning forests for circulant foliation over a graph*, Tohoku Math. J., **74**:4 (2022), 535–548.
- [9] Y. S. Kwon, A. D. Mednykh, I. A. Mednykh, *Complexity of the circulant foliation over a graph*, J. Algebr. Comb., **53** (2020), 115–129.
- [10] E. A. Monakhova, *A Survey on Undirected Circulant Graphs*, Discret. Math. Algorithms Appl., **4**:1 (2012)
- [11] N. Saxena, S. Severini, I. Shparlinski, *Parameters of integral circulant graphs and periodic quantum dynamics*, International Journal of Quantum Information, **5** (2007), 417–430.
- [12] M. Imran, A. Baig, S.A.U.H. Bokhary, I. Javaid, *On the metric dimension of circulant graphs*, Appl. Math. Lett., **25** (2012), 320–325.
- [13] J. Sander, T. Sander, *Recent developments on the edge between number theory and graph theory. From Arithmetic to Zeta-Functions: Number Theory in Memory of Wolfgang Schwarz* (2016), 405–425.
- [14] L. A. Grunwald, I. A. Mednykh, *On the Jacobian group of a cone over a circulant graph*, Mat. Zametki Sev. Vost. federal. Univ., **28**:2 (2021), 88–101.
- [15] P. J. Davis, *Circulant Matrices: 2nd ed.*, Chelsea Publishing, New York, 1994.
- [16] M. B. van Gijzen, Y.A. Erlangga, C. Vuik, *Spectral Analysis of the Discrete Helmholtz Operator Preconditioned with a Shifted Laplacian*, SIAM J. Sci. Comput., **29**:5 (2007), 1942–B“1958.
- [17] R. A. Horn, C. R. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1985.

LILIYA ALEXANDROVNA GRUNWALD
 SOBOLEV INSTITUTE OF MATHEMATICS,
 PR. KOPTYUGA, 4,
 630090, NOVOSIBIRSK, RUSSIA
 Email address: mathmanlily@gmail.com