

# The sum of orders of elements in nonabelian groups of odd order

Alexander Buturlakin\*, Anastasiya Tereshchenko†

## Abstract

Denote by  $\psi(G)$  the sum of the orders of the elements of a finite group  $G$ . We obtain an exact upper bound for  $\psi(G)$  on the set of nonabelian groups of given odd order  $n$  in terms of the minimal prime divisor of  $n$ . We also describe the finite groups on which this bound is achieved.

**Keywords:** orders of elements, solvable groups.

**MSC2010:** 20D10, 20D60.

## 1. Introduction

Denote by  $\psi(G)$  the sum of the orders of the elements of a finite group  $G$ . In [1] the authors show that the maximum of  $\psi(G)$  on the set of groups of the given order  $n$  is attained at a cyclic group. This maximum is strict, that is, if  $H$  is a non-cyclic group of order  $n$ , then  $\psi(H) < \psi(G)$ . Using the function  $\psi$ , it is possible to formulate sufficient conditions for a finite group to be cyclic, abelian, nilpotent, solvable, or supersolvable [2]. The paper [2], in particular, gives an upper bound for the values of  $\psi(G)$  on the set of nonabelian groups  $G$  of even order. The paper [3] contains a similar result for the non-cyclic  $q^*$ -groups. A finite group  $G$  is a  $q^*$ -group if  $q$  is the smallest prime divisor of the order of  $G$ . The main result of [3] states that if  $G$  is a finite non-cyclic  $q^*$ -group of order  $n$  and  $C_n$  is a cyclic group of order  $n$ , then

$$\psi(G) \leq \frac{(q^3 - q + 1)(q + 1)}{q^5 + 1} \psi(C_n).$$

For a positive integer  $m$  and a prime  $q$ , put

$$M_{q^{m+1}} = \langle a, b | a^{q^m} = b^q = 1, a^b = a^{1+q^{m-1}} \rangle. \quad (1)$$

Here we prove the following statement.

**Theorem.** *Let  $q$  be an odd prime and let  $G$  be a nonabelian  $q^*$ -group of order  $n$ . Then*

$$\psi(G) \leq \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1} \psi(C_n)$$

*with the equality if and only if  $G$  is a direct product of  $M_{q^3}$  and a cyclic group of order coprime to  $q$ .*

---

\*buturlakin@math.nsc.ru; Sobolev Institute of Mathematics, 4 Acad. Koptyug avenue, 630090 Novosibirsk, Russia.

†a.f.tereshchenko@gmail.com; Siberian State University of Telecommunications and Informatics, 86 Kirov str., 630102 Novosibirsk, Russia.

## 2. Preliminaries

The following lemma contains some useful facts about the function  $\psi$ .

**Lemma 1.** (1) [4, Lemma 2.2] *If  $A, B$  are finite groups of coprime orders, then  $\psi(A \times B) = \psi(A)\psi(B)$ .*

(2) [1, Corollary B] *If  $P$  is a normal cyclic Sylow subgroup of a finite group  $G$ , then  $\psi(G) \leq \psi(P)\psi(G/P)$ , with equality if and only if  $P$  is central in  $G$ .*

(3) [4, Lemma 2.9] *If  $n$  is a natural number,  $p$  and  $q$  are the largest and smallest prime divisors of  $n$ , then*

$$\psi(C_n) > \frac{q}{p+1}n^2.$$

(4) [4, Lemma 2.9] *If  $p$  is a prime and  $m$  is a non-negative integer, then*

$$\psi(C_{p^m}) = \frac{p^{2m+1} + 1}{p + 1}.$$

(5) [4, Lemma 2.2] *Let  $p$  be a prime. Assume that  $G$  is a semidirect product of a normal cyclic  $p$ -subgroup  $C$  and a nontrivial subgroup  $F$  whose order is coprime to  $p$ . Put  $Z = C_F(C)$ . Then*

$$\psi(G) = \psi(C)\psi(Z) + |C|(\psi(F) - \psi(Z)).$$

Observe that [4, Lemma 2.9] states that  $\psi(C_n) \geq 2/(p+1)$ , but the authors prove strict inequality. Also it is easy to see that the number 2 in the numerator can be replaced with  $q$ .

The following statement easily follows from Item (4) of the previous lemma.

**Corollary 1.** *Let  $p$  be a prime and  $m$  be a positive integer. Then the following statements hold.*

(1)

$$\frac{p^m}{\psi(C_{p^m})} < \frac{p+1}{p^2}.$$

(2) *If  $H$  is a proper subgroup of a cyclic group  $C$  and  $p$  is a prime divisor of the index  $|C : H|$ , then*

$$\frac{\psi(H)}{\psi(C)} \leq \frac{1}{p^2 - p + 1}.$$

**Lemma 2.** *If  $H$  is a normal abelian Hall subgroup of a finite group  $G$ , then  $\psi(G) \leq \psi(H)\psi(G/H)$ .*

**Proof.** By the Schur–Zassenhaus theorem,  $G \simeq H \rtimes (G/H)$ . It follows from Item (1) of Lemma 1 that it is sufficient to prove that  $\psi(H \rtimes (G/H)) \leq \psi(H)\psi(G/H)$ . Choose  $h \in H$  and  $g \in G/H$  and denote by  $gh$  and  $g \cdot h$  their products as elements of semidirect and direct products respectively. Since  $H$  is a Hall subgroup,  $|g \cdot h| = |g||h|$ . Now

$$(gh)^{|g|} = h^{g^{|g|-1}} h^{g^{|g|-2}} \dots h^g h.$$

So  $|gh|$  divides  $|g \cdot h|$ , and the lemma is proved.

**Lemma 3.** [4, Proposition 2.5] *Let  $G$  be a finite group and suppose that there exists  $x \in G$  such that  $|G : \langle x \rangle| < 2p$ , where  $p$  is the maximal prime divisor of  $|G|$ . Then one of the following holds:*

- (1)  $G$  has a normal cyclic Sylow  $p$ -subgroup;
- (2)  $G$  is solvable and  $\langle x \rangle$  is a maximal subgroup of  $G$  of index either  $p$  or  $p + 1$ .

**Lemma 4.** [5, Theorem 4.2] *If  $q$  is an odd prime and  $m$  is a positive integer, then*

$$\psi(M_{q^{m+1}}) = \psi(C_{q^m} \times C_q) = \frac{q^{2m+2} + q^3 - q^2 + 1}{q + 1}.$$

*In particular,*

$$\psi(M_{q^3}) = \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1} \psi(C_{q^3}).$$

### 3. Proof of Theorem

Let  $q$  be an odd prime and let  $G$  be a nonabelian  $q^*$ -group of order  $n$ . Suppose that

$$\psi(G) \geq \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1} \psi(C_n). \quad (2)$$

Let  $p$  be the greatest prime divisor of  $|G|$ . Since  $\psi(C_n) > \frac{q}{p+1}n^2$  by Item (3) of Lemma 1,

$$\psi(G) > \frac{(q^6 + q^3 - q^2 + 1)q}{(q^7 + 1)(p + 1)} n^2.$$

So  $G$  contains an element  $x$  such that

$$|x| \geq \frac{\psi(G)}{n} > \frac{(q^6 + q^3 - q^2 + 1)q}{(q^7 + 1)(p + 1)} n.$$

Then

$$|G : \langle x \rangle| < \frac{(q^7 + 1)(p + 1)}{(q^6 + q^3 - q^2 + 1)q}.$$

We proceed by induction on the number of prime divisors of the order of  $G$ . Assume that  $p = q$ . Then

$$|G : \langle x \rangle| < \frac{(q^7 + 1)(q + 1)}{(q^6 + q^3 - q^2 + 1)q}.$$

Since  $G$  is nonabelian and

$$\frac{(q^7 + 1)(q + 1)}{(q^6 + q^3 - q^2 + 1)q} < q^2,$$

the index  $|G : \langle x \rangle|$  is  $q$ . It follows that  $G$  is isomorphic to  $M_{q^{m+1}}$  for  $m \geq 2$  (see, for example, [6, Theorem 1.2]). By Lemma 4 and Item (4) of Lemma 1, the inequality (2) leads to

$$\begin{aligned} \psi(G) &= \frac{q^{2m+2} + q^3 - q^2 + 1}{q + 1} \geq \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1} \psi(C_n) = \\ &= \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1} \cdot \frac{q^{2m+3} + 1}{q + 1}, \end{aligned}$$

or equivalently

$$\begin{aligned}
& (q^{2m+2} + q^3 - q^2 + 1)(q^7 + 1) - (q^6 + q^3 - q^2 + 1)(q^{2m+3} + 1) = \\
& = -q^{2m+6} + q^{2m+5} - q^{2m+3} + q^{2m+2} + q^{10} - q^9 + q^7 - q^6 = \\
& = (-q^{2m+2} + q^6)(q^3 + 1)(q - 1) \geq 0.
\end{aligned}$$

This inequality holds and becomes equality only if  $m = 2$  as stated.

Suppose that  $p > q$ . Then

$$|G : \langle x \rangle| < \frac{(q^7 + 1)(p + 1)}{(q^6 + q^3 - q^2 + 1)q} < p + 1.$$

It follows from Lemma 3 that either  $G$  contains a normal cyclic Sylow  $p$ -subgroup or  $|G : \langle x \rangle| = p$ .

Assume that  $G$  contains a normal cyclic Sylow  $p$ -subgroup  $P$ . By Item (2) of Lemma 1, we have  $\psi(G) \leq \psi(P)\psi(G/P)$  with the equality if and only if  $P$  is central in  $G$ .

If  $G/P$  is a nonabelian group, then

$$\psi(G/P) \leq \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1} \psi(C_{|G/P|})$$

by the inductive hypothesis. Hence

$$\psi(G) \leq \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1} \psi(P)\psi(C_{|G/P|}) = \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1} \psi(C_n).$$

Since we assume (2), the latter inequality must be equality. It is possible only if  $P$  is central. Then  $G$  has the stated structure, that is, it is a direct product of  $M_{q^3}$  and a cyclic group of order coprime to  $q$ .

Let  $G/P$  be an abelian group. Denote by  $H$  a  $p$ -complement of  $G$ . We write  $Z$  for  $C_H(P)$ . By Item (5) of Lemma 1,

$$\psi(G) < \psi(C_{p^m})\psi(H) \left( \frac{\psi(Z)}{\psi(H)} + \frac{p^m}{\psi(C_{p^m})} \right).$$

If  $H$  is cyclic, then

$$\psi(G) < \left( \frac{\psi(Z)}{\psi(H)} + \frac{p^m}{\psi(C_{p^m})} \right) \psi(C_n).$$

Let us bound the right-hand side from above. Consider the fraction  $\psi(Z)/\psi(H)$ . If  $Z = H$ , then  $G = P \times H$  and  $G$  is abelian, contradicting the assumption. So  $Z$  is a proper subgroup of  $H$ . It follows from Item (2) of Corollary 1 that

$$\frac{\psi(Z)}{\psi(H)} \leq \frac{1}{r^2 - r + 1} \tag{3}$$

for a prime divisor  $r$  of  $|H : Z|$ . Since  $G/C_G(P) \leq \text{Aut}(C_{p^m})$ , we have  $H/C_H(P) \leq \text{Aut}(C_{p^m})$ . Therefore,  $|H/C_H(P)|$  divides  $p - 1$ . In particular,  $r$  divides  $p - 1$  and so  $p \geq 2r + 1$ . Since  $r \geq q$ , it follows that  $p \geq 2q + 1$ . Now it follows from (3) and Item (1) of Corollary 1 that

$$\left( \frac{\psi(Z)}{\psi(H)} + \frac{p^m}{\psi(C_{p^m})} \right) \psi(C_n) \leq \left( \frac{1}{(2q + 1)^2} + \frac{1}{2q + 1} + \frac{1}{q^2 - q + 1} \right) \psi(C_n).$$

Since

$$\frac{1}{(2q+1)^2} + \frac{1}{2q+1} + \frac{1}{q^2 - q + 1} < \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1}$$

holds for all  $q > 2$ , we have a contradiction.

If  $H$  is a non-cyclic subgroup of  $G$ , then by Item (5) of Lemma 1, we have

$$\psi(G) = |P|\psi(H) + (\psi(P) - |P|)\psi(Z).$$

Dividing the both sides by  $\psi(C_n)$ , we get

$$\frac{\psi(G)}{\psi(C_n)} = \frac{|P|}{\psi(P)} \frac{\psi(H)}{\psi(C_{|H|})} + \left(1 - \frac{|P|}{\psi(P)}\right) \frac{\psi(Z)}{\psi(C_{|Z|})} \frac{\psi(C_{|Z|})}{\psi(C_{|H|})}.$$

Since  $\psi(Z) \leq \psi(C_{|Z|})$ ,

$$\begin{aligned} \frac{\psi(G)}{\psi(C_n)} &\leq \frac{|P|}{\psi(P)} \frac{\psi(H)}{\psi(C_{|H|})} + \left(1 - \frac{|P|}{\psi(P)}\right) \frac{\psi(C_{|Z|})}{\psi(C_{|H|})} = \\ &= \frac{|P|}{\psi(P)} \left( \frac{\psi(H)}{\psi(C_{|H|})} - \frac{\psi(C_{|Z|})}{\psi(C_{|H|})} \right) + \frac{\psi(C_{|Z|})}{\psi(C_{|H|})}. \end{aligned}$$

Since  $H$  is a non-cyclic  $q^*$ -group, according to [3, Theorem 4]

$$\frac{\psi(H)}{\psi(C_{|H|})} \leq \frac{(q^3 - q + 1)(q + 1)}{q^5 + 1}.$$

Thus

$$\frac{\psi(G)}{\psi(C_n)} \leq \frac{|P|}{\psi(P)} \left( \frac{(q^3 - q + 1)(q + 1)}{q^5 + 1} - \frac{\psi(C_{|Z|})}{\psi(C_{|H|})} \right) + \frac{\psi(C_{|Z|})}{\psi(C_{|H|})}.$$

Since  $\psi(C_{|Z|})/\psi(C_{|H|}) > 0$ , Corollary 1 implies that

$$\frac{\psi(G)}{\psi(C_n)} < \frac{(q+1)(q^3 - q + 1)(q + 1)}{q^2(q^5 + 1)} + \frac{1}{q^2 - q + 1}.$$

The inequality

$$\frac{(q+1)(q^3 - q + 1)(q + 1)}{q^2(q^5 + 1)} + \frac{1}{q^2 - q + 1} < \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1}$$

is equivalent to

$$q^{15} - 3q^{14} + 2q^{12} - 3q^{11} + q^{10} - q^9 - q^7 - 3q^6 + 3q^5 - q^4 - 3q^3 + q^2 - 1 > 0$$

which holds for all  $q > 2$  (indeed,  $q^{15} \geq 3q^{14}$ ,  $2q^{12} > 3q^{11}$ , and so on); that is a contradiction and we finished the case when  $G$  contains a normal cyclic Sylow  $p$ -subgroup.

Assume that  $|G : \langle x \rangle| = p$ . Let  $P \in \text{Syl}_p(G)$  and  $|P| = p^{m+1}$ . There are three options:  $P$  is cyclic (this situation has been considered already),  $P \simeq C_{p^m} \times C_p$ , or  $P \simeq M_{p^{m+1}}$ .

Let us show that  $P$  is normal in  $G$ . If  $Q \in \text{Syl}_q(G)$  then  $Q$  lies in  $\langle x \rangle$ . So  $Q$  is cyclic and  $G$  contains a normal  $q$ -complement  $N$  (see, for example, [7, Theorem 5.14]). By the inductive hypothesis,  $P$  is normal in  $N$  and, therefore, in  $G$ . So  $G = P \rtimes H$  where  $H$  is a cyclic subgroup of  $G$ .

If  $P$  is isomorphic to  $C_{p^m} \times C_p$ , then  $\psi(G) \leq \psi(P)\psi(H)$  by Lemma 2. It follows from Lemma 4 that

$$\frac{\psi(G)}{\psi(C_n)} \leq \frac{\psi(P)}{\psi(C_{p^{m+1}})} = \frac{p^{2m+2} + p^3 - p^2 + 1}{p^{2m+3} + 1} \leq \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1}.$$

The last inequality holds and is strict for all  $m \geq 2$ . If  $m = 0$ , then  $P$  is cyclic. Assume that  $m = 1$ . Since  $H$  acts nontrivially on the cyclic group  $P/C_P(H)$  of order  $p$ , its order has a non-identity common divisor with  $p - 1$ . So  $p \geq 2q + 1$ . The inequality

$$\frac{(2q + 1)^4 + (2q + 1)^3 - (2q + 1)^2 + 1}{(2q + 1)^5 + 1} < \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1}$$

is equivalent to

$$8q^9 + 20q^8 + 24q^7 + 31q^6 + 28q^5 + q^4 - 4q^3 + 17q^2 + 16q + 3 > 0,$$

which holds for all  $q \geq 1$ ; that is a contradiction.

Finally, consider the case  $P \simeq M_{p^{m+1}}$ . Let  $h$  be an arbitrary element of the group  $H$  and  $\varphi$  be the automorphism of  $P$  induced by conjugation by  $h$ . Let  $a, b$  be generators for  $P$  such that

$$a^{p^m} = b^p = 1, a^b = a^{1+p^{m-1}}.$$

Obviously,  $\varphi(g) = g$  for every  $g \in \langle a \rangle$ . If  $\varphi(b) = b^\gamma a^\alpha$  for  $0 \leq \gamma \leq p - 1$  and  $0 \leq \alpha \leq p^m - 1$ , then

$$a^{(1+p^{m-1})} = \varphi(a^b) = a^{\varphi(b)} = a^{b^\gamma a^\alpha} = a^{b^\gamma} = a^{(1+p^{m-1})^\gamma}.$$

Hence

$$(1 + p^{m-1})^\gamma \equiv 1 + p^{m-1} \pmod{p^m},$$

or equivalently

$$(1 + p^{m-1})((1 + p^{m-1})^{\gamma-1} - 1) \equiv 0 \pmod{p^m}.$$

So  $\gamma - 1 \equiv 0 \pmod{p}$ , i.e.  $\gamma = 1$ .

Since  $h$  is an arbitrary element of  $H$ , the latter centralizes the normal series

$$1 \leq \langle a \rangle \leq P.$$

Since  $H$  is a  $p'$ -group, we have  $G = P \times H$ .

By Item (1) of Lemma 1

$$\frac{\psi(G)}{\psi(C_n)} = \frac{\psi(P)\psi(H)}{\psi(C_{|P|})\psi(C_{|H|})}.$$

Recall that  $H$  is a cyclic subgroup, and therefore

$$\frac{\psi(G)}{\psi(C_n)} = \frac{\psi(P)}{\psi(C_{|P|})} \leq \frac{p^6 + p^3 - p^2 + 1}{p^7 + 1} < \frac{q^6 + q^3 - q^2 + 1}{q^7 + 1},$$

that is a contradiction, and the proof is complete.

#### 4. Acknowledgments

The reported study was funded by RFBR and BRFR (project number 20-51-00007), and was carried out within the framework of the state contract of the Sobolev Institute of Mathematics (project no. 0314-2019-0001).

The authors are grateful to the referee for careful reading of the paper and valuable suggestions and comments.

## References

- [1] H. Amiri, S.M. Jafarian Amiri, I.M. Isaacs, *Sums of element orders in finite groups*, Comm. Algebra. **37** (2009), 2978–2980.
- [2] M. Tărnăuceanu, *Detecting structural properties of finite groups by the sum of element orders*, Israel Journal of Mathematics, **238** (2020), 629–637.
- [3] M. Herzog, P. Longobardi, M. Maj, *The second maximal groups with respect to the sum of element orders*, J. Pure Appl. Algebra, **225**, no. 3 (2020), 1–11.
- [4] M. Herzog, P. Longobardi, M. Maj, *An exact upper bound for sums of element orders in non-cyclic finite groups*, J. Pure Appl. Algebra, **222**, no. 7 (2018), 1628–1642.
- [5] R. Shen, G. Chen and C. Wu, *On groups with the second largest value of the sum of element orders*, Comm. Algebra, **43**, no. 6 (2015), 2618–2631.
- [6] Y. Berkovich, *Groups of prime power order*, Walter de Gruyter, Berlin, New York (2008).
- [7] I.M. Isaacs, *Finite Group Theory*. American Mathematical Society, Providence, Rhode Island (2008).