

СИБИРСКИЕ ЭЛЕКТРОННЫЕ  
МАТЕМАТИЧЕСКИЕ ИЗВЕСТИЯ

Siberian Electronic Mathematical Reports

<http://semr.math.nsc.ru>

Том 19, вып. ???-??? (2022)

УДК 510.67

DOI 10.33048/semi.2019.16.xxx

MSC 03C07, 03C52

PROPERTIES OF RANKS FOR FAMILIES OF STRONGLY  
MINIMAL THEORIES

B.SH. KULPESHOV, S.V. SUDOPLATOV

ABSTRACT. We study rank properties for families of strongly minimal theories. A criterion for  $e$ -total transcendence of families of strongly minimal theories is obtained in terms of the description of the language symbols.

**Keywords:** strongly minimal theory, sentence, family of theories, rank for a family of theories.

## 1. INTRODUCTION

In the study of elementary theories, an important role is played by the description of their interrelationships and derived objects, including main characteristics. Essential interrelationships between theories arise when considering their various combinations [1, 2], closures and generating sets for families of theories [3, 4], approximations of theories [5]. Along with combinations of theories in general, various combinations of ordered theories are considered, which allow describing their essential structural properties [6, 7, 8, 9]. Among various approximations of theories, one of the central places is occupied by approximations by theories of finite structures, which define pseudofinite structures [10, 11, 12, 13]. Important derived objects are algebras of definable subfamilies [14, 15]. Main characteristics related to the families of theories are the  $e$ -spectra [1, 16], the ranks for the families of theories [14, 17, 18] and for the formulas [19, 20]. Rank values allow one to determine how complex certain types of families of theories are, both in general case and in relation to natural classes of theories, including classes of substitution theories [21], theories of abelian groups [22], ordered theories [23, 24].

---

KULPESHOV, B.SH., SUDOPLATOV, S.V., PROPERTIES OF RANKS FOR FAMILIES OF STRONGLY MINIMAL THEORIES.

© 2022 KULPESHOV B.SH., SUDOPLATOV S.V..

The research is supported by Russian Scientific Foundation, Project No. 22-21-00044.

Received December, ??, 2021, published ???, ??, 2022.

The present article is concerned the notion of *strong minimality* initially deeply investigated in [25]. We study rank properties for families of strongly minimal theories. A criterion for *e*-total transcendence of families of strongly minimal theories is obtained in terms of the description of the language symbols.

## 2. PRELIMINARY NOTIONS

**Definition 1.** [25] Let  $T$  be a complete theory without finite models. The theory  $T$  is said to be *strongly minimal* if for any model  $\mathcal{M} \models T$ , any formula  $\phi(x, \bar{y})$  and any tuple  $\bar{a} \in M$  either the set  $\phi(\mathcal{M}, \bar{a})$  is finite or the set  $\neg\phi(\mathcal{M}, \bar{a})$  is finite.

**Definition 2.** [17] Let  $\mathcal{T}$  be a family of complete theories in a fixed language  $\Sigma$ ,  $\phi$  be an arbitrary  $\Sigma$ -sentence. Then the set  $\mathcal{T}_\phi := \{T \in \mathcal{T} \mid T \models \phi\}$  is said to be the  $\phi$ -neighbourhood of the family  $\mathcal{T}$ , or the *s-definable* subfamily of  $\mathcal{T}$  defined by the sentence  $\phi$ .

We define the *rank* RS for the family  $\mathcal{T}$  as follows:

- (1)  $\text{RS}(\mathcal{T}) = -1$ , if  $\mathcal{T} = \emptyset$ .
- (2)  $\text{RS}(\mathcal{T}) = 0$ , if  $\mathcal{T}$  is a finite nonempty family.
- (3)  $\text{RS}(\mathcal{T}) \geq 1$ , if  $\mathcal{T}$  is infinite.
- (4)  $\text{RS}(\mathcal{T}) \geq \alpha + 1$ , if there are pairwise inconsistent  $\Sigma$ -sentences  $\phi_n, n \in \omega$ , such that  $\text{RS}(\mathcal{T}_{\phi_n}) \geq \alpha$ .
- (5) If  $\delta$  is a limit ordinal, then  $\text{RS}(\mathcal{T}) \geq \delta$ , if  $\text{RS}(\mathcal{T}) \geq \beta$  for any  $\beta < \delta$ .

We put  $\text{RS}(\mathcal{T}) = \alpha$ , if  $\text{RS}(\mathcal{T}) \geq \alpha$  and  $\neg[\text{RS}(\mathcal{T}) \geq \alpha + 1]$ .

If  $\text{RS}(\mathcal{T}) \geq \alpha$  for any  $\alpha$ , then we put  $\text{RS}(\mathcal{T}) = \infty$ .

A family  $\mathcal{T}$  is called *e-totally transcendental*, or *totally transcendental*, if  $\text{RS}(\mathcal{T})$  is an ordinal.

If a family  $\mathcal{T}$  is *e-totally transcendental*, with  $\text{RS}(\mathcal{T}) = \alpha \geq 0$ , then the *degree*  $\text{ds}(\mathcal{T})$  of  $\mathcal{T}$  is the maximal number of pairwise inconsistent sentences  $\phi_i$  with  $\text{RS}(\mathcal{T}_{\phi_i}) = \alpha$ .

By the definition, if  $\text{RS}(\mathcal{T}) = \alpha \in \text{Ord}$  then  $\text{ds}(\mathcal{T}) \in \omega \setminus \{0\}$ .

## 3. FAMILIES OF STRONGLY MINIMAL THEORIES AND THEIR RANKS

Throughout this article we denote by  $\mathcal{T}_\Sigma$  the family of all strongly minimal theories in a language  $\Sigma$ .

**Proposition 1.** Let  $\Sigma_\kappa^1 := \{P_i^1\}_{i < \kappa}$ , where  $\kappa$  is a cardinal. Then the following holds:

- (1) If  $\kappa < \omega$  then  $\text{RS}(\mathcal{T}_{\Sigma_\kappa^1}) = \kappa$ .
- (2) If  $\kappa \geq \omega$  then  $\text{RS}(\mathcal{T}_{\Sigma_\kappa^1}) = \infty$ .

*Proof.* (1) By virtue of strong minimality, each unary predicate selects a finite or cofinite set in any model of an arbitrary strongly minimal theory.

We construct an infinitely branching tree of length  $\kappa$  for the family  $\mathcal{T}_{\Sigma_\kappa^1}$  of theories. At the first level, we distinguish theories in which the first predicate  $P_1$  selects in any model a finite set consisting of  $n$  elements for each  $n < \omega$ . Further, at the  $i$ -th level ( $i \leq \kappa$ ), we distinguish theories in which the predicate  $P_i$  selects a finite set consisting of  $n$  elements for each  $n < \omega$ . Therefore, we obtain  $\text{RS}(\mathcal{T}_{\Sigma_\kappa^1}) = \kappa$ .

(2) Since there exist infinitely many unary predicates, then one can construct an infinitely branching tree of infinite length, whence  $\text{RS}(\mathcal{T}_{\Sigma_\kappa^1}) = \infty$ .  $\square$

Let  $\Sigma_1^f := \{f^1\}$ , where  $f$  is a unary functional symbol, i.e. any theory  $T$  of the language  $\Sigma_1^f$  satisfies the following sentence:  $T \models (\forall x)(\exists! y)f(x) = y$ .

Let  $T$  be a complete theory of the language  $\Sigma_1^f$ ,  $M \models T$ .

We say that an element  $a \in M$  forms a cycle of length 1 if  $f(a) = a$ .

We say that elements  $a_1, a_2, \dots, a_n \in M$  form a cycle of length  $n$  if  $a_i \neq a_j$  for any  $1 \leq i, j \leq n$  with  $i \neq j$ ,  $f(a_1) = a_2, f(a_2) = a_3, \dots, f(a_{n-1}) = a_n$  and  $f(a_n) = a_1$ .

Consider the following sentences:

$$\begin{aligned} \theta_n^1 &:= \exists x_1 \dots \exists x_n [\wedge_{i \neq j} x_i \neq x_j \wedge \wedge_{i=1}^{n-1} f(x_i) = x_{i+1} \wedge f(x_n) = x_1 \wedge \\ &\forall y_1 \dots \forall y_n (\wedge_{i \neq j} y_i \neq y_j \wedge \wedge_{i=1}^{n-1} f(y_i) = y_{i+1} \wedge f(y_n) = y_1 \rightarrow \vee_{i=1}^n y_i = x_i)], \\ \theta_n^2 &:= \exists x_1 \dots \exists x_n \exists y_1 \dots \exists y_n [\wedge_{i \neq j} x_i \neq x_j \wedge \wedge_{i=1}^{n-1} f(x_i) = x_{i+1} \wedge f(x_n) = x_1 \wedge \\ &\wedge_{i \neq j} y_i \neq y_j \wedge \wedge_{i=1}^{n-1} f(y_i) = y_{i+1} \wedge f(y_n) = y_1 \wedge \\ &\forall z_1 \dots \forall z_n (\wedge_{i \neq j} z_i \neq z_j \wedge \wedge_{i=1}^{n-1} f(z_i) = z_{i+1} \wedge f(z_n) = z_1 \rightarrow \vee_{i=1}^n (z_1 = x_i \vee z_1 = y_i))]. \end{aligned}$$

The sentence  $\theta_n^1$  asserts that there exists a unique cycle of length  $n$ , and the sentence  $\theta_n^2$  asserts that there exist exactly two cycles of length  $n$ . Similarly, we can construct a sentence  $\theta_n^k$  asserting that there exist exactly  $k$  cycles of length  $n$ , for any  $k, n < \omega$ .

**Proposition 2.** Let  $\Sigma_1^f := \{f^1\}$ , where  $f$  is a unary functional symbol. Then  $\text{RS}(\mathcal{T}_{\Sigma_1^f}) = \infty$ .

*Proof.* We construct an infinitely branching tree of infinite length for  $\mathcal{T}_{\Sigma_1^f}$ .

At the first level, we distinguish theories in which there are exactly  $n$  cycles of length 1 for each  $n < \omega$ . At the second level, we distinguish theories in which there are exactly  $n$  cycles of length 2 for each  $n < \omega$ . At the  $k$ -th level ( $k < \omega$ ), we distinguish theories in which there are exactly  $n$  cycles of length  $k$  for each  $n < \omega$ . Thus, we obtain  $\text{RS}(\mathcal{T}_{\Sigma_1^f}) = \infty$ .  $\square$

Since  $k$ -ary predicates and  $k$ -ary operations, for  $k \geq 2$ , allow to interpret unary operations, Proposition 2 implies:

**Corollary 1.** If a language  $\Sigma$  contains a  $k$ -ary predicate symbol or a  $k$ -ary functional symbol, where  $k \geq 2$ , then  $\text{RS}(\mathcal{T}_{\Sigma}) = \infty$ .

**Proposition 3.** [18] If  $\Sigma$  is a language consisting of constant symbols then for the family  $\mathcal{T}$  of all theories in the language  $\Sigma$  either  $\text{RS}(\mathcal{T}) = 1$  and  $\text{ds}(\mathcal{T}) = P(n)$ , where  $P(n)$  is the number of partitions of  $n$ -element sets, if  $\Sigma$  consists of  $n$  symbols, where  $n < \omega$ , or  $\text{RS}(\mathcal{T}) = \infty$ , if  $\Sigma$  has infinitely many symbols.

Proposition 3 implies:

**Fact 1.** Let  $\Sigma_{\kappa}^c := \{c_i\}_{i < \kappa}$ , where  $\kappa$  is a cardinal. Then  $\text{RS}(\mathcal{T}_{\Sigma_{\kappa}^c}) = 0$  and  $\text{ds}(\mathcal{T}_{\Sigma_{\kappa}^c}) = P(\kappa)$ , where  $P(\kappa)$  is the number of partitions of  $n$ -element set, if  $\kappa < \omega$ , or  $\text{RS}(\mathcal{T}_{\Sigma_{\kappa}^c}) = \infty$ , if  $\kappa \geq \omega$ .

**Proposition 4.** Let  $\Sigma_{k,m}^{1,c} := \{P_1^1, \dots, P_k^1, c_1, \dots, c_m\}$ , where  $k, m < \omega$ . Then  $\text{RS}(\mathcal{T}_{\Sigma_{k,m}^{1,c}}) = k$  and  $\text{ds}(\mathcal{T}_{\Sigma_{k,m}^{1,c}}) = P'(k, m)$ , where  $P'(k, m)$  is the total number of actions by predicates  $P_1^1, \dots, P_k^1$  on each of  $P(m)$  partitions of the set  $\{c_1, \dots, c_m\}$  on  $r$  parts,  $1 \leq r \leq m$ :  $P'(k, m) = \sum_r 2^{kr}$ .

*Proof.* By virtue of Proposition 1 and Fact 1, we assert that  $\text{RS}(\mathcal{T}_{\Sigma_{k,m}^{1,c}}) = k$ . Let us now understand that  $\text{ds}(\mathcal{T}_{\Sigma_{k,m}^{1,c}}) = 2^{km}$ . Each of the unary predicates  $P_i^1$  can act on the set of identified constants  $c_j$  in two ways, i.e. either  $P_i^1(c_j)$ , for all identified constants  $c_j$ , or  $\neg P_i^1(c_j)$  (each constant has only two possibilities for a chosen unary predicate). Therefore, taking an arbitrary partition of the set  $\{c_1, \dots, c_m\}$  into  $r$  parts, we obtain the multiplication  $r$  times  $2^k$  of the actions of the predicates  $P_1^1, \dots, P_k^1$  on each element partition equal to  $2^{kr}$ . Running through all the partitions of constants and all variants of the influence of the predicates  $P_1^1, \dots, P_k^1$  on the elements of the partitions of the set  $\{c_1, \dots, c_m\}$ , we obtain the desired equality  $\text{ds}(\mathcal{T}_{\Sigma_{k,m}^{1,c}}) = P'(k, m)$ .  $\square$

**Proposition 5.** [18] *If  $\Sigma$  is a language of 0-ary predicates, then for the family  $\mathcal{T}$  of all theories in the language  $\Sigma$  either  $\text{RS}(\mathcal{T}) = 0$  and  $\text{ds}(\mathcal{T}) = 2^m$ , if  $\Sigma$  consists of  $m$  0-ary predicates, where  $m < \omega$ , or  $\text{RS}(\mathcal{T}) = \infty$ , if  $\Sigma$  has infinitely many symbols.*

Proposition 5 implies:

**Fact 2.** *Let  $\Sigma_\kappa^0 := \{P_i^0\}_{i < \kappa}$ , where  $\kappa$  is a cardinal. Then either  $\text{RS}(\mathcal{T}_{\Sigma_\kappa^0}) = 0$  and  $\text{ds}(\mathcal{T}_{\Sigma_\kappa^0}) = 2^\kappa$ , if  $\kappa < \omega$ , or  $\text{RS}(\mathcal{T}_{\Sigma_\kappa^0}) = \infty$ , if  $\kappa \geq \omega$ .*

In view of Fact 2 if we add to the language  $\Sigma$  finitely many 0-ary predicate symbols then we preserve the  $e$ -total transcendence of the family of all strongly minimal theories in the expanded language  $\Sigma'$ , whereas adding to the language  $\Sigma$  infinitely many 0-ary predicate symbols we obtain the family  $\mathcal{T}'_\Sigma$  with  $\text{RS}(\mathcal{T}'_\Sigma) = \infty$ . Thus, Propositions 1, 2, 4, Corollary 1, and Facts 1, 2 imply the following:

**Theorem 1.** *A family  $\mathcal{T}_\Sigma$  of strongly minimal theories is  $e$ -totally transcendental if and only if the language  $\Sigma$  consists of finitely many constant symbols as well as of finitely many 0-ary and unary predicate symbols.*

**Remark 1.** Assuming that distinct constant symbols are interpreted by distinct elements,  $e$ -totally transcendental families of strongly minimal theories in Theorem 1 admit arbitrarily many constant symbols in the language.

**Remark 2.** Since  $\text{RS}(\mathcal{T}) \leq \text{RS}(\mathcal{T}')$  for  $\mathcal{T} \subseteq \mathcal{T}'$  then the criterion for  $e$ -total transcendence of families of strongly minimal theories in Theorem 1 can be transformed for any family  $\mathcal{T}'$  containing the family  $\mathcal{T}_\Sigma$  of strongly minimal theories of given language  $\Sigma$ . In particular, it holds for families of  $\omega$ -stable, superstable, and stable theories.

REFERENCES

[1] S.V. Sudoplatov, *Combinations of structures*, The Bulletin of Irkutsk State University. Series Mathematics, **24** (2018), 65–84.  
 [2] S.V. Sudoplatov, *Combinations related to classes of finite and countably categorical structures and their theories*, Siberian Electronic Mathematical Reports, **14** (2017), 135–150.  
 [3] S.V. Sudoplatov, *Closures and generating sets related to combinations of structures*, The Bulletin of Irkutsk State University. Series Mathematics, **16** (2016), 131–144.  
 [4] S.V. Sudoplatov, *Families of language uniform theories and their generating sets*, The Bulletin of Irkutsk State University. Series Mathematics, **17** (2016), 62–76.  
 [5] S.V. Sudoplatov, *Approximations of theories*, Siberian Electronic Mathematical Reports, **17** (2020), 715–725.

- [6] A.B. Altayeva, B.Sh. Kulpeshov, S.V. Sudoplatov, *E-combinations of  $\aleph_0$ -categorical linear orders*, Journal of Mathematics, Mechanics and Computer Science, **103**:3 (2019), 3–12.
- [7] B.Sh. Kulpeshov, S.V. Sudoplatov, *P-combinations of ordered theories*, Lobachevskii Journal of Mathematics, **41**:2 (2020), 227–237.
- [8] B.Sh. Kulpeshov, S.V. Sudoplatov, *P\*-combinations of almost  $\omega$ -categorical weakly o-minimal theories*, Lobachevskii Journal of Mathematics, **42**:4 (2021), 743–750.
- [9] B.Sh. Kulpeshov, S.V. Sudoplatov, *E-combinations of almost omega-categorical weakly o-minimal theories*, Algebra and model theory 13, Collection of papers edited by A.G. Pinus, E.N. Poroshenko, and S.V. Sudoplatov, Novosibirsk State Technical University, Novosibirsk, 2021. — P. 83–89.
- [10] E. Rosen, *Some Aspects of Model Theory and Finite Structures*, The Bulletin of Symbolic Logic, **8**:3 (2002), 380–403.
- [11] J. Väänänen, *Pseudo-finite model theory*, Matematica Contemporanea, **24** (2003), 169–183.
- [12] G. Cherlin, E. Hrushovski, *Finite Structures with Few Types*, Annals of Mathematics Studies, No. 152, Princeton University Press, Princeton, Oxford, 2003.
- [13] H.D. Macpherson, Ch. Steinhorn, *Definability in the classes of finite structures*, Finite and Algorithmic Model Theory. London Mathematical Society Lecture Notes series: 379 / eds.: J. Esparza, C. Michaux, Ch. Steinhorn, Cambridge University Press, Cambridge, 2011. — P. 140–176.
- [14] N.D. Markhabatov, S.V. Sudoplatov, *Definable subfamilies of theories, related calculi and ranks*, Siberian Electronic Mathematical Reports, **17** (2020), 700–714.
- [15] N.D. Markhabatov, S.V. Sudoplatov, *Algebras for definable families of theories*, Siberian Electronic Mathematical Reports, **16** (2019), 600–608.
- [16] S.V. Sudoplatov, *Relative e-spectra and relative closures for families of theories*, Siberian Electronic Mathematical Reports, **14** (2017), 296–307.
- [17] S.V. Sudoplatov, *Ranks for families of theories and their spectra*, Lobachevskii Journal of Mathematics, **42**:12 (2021), 2959–2968.
- [18] N.D. Markhabatov, S.V. Sudoplatov, *Ranks for families of all theories of given languages*, Eurasian Mathematical Journal, **12**:2 (2021), 52–58.
- [19] S.V. Sudoplatov, *Formulas and properties, their links and characteristics*, Mathematics, **9**:12 (2021), 16 pp.
- [20] In.I. Pavlyuk, S.V. Sudoplatov, *Formulas and properties for families of theories of Abelian groups*, Bulletin of Irkutsk State University. Series Mathematics, **36** (2021), 95–109.
- [21] N.D. Markhabatov, *Ranks for families of permutation theories*, The Bulletin of Irkutsk State University. Series Mathematics, **28** (2019), 86–95.
- [22] In.I. Pavlyuk, S.V. Sudoplatov, *Ranks for families of theories of abelian groups*, The Bulletin of Irkutsk State University. Series Mathematics, **28** (2019), 96–113.
- [23] B.Sh. Kulpeshov, S.V. Sudoplatov, *Ranks and approximations for families of ordered theories*, Algebra and model theory 12, Collection of papers edited by A.G. Pinus, E.N. Poroshenko, and S.V. Sudoplatov, Novosibirsk State Technical University, Novosibirsk, 2019. — P. 32–40.
- [24] B.Sh. Kulpeshov, In.I. Pavlyuk, S.V. Sudoplatov, *On criterion of total transcendency for families of ordered theories*, Abstracts of International conference "Maltsev Meetings", Novosibirsk, 2021, P. 152.
- [25] J.T. Baldwin, A.H. Lachlan, *On strongly minimal sets*, The Journal of Symbolic Logic, **36**:1 (1971), 79–96.

BEIBUT SHAIYKOVICH KULPESHOV  
 KAZAKH BRITISH TECHNICAL UNIVERSITY  
 TOLE BI STREET, 59  
 050000, ALMATY, KAZAKHSTAN.  
*E-mail address:* [b.kulpeshov@kbtu.kz](mailto:b.kulpeshov@kbtu.kz)

NOVOSIBIRSK STATE TECHNICAL UNIVERSITY  
 K. MARX AVENUE, 20  
 630073, NOVOSIBIRSK, RUSSIA.  
*E-mail address:* [kulpeshov@corp.nstu.ru](mailto:kulpeshov@corp.nstu.ru)

SERGEY VLADIMIROVICH SUDOPLATOV  
NOVOSIBIRSK STATE TECHNICAL UNIVERSITY  
K. MARX AVENUE, 20  
630073, NOVOSIBIRSK, RUSSIA.  
*E-mail address:* `sudoplatov@corp.nstu.ru`

SOBOLEV INSTITUTE OF MATHEMATICS,  
ACADEMICIAN KOPTYUG AVENUE, 4  
630090, NOVOSIBIRSK, RUSSIA.  
*E-mail address:* `sudoplat@math.nsc.ru`