

СИБИРСКИЕ ЭЛЕКТРОННЫЕ
МАТЕМАТИЧЕСКИЕ ИЗВЕСТИЯ

Siberian Electronic Mathematical Reports

<http://semr.math.nsc.ru>

Том 19, вып. ???-??? (2022)

УДК 510.67

DOI 10.33048/semi.2019.16.xxx

MSC 03C07, 03C10, 03C68

ARITIES AND ARITIZABILITIES OF FIRST-ORDER THEORIES

S.V. SUDOPLATOV

ABSTRACT. We study and describe possibilities for arities of elementary theories and of their expansions. Links for arities with respect to Boolean algebras, to disjoint unions and to compositions of structures are shown. Arities and aritizabilities are semantically characterized. The dynamics for arities of theories is described. Possibilities for arities and aritizabilities of theories are illustrated by a series of natural geometric, combinatorial and model-theoretic examples.

Keywords: elementary theory, arity, expansion, aritizability.

Arities of theories are important characteristics showing complexity measures of theories [1, 2] and reducing all definable sets to definable ones generated by cylinders of special forms. It is closely linked with cylindric algebras reflecting semantically first-order calculi [3, 4, 5, 6, 7].

Special cases for arities of theories, especially binary, ternary and related ordered theories are studied in a series of papers including [8, 9, 10, 11, 12, 13, 14]. Structures and links with respect to binary formulas are investigated both in general case [15, 16] and for a series of natural classes of theories [17, 18, 19, 20, 21, 22, 23].

In the present paper we adapt the general cylindric approach and describe semantically arities of theories, properties related to the n -arity and n -aritizability of theories and their dynamics.

The paper is organized as follows. In Section 1, we consider arities of formulae and arities of theories, describe possibilities of arities, describe arities for a series of natural theories including n -transitive theories (Proposition 1.6), characterize

SUDOPLATOV, S.V., ARITIES AND ARITIZABILITIES OF FIRST-ORDER THEORIES.

© 2022 SUDOPLATOV S.V.

The work of the author was carried out in the framework of the State Contract of the Sobolev Institute of Mathematics, Project No. FWNF-2022-0012 (Section 1), and of Russian Scientific Foundation, Project No. 22-21-00044 (Sections 2 and 3).

Received ???, ??, 2022, published ???, ??, 2022.

the ω -categoricity and the stability of n -ary theories (Theorems 1.13 and 1.14). In Section 2, we introduce a series of notions for the aritizability of a theory, describe sufficient conditions and criteria for the aritizabilities in terms of Boolean algebras (Proposition 2.7), disjoint unions and compositions (Theorems 2.8 and 2.10). In Section 3, possibilities for aritizabilities are characterized semantically (Theorems 3.1–3.4), constantizable theories are studied (Proposition 3.6, Corollaries 3.7, 3.8), the existence of a non-aritizable expansion is characterized (Proposition 3.9) and the dynamics for arities of theories is described (Theorem 3.10).

Throughout we consider complete first-order theories T .

1. n -ARY FORMULAE AND THEORIES

Definition [24]. A theory T is said to be Δ -based, where Δ is some set of formulae without parameters, if any formula of T is equivalent in T to a Boolean combination of formulae in Δ .

For Δ -based theories T , it is also said that T has *quantifier elimination* or *quantifier reduction* up to Δ . It is said that T has *quantifier elimination* if T has quantifier elimination up to the set of quantifier free formulae.

Definition [15, 24]. Let Δ be a set of formulae of a theory T , and $p(\bar{x})$ a type of T lying in $S(T)$. The type $p(\bar{x})$ is said to be Δ -based if $p(\bar{x})$ is isolated by a set of formulas $\varphi^\delta \in p$, where $\varphi \in \Delta$, $\delta \in \{0, 1\}$.

The following lemma, being a corollary of Compactness Theorem, noticed in [24].

Lemma 1.1. *A theory T is Δ -based if and only if, for any tuple \bar{a} of any (some) weakly saturated model of T , the type $\text{tp}(\bar{a})$ is Δ -based.*

Definition (cf. [1]). An elementary theory T is called *unary*, or *1-ary*, if any T -formula $\varphi(\bar{x})$ is T -equivalent to a Boolean combination of T -formulas, each of which is of one free variable, and of formulas of form $x \approx y$.

For a natural number $n \geq 1$, a formula $\varphi(\bar{x})$ of a theory T is called *n -ary*, or an *n -formula*, if $\varphi(\bar{x})$ is T -equivalent to a Boolean combination of T -formulas, each of which is of n free variables.

For a natural number $n \geq 2$, an elementary theory T is called *n -ary*, or an *n -theory*, if any T -formula $\varphi(\bar{x})$ is n -ary.

A theory T is called *binary* if T is 2-ary, it is called *ternary* if T is 3-ary, etc.

We will admit the case $n = 0$ for n -formulae $\varphi(\bar{x})$. In such a case $\varphi(\bar{x})$ is just T -equivalent to a sentence $\forall \bar{x} \varphi(\bar{x})$.

If T is a theory such that T is n -ary and not $(n - 1)$ -ary then the value n is called the *arity* of T and it is denoted by $\text{ar}(T)$. If T does not have any arity we put $\text{ar}(T) = \infty$.

Similarly, for a formula φ of a theory T we denote by $\text{ar}_T(\varphi)$ the natural value n if φ is n -ary and not $(n - 1)$ -ary. If φ does not have any arity we put $\text{ar}_T(\varphi) = \infty$. If a theory T is fixed we write $\text{ar}(\varphi)$ instead of $\text{ar}_T(\varphi)$.

By the definition any n -theory is Δ_n -based, where Δ_n consists of formulae with n free variables and formulae of the form $x \approx y$. It implies that theories of n -element models \mathcal{M} are n -ary and based by formulae describing these n -element structures and differences/coincidences of elements.

Using Lemma 1.1 we obtain the following characterization for the n -arity of a formula:

Proposition 1.2. *A T -formula $\varphi(\bar{x})$ is not n -ary if and only if for any T -formulae $\psi_i(\bar{x}_i)$ with subtuples \bar{x}_i of the tuple \bar{x} having $l(\bar{x}_i) = n$ and $T \vdash \varphi(\bar{x}) \rightarrow \psi_i(\bar{x}_i)$, there exists a tuple $\bar{a} \in \mathcal{M} \models T$ such that $\mathcal{M} \models \psi_i(\bar{a}_i) \wedge \neg\varphi(\bar{a})$, where \bar{a}_i is a subtuple of \bar{a} consisting of substitutions of elements of \bar{a} instead of correspondent elements of \bar{x}_i .*

By the definition the notion of n -arity is local and reduced to finite sublanguages:

Proposition 1.3. *A theory T of a language Σ is n -ary if and only if for any T -formula $\varphi(\bar{x})$ there is a finite sublanguage $\Sigma' \subseteq \Sigma$ such that $T \vdash \varphi(\bar{x}) \leftrightarrow \psi(\bar{x})$, where $\psi(\bar{x})$ is a Boolean combination of n -formulae.*

Proposition 1.4. *If \mathcal{M} is a n -element structure, for $n \in \omega$, then $\text{ar}(\text{Th}(\mathcal{M})) \leq n$.*

Proof. Since $|\mathcal{M}| = n$ each $\text{Th}(\mathcal{M})$ -formula $\varphi := \varphi(x_1, \dots, x_m)$ is $\text{Th}(\mathcal{M})$ -equivalent to a disjunction of substitutions of variables x_{i_1}, \dots, x_{i_n} instead of x_1, \dots, x_m into the formula φ , as required.

Remark 1.5. (cf. [4, 7]) Since negations of formulas with n free variables again have n free variables, witnessing the n -arity of a formula it suffices to consider positive Boolean combinations of formulas with n free variables, i.e., conjunctions and disjunctions of formulas with n free variables.

Thus for the description of definable sets for models \mathcal{M} of n -theories it suffices describe links between definable sets A and B for n -formulae $\varphi(\bar{x})$ and $\psi(\bar{y})$, respectively, and definable sets C and D for $\varphi(\bar{x}) \wedge \psi(\bar{y})$ and $\varphi(\bar{x}) \vee \psi(\bar{y})$, respectively.

If $\bar{x} = \bar{y}$ then $C = A \cap B$ and $D = A \cup B$, i.e., conjunctions and disjunctions work as set-theoretic intersections and unions.

If \bar{x} and \bar{y} are disjoint then $C = A \times B$ and $D = (A + B)_{\mathcal{M}} = \{\langle \bar{a}, \bar{b} \rangle \mid \bar{a} \in A \text{ and } \bar{b} \in B, \text{ or } \bar{a} \in M \text{ and } \bar{b} \in B\}$, i.e., C is the Cartesian product of A and B , and D is the (*generalized*) Cartesian sum of A and B in the model \mathcal{M} .

If $\bar{x} \neq \bar{y}$, and \bar{x} and \bar{y} have common variables, then C and D are represented as a *mixed product* and a *mixed sum*, respectively, working partially as intersection and union, for common variables, and partially as Cartesian product and Cartesian sum, for disjoint variables.

If \bar{x} and \bar{y} consist of pairwise disjoint variables and $\bar{x} \subseteq \bar{y}$ and $\bar{x} \neq \bar{y}$ then for any formula $\varphi(\bar{x})$ the set of solution of the formula $\varphi(\bar{x}) \wedge (\bar{y} \approx \bar{y})$ in \mathcal{M} is called a *cylinder* with respect to $M^{l(\bar{y})}$ and generated by the set of solutions $\varphi(\mathcal{M})$. In any case generating sets for cylinders coincide their *projections*, i.e., sets of solutions for formulae $\exists \bar{z} \varphi(\bar{x})$, where $\bar{z} \subset \bar{x}$.

Since n -formulae produce cylinders on Cartesian products of universes, definable sets of n -ary theories are composed by Boolean combinations of definable cylinders, i.e., of elements of cylindric algebras.

Definition (cf. [2]). For a natural number n , a theory T is called n -transitive if each n -type $q(x_1, \dots, x_n) \in S(T)$ is forced by its restriction to the empty language.

By the definition the n -transitivity of a theory T means that a saturated model \mathcal{M} of T has a n -transitive automorphism group, i.e., for any two tuples $(a_1, \dots, a_n), (b_1, \dots, b_n) \in M$ with $a_i \neq a_j$ and $b_i \neq b_j$ for $i \neq j$ there is an automorphism $f \in \text{Aut}(\mathcal{M})$ such that $f(a_1) = b_1, \dots, f(a_n) = b_n$.

Proposition 1.6. *If a theory T is n -transitive and non- $(n+1)$ -transitive then T is not an n -theory.*

Proof. Since T is n -transitive, cylinders defined by n -formulae are reduced to the cylinders defined by the formulae for the empty language, i.e., they are defined by equalities and inequalities. As T is not $(n+1)$ -transitive then there is a \emptyset -definable set $X \subset M^{n+1}$ in a model $\mathcal{M} \models T$ which is not reduced to the cylinders defined by the formulae for the empty language. It means that a formula $\varphi(x_1, \dots, x_{n+1})$ defining X is not T -equivalent to n -formulae. Thus T is not an n -theory, as required.

Clearly, generic constructions [15, 25] allow to produce, for each $n \geq 1$, n -transitive and non- $(n+1)$ -transitive theories with unique $(n+1)$ -ary predicates and having quantifier elimination.

For instance, the theory T_2 of structure $\mathcal{M}_2 = \langle \{a, b, c\}; R_2 \rangle$ with the binary relation $R_2 = \{(a, b), (b, c), (c, a)\}$ has quantifier elimination, is 1-transitive, not 2-transitive, and thus $\text{ar}(T) = 2$.

Indeed, the automorphism group $\text{Aut}(\mathcal{M}_2)$ is 1-transitive, whereas it is not 2-transitive, since the pairs in R_2 and in its complement are not connected by automorphisms. Finally T_2 has quantifier elimination since n -tuples with pairwise distinct coordinates, for $n \geq 2$, are connected by an automorphism iff their correspondent coordinates simultaneously satisfy or not satisfy R_2 .

Similarly the theory T_3 of structure $\mathcal{M}_3 = \langle \{a, b, c, d\}; R_3 \rangle$ with the ternary relation

$$R_3 = \{(a, b, c), (b, a, d), (b, c, d), (c, b, a), (a, c, d), (c, a, b), \\ (c, d, a), (d, c, b), (d, a, b), (a, d, c), (b, d, a), (d, b, c)\}$$

has quantifier elimination, is 2-transitive, not 3-transitive, and thus $\text{ar}(T_3) = 3$.

Indeed, any two pairs (a_1, a_2) and (b_1, b_2) of distinct elements in $\{a, b, c, d\}$ are connected by an automorphism. Say, taking the pairs (a, b) and (b, a) we construct $f \in \text{Aut}(\mathcal{M}_3)$ mapping $f(a) = b, f(b) = a, f(c) = d, f(d) = c$; taking the pairs (a, b) and (b, c) we define $f \in \text{Aut}(\mathcal{M}_3)$ mapping $f(a) = b, f(b) = c, f(c) = d, f(d) = a$; taking the pairs (a, b) and (c, d) we construct $f \in \text{Aut}(\mathcal{M}_3)$ mapping $f(a) = c, f(b) = d, f(c) = a, f(d) = b$, etc., producing the 2-transitivity of T_3 . At the same time T_3 is not 3-transitive since the triples (a, b, c) and (a, b, d) are not connected by an automorphism. Finally T_3 has quantifier elimination since n -tuples with pairwise distinct coordinates, for $n \geq 3$, are connected by an automorphism iff their correspondent coordinates simultaneously satisfy or not satisfy R_3 .

These examples can be naturally spread for n -ary relations R_n with $n \geq 4$ and $|R_n| = \frac{n!}{2}$. In view of Proposition 1.6 it implies the following:

Corollary 1.7. *For any natural $n \geq 1$ there is a theory T_n with $\text{ar}(T_n) = n$.*

The following examples illustrate values $\text{ar}(T) = n$.

Example 1.8. [26] For any theory T_f of an unar, i.e., of one unary operation f , $\text{ar}(T) \leq 2$. There are both theories T_{f_1} with $\text{ar}(T_{f_1}) = 1$ and theories T_{f_2} with $\text{ar}(T_{f_2}) = 2$. For instance, f_1 can be taken identical, and f_2 — a successor function on at least 3-element set.

Example 1.9. [26] For any theory T_Γ of an acyclic graph Γ with unary predicates, $\text{ar}(T_\Gamma) \leq 2$.

Example 1.10. Let E be the following equivalence relation on the set \mathbf{R}^n :

$$\{(M, N) \mid M(x_1, \dots, x_n), N(y_1, \dots, y_n) \in \mathbf{R}^n, x_1^2 + \dots + x_n^2 = y_1^2 + \dots + y_n^2\}.$$

Equivalence classes for the concentric spheres in \mathbf{R}^n can not be reconstructed via cylinders defined by projections which form concentric balls and circles. The homogeneity of equivalence classes implies that each formula in the language $\langle E \rangle$ is reduced to a Boolean combination of $2n$ -formulas. Thus $\text{Th}(\langle \mathbf{R}, E \rangle)$ is a $2n$ -theory which is not an $(2n - 1)$ -theory.

Adding a disjoint unary predicate P and a bijection f between the set of spheres and P we obtain names for spheres and an additional coordinate for generating formulas for a basedness. Thus we form a $(2n + 1)$ -theory which is not an $2n$ -theory.

Hence all possibilities for $\text{ar}(T) = n$ are realized.

Example 1.11. Taking a non-degenerated algebraic surface at \mathbf{R}^n which is not reduced to cylinders we obtain a defining formula $\varphi(\bar{x})$, $l(\bar{x}) = n + 1$, which is $(n+1)$ -formula and not an n -formula. In particular, non-degenerated non-cylindrical surfaces of the second order in \mathbf{R}^3 are defined by formulas φ with $\text{ar}(\varphi) = 3$. For instance, taking the formula $x^2 + y^2 + z^2 = 1$ for the sphere S we obtain projections $x^2 + y^2 \leq 1$, $x^2 + z^2 \leq 1$, $y^2 + z^2 \leq 1$ which can not allow to reconstruct S by their Boolean combinations.

Example 1.12. Recall [8, 11, 12] that a *circular*, or *cyclic* order relation is described by a ternary relation K_3 satisfying the following conditions:

- (co1) $\forall x \forall y \forall z (K_3(x, y, z) \rightarrow K_3(y, z, x))$;
- (co2) $\forall x \forall y \forall z (K_3(x, y, z) \wedge K_3(y, x, z) \leftrightarrow x = y \vee y = z \vee z = x)$;
- (co3) $\forall x \forall y \forall z (K_3(x, y, z) \rightarrow \forall t [K_3(x, y, t) \vee K_3(t, y, z)])$;
- (co4) $\forall x \forall y \forall z (K_3(x, y, z) \vee K_3(y, x, z))$.

Clearly, $\text{ar}(K_3(x, y, z)) = 3$ if the relation has at least three element domain.

Indeed, the relation K_3 on a 3-element set $M_3 = \{a, b, c\}$ consists of 24 triples, with

$$K_3 = \{(a_1, a_2, a_3) \in M_3 \mid a_i = a_j \text{ for some } i \neq j\} \cup \{(a, b, c), (b, c, a), (c, a, b)\}$$

producing a 2-transitive theory with quantifier elimination, which is not 3-transitive.

Hence, theories with infinite circular order relations are at least 3-ary.

The following generalization of circular order produces a *n-ball*, or *n-spherical*, or *n-circular* order relation, for $n \geq 4$, which is described by a n -ary relation K_n satisfying the following conditions:

- (nso1) $\forall x_1, \dots, x_n (K_n(x_1, x_2, \dots, x_n) \rightarrow K_n(x_2, \dots, x_n, x_1))$;
- (nso2) $\forall x_1, \dots, x_n \left((K_n(x_1, \dots, x_i, \dots, x_j, \dots, x_n) \wedge \right.$

$$\left. \wedge K_n(x_1, \dots, x_j, \dots, x_i, \dots, x_n) \right) \leftrightarrow \bigvee_{1 \leq k < l \leq n} (x_k \approx x_l) \text{ for any } 1 \leq i < j \leq n;$$

- (nso3) $\forall x_1, \dots, x_n \left(K_n(x_1, \dots, x_n) \rightarrow \forall t \left(\bigvee_{i=1}^n K_n(x_1, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n) \right) \right)$;
- (nso4) $\forall x_1, \dots, x_n (K_n(x_1, \dots, x_i, \dots, x_j, \dots, x_n) \vee K_n(x_1, \dots, x_j, \dots, x_i, \dots, x_n))$, $1 \leq i < j \leq n$.

Clearly, $\text{ar}(K_n(x_1, \dots, x_n)) = n$ if the relation has at least n -element domain. Thus, theories with infinite n -spherical order relations are at least n -ary.

Theorem 1.13. *An n -ary theory T is ω -categorical if and only if there are finitely many T -non-equivalent formulas with n free variables.*

Proof. If T is ω -categorical then by Ryll-Nardzewski Theorem there are finitely many T -non-equivalent formulas with m free variables for every m , in particular, for $m = n$. Conversely, we again apply Ryll-Nardzewski Theorem showing that there are finitely many T -non-equivalent formulas with m free variables for every m . If $m \leq n$ then there are finitely many T -non-equivalent formulas with m free variables by the monotony of this property with respect to the number of free variables. If $m > n$ then by the n -arity of T each T -formula $\varphi(x_1, \dots, x_m)$ is T -equivalent to a Boolean combination of formulas with n free variables. Since there are finitely many T -non-equivalent possibilities for these formulas, Boolean combinations produce finitely many possibilities, too. As there are finitely many T -non-equivalent formulas with m free variables for every m then T is ω -categorical by Ryll-Nardzewski Theorem, as required.

Recall [27] that a formula $\varphi(\bar{x}, \bar{y})$ of a theory T is *stable* if there are no tuples $\bar{a}_n, \bar{b}_n, n \in \omega$, such that $\models \varphi(\bar{a}_i, \bar{b}_j) \Leftrightarrow i \leq j$. The theory T is called *stable* if every T -formula is stable.

In [28], it was shown that any Boolean combination of stable formulas is again a stable formula. Thus, using the definition of n -ary theory we obtain the following:

Theorem 1.14. *An n -ary theory T is stable if and only if each T -formula with n free variables is stable.*

2. ARITIZABLE FORMULAE AND THEORIES

Definition. A T -formula $\varphi(\bar{x})$ is called *n -expansible*, or *n -arizable*, or *n -aritzable*, if T has an expansion T' such that $\varphi(\bar{x})$ is T' -equivalent to a Boolean combination of T' -formulas with n free variables.

A theory T is called *n -expansible*, or *n -arizable*, or *n -aritzable*, if there is an n -ary expansion T' of T .

A theory T is called *arizable* or *aritzable*, if T is n -aritzable for some n .

A 1-aritzable theory is called *unary-able*, or *unary-tizable*. A 2-aritzable theory is called *binary-tizable* or *binarizable*, a 3-aritzable theory is called *ternary-tizable* or *ternarizable*, etc.

By the definition any n -theory is n -expansible, by itself, and if T is n -expansible then T is m -expansible for each $m > n$.

Besides each formula of an n -expansible theory is n -expansible, too, but not vice versa in the following sense: if each formula of a theory T is n -expansible, it can not guarantee that a resulting expansion T' , witnessing that n -expansibility, is coordinated enough such that it is n -ary or at least n -expansible.

Proposition 2.1. *Any theory of a finite structure \mathcal{M} is binarizable.*

Proof. Let $M = \{a_1, \dots, a_m\}$. For any pair $\langle a_i, a_j \rangle, i, j \leq m$, we introduce new binary singleton predicate $B_{i,j} = \{\langle a_i, a_j \rangle\}$. We denote the resulted expansion of \mathcal{M} by \mathcal{M}' , and the theory $\text{Th}(\mathcal{M}')$ expanding given theory $T = \text{Th}(\mathcal{M})$ by T' . Now an arbitrary T' -formula $\varphi(\bar{x})$, with $\bar{x} = \langle x_1, \dots, x_n \rangle$, has finitely many solutions $\bar{b} = \langle a_{k_1}, \dots, a_{k_n} \rangle$ in \mathcal{M}' . We collect these solutions into a set Z . Without loss of generality $Z \neq \emptyset$ since for $Z = \emptyset, T \vdash \varphi(\bar{x}) \leftrightarrow \neg x_i \approx x_i$ for any x_i . Now the formula $\varphi(\bar{x})$ is T' -equivalent to the following Boolean combination of binary formulae:

$$\bigvee_{\langle a_{k_1}, \dots, a_{k_n} \rangle \in Z} \bigwedge_{i, j \leq n} B_{k_i, k_j}(x_i, x_j),$$

as required.

Remark 2.2. If $m, n \in \omega \setminus \{0\}$ and M is an m -element set, then M^n has $2^{(m^n)}$ subsets producing distinct n -ary predicates Q_i , $i < 2^{(m^n)}$. Since by Stone Theorem any finite Boolean has 2^l elements with l generators, there are m^n independent predicates Q'_j whose Boolean combinations produce all these predicates. Taking a quantifier free formula $\varphi(x_1, \dots, x_k)$, for $k \geq n$, composed by these independent predicates Q'_j and having a perfect disjunctive normal form we obtain $A_k^n \cdot 2^{m^n}$ possibilities for disjunctive members, where A_k^n is used to calculate the number of choice of n variables among x_1, \dots, x_k and there are 2^{m^n} possibilities for positive and negative entries of Q'_j . Now there are $2^{A_k^n \cdot 2^{m^n}}$ possibilities for $\varphi(x_1, \dots, x_k)$, big enough. At the same time, using the arguments for Proposition 2.1 we can obtain all definable subsets of M^n just using m^2 singleton binary relations.

Applying arguments for Proposition 2.1 we immediately obtain the following:

Proposition 2.3. *Any formula of a theory having finitely many solutions is binarizable.*

Proposition 2.1 can be strengthened as follows:

Proposition 2.4. *Any theory of a finite structure \mathcal{M} is unary-tizable.*

Proof. Let $M = \{a_1, \dots, a_m\}$. For element a_i , $i \leq m$, we introduce new unary singleton predicate $U_i = \{a_i\}$. We denote the resulted expansion of \mathcal{M} by \mathcal{M}' , and the theory $\text{Th}(\mathcal{M}')$ expanding given theory $T = \text{Th}(\mathcal{M})$ by T' . Now an arbitrary T' -formula $\varphi(\bar{x})$, with $\bar{x} = \langle x_1, \dots, x_n \rangle$, has finitely many solutions $\bar{b} = \langle a_{k_1}, \dots, a_{k_n} \rangle$ in \mathcal{M}' . We collect these solutions into a set Z . Without loss of generality $Z \neq \emptyset$. Now the formula $\varphi(\bar{x})$ is T' -equivalent to the following Boolean combination of formulae each of which with one free variable:

$$\bigvee_{\langle a_{k_1}, \dots, a_{k_n} \rangle \in Z} \bigwedge_{i, j \leq n} U_{k_i}(x_i),$$

as required.

Thus $|\mathcal{M}|$ -many unary predicates produce a unary expansion of the theory $\text{Th}(\mathcal{M})$ with finite \mathcal{M} . Besides using the proof of Proposition 2.4 we have:

Proposition 2.5. *Any formula of a theory having finitely many solutions is unary-tizable.*

Remark 2.6. By the definition for any natural n both n -ary formulae and n -arizable formulae of a fixed theory T are closed under Boolean combinations. Therefore taking a model $\mathcal{M} \models T$ and collecting in sets $\text{BA}_{kn}(\mathcal{M})$ and $\text{BA}'_{kn}(\mathcal{M})$ definable sets which are defined by n -ary, respectively, n -arizable formulae with k free variables we obtain Boolean algebras $\mathcal{BA}_{kn}(\mathcal{M})$ and $\mathcal{BA}'_{kn}(\mathcal{M})$ of these definable sets.

Clearly, $\mathcal{BA}_{kn}(\mathcal{M}) \subseteq \mathcal{BA}'_{kn}(\mathcal{M})$, and the equality $\mathcal{BA}_{kn}(\mathcal{M}) = \mathcal{BA}'_{kn}(\mathcal{M})$ means that any n -arizable formula $\varphi(x_1, \dots, x_n)$ of T is already n -ary.

In view of Proposition 2.5 the algebra $\text{BA}'_{kn}(\mathcal{M})$ satisfies the following condition: if X and Y are \emptyset -definable subsets of M^k with finite symmetric difference $X \div Y$ then $X \in \text{BA}'_{kn}(\mathcal{M})$ iff $Y \in \text{BA}'_{kn}(\mathcal{M})$. At the same time $\text{BA}_{kn}(\mathcal{M})$ can be not closed under finite symmetric difference since, for instance, there are theories of finite structures which are not n -ary but by Proposition 2.4 all theories of finite structures are unary-tizable.

The Boolean algebras $\mathcal{BA}_{kn}(\mathcal{M})$ and $\mathcal{BA}'_{kn}(\mathcal{M})$ have extensions $\mathcal{BA}_k(\mathcal{M})$ and $\mathcal{BA}'_k(\mathcal{M})$, respectively, consisting of definable sets for n -ary/ n -aritizable formulae with m free variables, for some n . Clearly, both $\mathcal{BA}_k(\mathcal{M})$ and $\mathcal{BA}'_k(\mathcal{M})$ equal the Boolean algebra $\mathcal{B}_k(\mathcal{M})$ of all \emptyset -definable subsets of M^k . Both these inclusions $\mathcal{BA}_{kn}(\mathcal{M}) \subseteq \mathcal{B}_k(\mathcal{M})$ and $\mathcal{BA}'_{kn}(\mathcal{M}) \subseteq \mathcal{B}_k(\mathcal{M})$ can be proper for $k > n$.

It was noticed above that aritizabilities of separated formulae of a theory can not guarantee that witnesses of these aritizabilities produce a n -ary theory, the possibility of *coordinated* expansion of aritizable formulae is necessary. We denote by $\mathcal{BA}''_{kn}(\mathcal{M})$ the Boolean algebra $\mathcal{BA}'_{kn}(\mathcal{M})$ with a coordinated n -ary expansion for all n -aritizable formulae with k free variables.

Using Remark 2.6 we have the following characterizations of n -arity and of n -aritizability of a theory T in terms of Boolean algebras of a model for T .

Proposition 2.7. *For any theory T , its model \mathcal{M} , and $n \in \omega$ the following conditions hold:*

- (1) T is n -ary iff $\mathcal{BA}_{kn}(\mathcal{M}) = \mathcal{B}_k(\mathcal{M})$ for each $k > n$;
- (1) T is n -aritizable iff $\mathcal{BA}''_{kn}(\mathcal{M}) = \mathcal{B}_k(\mathcal{M})$ for each $k > n$.

Definition. [29] The *disjoint union* $\bigsqcup_{n \in \omega} \mathcal{M}_n$ of pairwise disjoint structures \mathcal{M}_n for pairwise disjoint predicate languages Σ_n , $n \in \omega$, is the structure of language $\bigcup_{n \in \omega} \Sigma_n \cup \{P_n^{(1)} \mid n \in \omega\}$ with the universe $\bigsqcup_{n \in \omega} M_n$, $P_n = M_n$, and interpretations of predicate symbols in Σ_n coinciding with their interpretations in \mathcal{M}_n , $n \in \omega$. The *disjoint union of theories* T_n for pairwise disjoint languages Σ_n accordingly, $n \in \omega$, is the theory

$$\bigsqcup_{n \in \omega} T_n \Leftrightarrow \text{Th} \left(\bigsqcup_{n \in \omega} \mathcal{M}_n \right),$$

where $\mathcal{M}_n \models T_n$, $n \in \omega$. Taking empty sets instead of some structures \mathcal{M}_k we obtain disjoint unions of finitely many structures and theories. In particular, we have the disjoint unions $\mathcal{M}_0 \sqcup \dots \sqcup \mathcal{M}_n$ and their theories $T_0 \sqcup \dots \sqcup T_n$.

Clearly, disjoint unions of theories does not depend on choice of correspondent disjoint unions of their models. Besides, disjoint unions $\bigsqcup_{n \in \omega} T_n$ are based by the unions of the basing sets Δ_n for T_n and by the formulae of the form $P_n(x)$. Thus we have the following:

Theorem 2.8. 1. *For any theories T_m , $m \in \omega$, and their disjoint union $\bigsqcup_{m \in \omega} T_m$, all T_m are n -theories iff $\bigsqcup_{m \in \omega} T_m$ is an n -theory, moreover,*

$$\text{ar} \left(\bigsqcup_{m \in \omega} T_m \right) = \max \{ \text{ar}(T_m) \mid m \in \omega \}.$$

2. *For any theories T_m , $m \in \omega$, and their disjoint union $\bigsqcup_{m \in \omega} T_m$, all T_m are n -aritizable iff $\bigsqcup_{m \in \omega} T_m$ is n -aritizable.*

Remark 2.9. The control of the arities with respect to disjoint unions of theories according to Theorem 2.8 shows that the arity of a theory T does not correlate with the degree n of the n -transitivity of automorphism groups of models $\mathcal{M} \models$

T . Indeed, following Proposition 1.6, taking languages with symbol arities $\leq n$ only, n -transitive theories T with quantifier elimination, which are not $(n+1)$ -transitive, satisfy $\text{ar}(T) = n+1$. At the same time, the degree n of the transitivity of automorphism group does not imply the equality $\text{ar}(T) = n+1$ in general since k -ary definable sets, for $k > n$, may be complicated enough, say, containing k -spherical orders for big k , and producing $\text{ar}(T) > n+1$, even $\text{ar}(T) = \infty$. Besides, taking a disjoint union of theories T_i which are marked by unary predicates P_i we loose the transitivity of automorphism group preserving the maximal value of the arity.

Definition [23]. Let \mathcal{M} and \mathcal{N} be structures of relational languages $\Sigma_{\mathcal{M}}$ and $\Sigma_{\mathcal{N}}$ respectively. We define the *composition* $\mathcal{M}[\mathcal{N}]$ of \mathcal{M} and \mathcal{N} satisfying the following conditions:

- 1) $\Sigma_{\mathcal{M}[\mathcal{N}]} = \Sigma_{\mathcal{M}} \cup \Sigma_{\mathcal{N}}$;
- 2) $M[\mathcal{N}] = M \times N$, where $M[\mathcal{N}]$, M , N are universes of $\mathcal{M}[\mathcal{N}]$, \mathcal{M} , and \mathcal{N} respectively;
- 3) if $R \in \Sigma_{\mathcal{M}} \setminus \Sigma_{\mathcal{N}}$, $\mu(R) = n$, then $((a_1, b_1), \dots, (a_n, b_n)) \in R_{\mathcal{M}[\mathcal{N}]}$ if and only if $(a_1, \dots, a_n) \in R_{\mathcal{M}}$;
- 4) if $R \in \Sigma_{\mathcal{N}} \setminus \Sigma_{\mathcal{M}}$, $\mu(R) = n$, then $((a_1, b_1), \dots, (a_n, b_n)) \in R_{\mathcal{M}[\mathcal{N}]}$ if and only if $a_1 = \dots = a_n$ and $(b_1, \dots, b_n) \in R_{\mathcal{N}}$;
- 5) if $R \in \Sigma_{\mathcal{M}} \cap \Sigma_{\mathcal{N}}$, $\mu(R) = n$, then $((a_1, b_1), \dots, (a_n, b_n)) \in R_{\mathcal{M}[\mathcal{N}]}$ if and only if $(a_1, \dots, a_n) \in R_{\mathcal{M}}$, or $a_1 = \dots = a_n$ and $(b_1, \dots, b_n) \in R_{\mathcal{N}}$.

The composition $\mathcal{M}[\mathcal{N}]$ is called *e-definable*, or *equi-definable*, if $\mathcal{M}[\mathcal{N}]$ has an \emptyset -definable equivalence relation E whose E -classes are universes of the copies of \mathcal{N} forming $\mathcal{M}[\mathcal{N}]$. If the equivalence relation E is fixed, the *e-definable* composition is called *E-definable*.

Using a nice basedness of E -definable compositions $T_1[T_2]$ (see [23]) till the formulas of form $E(x, y)$ and generating formulas for T_1 and T_2 we have the following:

Theorem 2.10. 1. *For any theories T_1 and T_2 and their E -definable composition $T_1[T_2]$, T_1 and T_2 are n -theories, for $n \geq 2$, iff $T_1[T_2]$ is an n -theory, moreover, $\text{ar}(T_1[T_2]) = \max\{\text{ar}(T_1), \text{ar}(T_2)\}$, if models of T_1 and of T_2 have at least two elements, and $\text{ar}(T_1[T_2]) = \max\{\text{ar}(T_1), \text{ar}(T_2), 2\}$, if a model of T_1 or T_2 is a singleton.*

2. For any theories T_1 and T_2 and their E -definable composition $T_1[T_2]$, T_1 and T_2 are n -aritzable iff $T_1[T_2]$ is n -aritzable.

Applying Proposition 2.4 and Theorem 2.10 we immediately obtain:

Corollary 2.11. *If each of theories T_1 and T_2 is a theory of a finite structure, or of an infinite structure and n -arizable, then their E -definable composition $T_1[T_2]$ is n -aritzable.*

3. UNARY-TIZABLE, BINARIZABLE AND ARITIZABLE THEORIES, THEIR DEFINABLE SETS AND DYNAMICS

Let T be a theory with a unary expansion T' . Since unary formulas $\varphi(x)$ and $\psi(y)$ have either equal or disjoint free variables we do not have essential mixed sums and mixed products forming definable sets for a model \mathcal{M} of T' , i.e., all definable

sets are formed using unions, intersections, Cartesian sums and Cartesian products of definable subsets of M , without parameters.

Conversely, having a system of definable sets formed by unions, intersections, Cartesian sums and Cartesian products of subsets of M , we can introduce names for these subsets and generate, using this introduced language, all given definable sets.

Thus using Remark 1.5 we obtain the following characterization for the unarity-tizability of a theory in terms of definable sets:

Theorem 3.1. *A theory T is unarity-tizable if and only if for any (some) model \mathcal{M} of T any definable set is formed by unions, intersections, Cartesian sums and Cartesian products of subsets of M .*

Similarly, all definable sets of binarizable theories are generated by unions, intersections, Cartesian sums and Cartesian products of subsets of M^2 , extended by mixed sums and mixed products of these subsets and their combinations:

Theorem 3.2. *A theory T is binarizable if and only if for any (some) model \mathcal{M} of T any \emptyset -definable set is formed by unions, intersections, Cartesian sums, Cartesian products, mixed sums and mixed products of subsets of M^2 .*

By Theorem 3.2 definable sets of binarizable theories are generated by combinations of 3-dimensional cylinders with two-dimensional generators.

Theorems 3.1 and 3.2 admit the following natural generalizations based on $(n + 1)$ -dimensional cylinders with n -dimensional generators.

Theorem 3.3. *A theory T is n -aritzable, for $n \geq 1$, if and only if for any (some) model \mathcal{M} of T any \emptyset -definable set is formed by unions, intersections, Cartesian sums, Cartesian products, mixed sums and mixed products of subsets of M^n .*

Theorem 3.4. *A theory T is aritzable if and only if for any (some) model \mathcal{M} of T any \emptyset -definable set is formed by unions, intersections, Cartesian sums, Cartesian products, mixed sums and mixed products of subsets of M^n , for some n .*

Remark 3.5. Using Theorem 3.3 one can form a definable subset $X \subset M^k$ of an infinite model \mathcal{M} of a n -theory, for $k > n$, such that X has an infinite complement and each projection of X equals M^m for some $m < n$. It implies that Boolean combinations of these projections can not reconstruct X . Thus the structure $\langle M, X \rangle$ has a n -expansive k -theory.

Definition. A T -formula $\varphi(\bar{x})$ is called *constantizable* if T has an expansion T' such that $\varphi(\bar{x})$ is T' -equivalent to a Boolean combination of formulae of forms $x \approx y$ and $x \approx c$ with variables x, y and constants c , i.e., T has an expansion T'' such that $\varphi(\bar{x})$ is T'' -equivalent to a Boolean combination of formulae of forms $x \approx y$ and of formulae of unary singleton predicates whose solutions consist of constants.

A theory T is called *constantizable* if any T -formula $\varphi(\bar{x})$ is constantizable.

By the definition any constantizable theory is unarity-tizable, but not vice versa, as the following assertions show.

Proposition 3.6. *A formula $\varphi(\bar{x})$ of a theory T is constantizable iff $\varphi(\bar{x})$ is T -equivalent to a Boolean combination of formulae of form $x \approx y$ and formulae with finitely many solutions.*

Proof. Let $\varphi(\bar{x})$ be a constantizable formula. Since $\varphi(\bar{x})$ is unary-tizable we can divide $\varphi(\bar{x})$ in some expansion of T onto cases with distinct/equal values for free variables. Thus without loss of generality $\varphi(\bar{x})$ is a Boolean combination of formulae $(x_i \approx x_j)^\delta$ and $\psi(x_i)$, for $x_i, x_j \in \bar{x}$, $i \neq j$, $\delta \in \{0, 1\}$, written in a disjunctive normal form. Since $\varphi(\bar{x})$ is constantizable, its definable set A in a model of T is represented by a Boolean combination of cylinders for $x \approx y$ and $x \approx c$. Thus, $\varphi(\bar{x})$ is T -equivalent to a Boolean combination of formulae of form $x \approx y$ and formulae with finitely many solutions.

Now let $\varphi(\bar{x})$ is T -equivalent to a Boolean combination of formulae of form $x \approx y$ and formulae $\psi(\bar{x})$ with finitely many solutions. We may assume that $\varphi(\bar{x})$ consistent and represented as a disjunctive normal form. We collect in a set Z all finite sets of solutions for formulae $\psi(\bar{x})$. Now for the finite set $\cup Z$ we apply the construction for Proposition 2.4 reducing the set Z' of all coordinates for tuples in $\cup Z$ to unary singleton predicates and so to the formulae $x \approx c$, for $c \in Z'$. It implies that a Boolean combination of these formulae and formulae of the form $x \approx y$ is equivalent to $\varphi(\bar{x})$, i.e., $\varphi(\bar{x})$ is constantizable, as required.

Proposition 3.6 immediately implies:

Corollary 3.7. *A theory T is constantizable iff each T -formula is T -equivalent to a Boolean combination of formulae of form $x \approx y$ and formulae with finitely many solutions.*

Definition [30]. A theory T is called *strongly minimal* if for any formula $\varphi(x, \bar{a})$ of language obtained by adding parameters of \bar{a} (in some model $\mathcal{M} \models T$) to the language of T , either $\varphi(x, \bar{a})$, or $\neg\varphi(x, \bar{a})$ has finitely many solutions.

Using Corollary 3.7 we obtain:

Corollary 3.8. *Any constantizable theory is strongly minimal.*

Now we consider some dynamics of arities of theories under expansions. Since the property of non- n -arizability forbids n -ary expansions it suffices to study possibilities for expansions of n -arizable theories.

Proposition 3.9. *A theory T has a non-arizable expansion iff T has an infinite model.*

Proof. If T has an infinite model there are expansions T' of T collecting, for instance, examples 1.10, 1.11, 1.12 forbidding n -arity for each n . Thus T' is not arizable.

Conversely, if T has a finite model then each expansion T' of T has a finite model producing arizability of T' by Proposition 2.4, as required.

Using examples above we observe that for each natural $n \geq 1$ there are theories T_n with $\text{ar}(T_n) = n$ and finite models. Thus there are theories T_{kn} with $\text{ar}(T_{kn}) = k$ and $\text{ar}(T'_{kn}) = n$ for some expansions T'_{kn} of T_{kn} .

Besides, for each natural $n \geq 1$ there are:

1) theories $T_{n,\infty}$ with $\text{ar}(T_{n,\infty}) = n$ and $\text{ar}(T'_{n,\infty}) = \infty$ for some expansions $T'_{n,\infty}$ of $T_{n,\infty}$: it suffices to expand a n -ary theory with infinite models by new predicates forbidding the k -arizability for each $k > n$;

2) theories $T_{\infty,n}$ with $\text{ar}(T_{\infty,n}) = \infty$ and $\text{ar}(T'_{\infty,n}) = n$ for some expansions $T'_{\infty,n}$ of $T_{\infty,n}$: it suffices to expand an n -arizable theory which is not m -ary for any m till a n -theory.

Thus the arities can be freely increased and decreased and we obtain the following:

Theorem 3.10. *For any $\mu, \nu \in (\omega \setminus \{0\}) \cup \{\infty\}$ there is a theory $T_{\mu, \nu}$ and its expansion $T'_{\mu, \nu}$ such that $\text{ar}(T_{\mu, \nu}) = \mu$ and $\text{ar}(T'_{\mu, \nu}) = \nu$.*

4. CONCLUSION

We considered possibilities for arities of theories and their dynamics, reductions of formulas to ones of special forms as well as definable sets connected with these reductions. Having a definable k -ary relation in n -ary theory, for $k > n$, this relation is reduced to a Boolean combination of n -ary relations. Thus a relation with many coordinates is simplified and reducible to ones with fewer coordinates. It can be used both for databases, simplifying them to ones with bounded dimensions, for geometric objects represented as finite combinations of cylinders, and for cryptographic constructions representing complicated configurations by simpler ones using key leading Boolean combinations for the representations. It would be interesting to describe values of arities and aritizabilities for various additional natural classes of theories.

REFERENCES

- [1] S.V. Sudoplatov, *On a certain complexity estimate in graph theory*, Siberian Mathematical Journal, **37**:3 (1996), 614–671.
- [2] S.V. Sudoplatov, *Transitive arrangements of algebraic systems*, Siberian Mathematical Journal, **40**:6 (1999), 1142–1145.
- [3] A.I. Mal'tsev, *Algebraic systems*, Nauka, Moscow, 1970.
- [4] L. Henkin, J.D. Monk, A. Tarski, *Cylindric Algebras. Part 1*, Amsterdam, London, 1971.
- [5] L. Henkin, J.D. Monk, A. Tarski, *Cylindric Set Algebras*, Lecture Notes in Mathematics. Vol. 883, Springer-Verlag, Berlin, Heidelberg, New York, 1981.
- [6] L. Henkin, J.D. Monk, A. Tarski, *Cylindric Algebras. Part 2*, Studies in Logic and Foundations of Mathematics. Vol. 115, Amsterdam, London, Oxford, 1985.
- [7] T. Imieliński, Jr.W. Lipski, *The relational model of data and cylindric algebras*, Journal of Computer and System Sciences, **28**:1 (1984), 80–102.
- [8] B.Sh. Kulpeshov, H.D. Macpherson, *Minimality conditions on circularly ordered structures*, Mathematical Logic Quarterly, **51**:4 (2005), 377–399.
- [9] B.S. Baizhanov, B.Sh. Kulpeshov, *On behaviour of 2-formulas in weakly o-minimal theories*, Mathematical Logic in Asia, Proceedings of the 9th Asian Logic Conference. Eds.: S. Goncharov, R. Downey, H. Ono, World Scientific: Singapore, 2006, 31–40.
- [10] B.Sh. Kulpeshov, *Criterion for binarity of \aleph_0 -categorical weakly o-minimal theories*, Annals of Pure and Applied Logic, **45** (2007), 354–367.
- [11] A.B. Altaeva, B.Sh. Kulpeshov, *On almost binary weakly circularly minimal structures*, Bulletin of Karaganda University, Mathematics, **78**:2 (2015), 74–82.
- [12] B.Sh. Kulpeshov, *On almost binarity in weakly circularly minimal structures*, Eurasian Mathematical Journal, **7**:2 (2016), 38–49.
- [13] A.B. Altaeva, B.Sh. Kulpeshov, *Binarity of almost ω -categorical quite o-minimal theories*, Siberian Mathematical Journal, **61**:3 (2020), 379–390.
- [14] B.Sh. Kulpeshov, *A criterion for binarity of almost ω -categorical weakly o-minimal theories*, Siberian Mathematical Journal, **62**:6 (2021), 1063–1075.
- [15] S.V. Sudoplatov, *Classification of Countable Models of Complete Theories*, NSTU, Novosibirsk, 2018.
- [16] I.V. Shulepov, S.V. Sudoplatov, *Algebras of distributions for isolating formulas of a complete theory*, Siberian Electronic Mathematical Reports, **11** (2014), 380–407.
- [17] B.Sh. Kulpeshov, S.V. Sudoplatov, *On algebras of distributions of binary formulas for quite o-minimal theories*, News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-Mathematical Series, **300**:2 (2015),
- [18] D.Yu. Emelyanov, *On algebras of distributions of binary formulas for theories of unars*, Bulletin of Irkutsk State University. Series Mathematics, **17** (2016), 23–36.

- [19] D.Yu. Emelyanov, B.Sh. Kulpeshov, S.V. Sudoplatov, *Algebras of distributions for binary formulas in countably categorical weakly o-minimal structures*, Algebra and Logic, **56**:1 (2017), 13–36.
- [20] D.Yu. Emelyanov, S.V. Sudoplatov, *On deterministic and absorbing algebras of binary formulas of polygonometrical theories*, Bulletin of Irkutsk State University. Series Mathematics, **20** (2017), 32–44.
- [21] K.A. Baikalova, D.Yu. Emelyanov, B.Sh. Kulpeshov, E.A. Palyutin, S.V. Sudoplatov, *On algebras of distributions of binary isolating formulas for theories of abelian groups and their ordered enrichments*, Russian Mathematics, **62**:4 (2018), 1–12.
- [22] D.Yu. Emelyanov, B.Sh. Kulpeshov, S.V. Sudoplatov, *On algebras of distributions for binary formulas for quite o-minimal theories*, Algebra and Logic, **57**:6 (2018), 429–444.
- [23] D.Yu. Emelyanov, B.Sh. Kulpeshov, S.V. Sudoplatov, *Algebras of binary formulas for compositions of theories*, Algebra and Logic, **59**:4 (2020), 295–312.
- [24] E.A. Palyutin, J. Saffe, S.S. Starchenko, *Models of superstable Horn theories*, Algebra and Logic, **24**:3 (1985), 171–210.
- [25] S.V. Sudoplatov, *Syntactic approach to constructions of generic models*, Algebra and Logic, **46**:2 (2007), 134–146.
- [26] S.V. Sudoplatov, *Basedness of stable theories and properties of countable models with powerful types*, Dis... cand. fiz.-mat. sc.: 01.01.06; Novosibirsk, 1990.
- [27] S. Shelah, *Classification theory and the number of non-isomorphic models*, North-Holland, Amsterdam, 1990.
- [28] V. Harnik, L. Harrington, *Fundamentals of forking*, Annals of Pure and Applied Logic, **26**:3 (1984), 245–286.
- [29] R.E. Woodrow, *Theories with a finite number of countable models and a small language*, Ph. D. Thesis, Simon Fraser University, 1976.
- [30] J.T. Baldwin, A. Lachlan, *On strongly minimal sets*, Journal of Symbolic Logic, **36**:1 (1971), 79–96.

SERGEY VLADIMIROVICH SUDOPLATOV
SOBOLEV INSTITUTE OF MATHEMATICS
ACADEMICIAN KOPTYUG AVENUE, 4
630090, NOVOSIBIRSK, RUSSIA.
E-mail address: sudoplat@math.nsc.ru

NOVOSIBIRSK STATE TECHNICAL UNIVERSITY
K. MARX AVENUE, 20
630073, NOVOSIBIRSK, RUSSIA.
E-mail address: sudoplatov@corp.nstu.ru