

СИБИРСКИЕ ЭЛЕКТРОННЫЕ  
МАТЕМАТИЧЕСКИЕ ИЗВЕСТИЯ

Siberian Electronic Mathematical Reports

<http://semr.math.nsc.ru>

---

---

Том 12, стр. 144–144 (2015)  
DOI 10.17377/semi.2015.12.xxx

УДК 512.554.7  
MSC 17C70

**DUAL COALGEBRA OF THE DIFFERENTIAL POLYNOMIAL  
ALGEBRA IN ONE VARIABLE AND RELATED COALGEBRAS**

V. N. ZHELYABIN, P.S. KOLESNIKOV

**ABSTRACT.** We show that the dual coalgebra of the polynomial algebra in one variable is the space of linearly recursive sequences over an arbitrary field. Moreover, this coalgebra is a differential one relative to the dual standard derivation and does not contain nonzero finite-dimensional differentially closed subcoalgebras if the characteristic of the ground field is zero. We construct a Novikov coalgebra which is the dual coalgebra of the left-symmetric Witt algebra of index one. Also, we construct a Jordan supercoalgebra which is dual to the Jordan superalgebra of vector type of the polynomial algebra in one variable. All these coalgebras do not contain non-zero finite-dimensional subcoalgebras if the characteristic of ground field is zero. It is shown that over a field of characteristic different from 2 the adjoint Lie coalgebra of the dual coalgebra of the left-symmetric Witt algebra of index one is isomorphic to the dual coalgebra of the Witt algebra of index one.

**Key words:** coalgebra, coderivation, associative commutative algebra, differential algebra, Novikov algebra, Lie algebra, Witt algebra, Jordan superalgebra, locally finite coalgebra

In general, an algebra is a vector space  $A$  over a ground field  $F$  equipped with a linear map  $m : A \otimes A \rightarrow A$ . The dual notion is known as a coalgebra. The theory of coalgebras has been initially developed within the framework of the theory of Hopf algebras [1]. The main result of the theory of associative coalgebras is the Fundamental Theorem on Coalgebras, which asserts that every associative coalgebra over a field is locally finite. The latter means that every finitely generated coalgebra is finite-dimensional.

---

ZHELYABIN, V.N., KOLESNIKOV, P.S. DUAL COALGEBRA OF THE DIFFERENTIAL POLYNOMIAL ALGEBRA IN ONE VARIABLE AND RELATED COALGEBRAS.

© 2022 ZHELYABIN V.N., KOLESNIKOV P.S. .

Supported by RAS Fundamental Research Program, project FWNF-2022-0002 .

Поступила 1 января 2015 г., опубликована 31 декабря 2015 г.

The interest to nonassociative coalgebras is related to the notion of a quantum group introduced by Drinfeld [2]. Initially, it was associated with the notion of a Lie bialgebra. The latter was introduced in [3] as one of the most important notions of quantum group theory. A Lie bialgebra is simultaneously a Lie algebra and a Lie coalgebra. In contrast to associative coalgebras there exist Lie coalgebras that are not locally finite [4].

It is known (see [1, 4]) that the dual algebra of an associative coalgebra or a Lie coalgebra is associative or Lie, respectively. In [5] (1994), the notion of a coalgebra related to some variety of algebras was introduced. In particular, alternative and Jordan coalgebras were defined, and their local finiteness was proved. An analogue of this result is true for structurable coalgebras [6], for Jordan copairs [7], for right alternative Malcev admissible coalgebras, and for binary  $(-1, 1)$ -coalgebras [8]. In [9] (1995), some necessary and sufficient conditions for a Lie coalgebra to be locally finite were found.

In the papers [10, 11] the first author found the connection between Jordan and Lie (super)coalgebras, which is an analogue of the well-known Kantor–Koecher–Tits construction for usual (super)algebras. In [12, 13], it was shown that every Malcev coalgebra is embedded into a Lie coalgebra with triality.

As mentioned above, there exist non-locally finite Lie coalgebras. In [14], an example of a non-locally finite right-symmetric coalgebra was constructed. In contrast to Jordan coalgebras, non-locally finite Jordan supercoalgebras exist [11].

In [15], it was constructed an example of a non-locally finite differential coalgebra. On this differential coalgebra we can define a Lie comultiplication so that the obtained Lie coalgebra coincides with the example of Michaelis presented in [4]. The dual analogue of the Gelfand–Dorfman construction was proposed in [15], which implied the construction of Novikov coalgebras based on differential associative commutative coalgebras. Using this construction, an example of a non-locally finite Novikov coalgebra was built. A dual analogue of the Kantor construction for usual Jordan superalgebra was also presented in [15], as a corollary, a new example of a non-locally finite Jordan supercoalgebra was constructed.

In [15] and [16], examples non-locally finite right alternative coalgebras were constructed.

In [17], it was shown that the dual coalgebra  $W_1^\circ$  of the Witt Lie algebra  $W_1$  is a non-zero Lie coalgebra which does not contain non-zero finite-dimensional subcoalgebras. An analogue of this result for Jordan supercoalgebra was obtained in [21]. The structure of the dual Lie coalgebra of the Witt algebra over field of characteristic not 2 and zero was also described in [18, 19]. These results were generalized to the case of several derivations in [20]. Namely in [18] was shown that  $W_1^\circ$  is the space of linearly recursive sequences, if  $\text{char } F = 0$  or  $\text{char } F = p > 2$ .

It is known that both the Witt Lie algebra  $W_1$  and the left-symmetric algebra  $\mathcal{L}_1$  can be obtained from the differential polynomial algebra in one variable by means of appropriate constructions. It follows from [19, Theorem 1] that the dual coalgebra  $W_1^\circ$  is obtained from  $P_1^\circ$  by means of the dual construction. The purpose of this paper is to obtain the dual analogue of these results for the dual coalgebra  $\mathcal{L}_1^\circ$ .

In particular, we prove that the dual coalgebra  $P_1^\circ$  of the algebra  $P_1$  in one variable is the space of linearly recursive sequences over an arbitrary field. Moreover, if the characteristic of ground field is zero then the coalgebra  $P_1^\circ$  does not contain finite-dimensional differentially closed subcoalgebras.

We also show that over field of characteristic different from 2 the dual coalgebra of the left-symmetric Witt algebra of index one can be obtained from the coalgebra  $P_1^\circ$  by means of the dual Gelfand–Dorfman construction.

Finally, we describe the dual supercoalgebra of a Jordan superalgebra obtained by the Kantor construction from the differential polynomial algebra in one variable.

### 1. COALGEBRAS AND CODERIVATIONS

Let  $F$  be an arbitrary field. Denote by  $\underbrace{V \otimes \dots \otimes V}_n$  the  $n$ th tensor power of the vector space  $V$  over  $F$ . Denote by  $V^*$  the dual vector space of  $V$ , i. e.,  $V^* = \text{Hom}_F(V, F)$ .

The map

$$\rho : \underbrace{V^* \otimes \dots \otimes V^*}_n \rightarrow \underbrace{(V \otimes \dots \otimes V)^*}_n$$

defined by

$$\rho(f_1 \otimes \dots \otimes f_n) \left( \sum_{i_1, \dots, i_n} v_{i_1} \otimes \dots \otimes v_{i_n} \right) = \sum_{i_1, \dots, i_n} f_1(v_{i_1}) \dots f_n(v_{i_n})$$

is injective. Therefore, we can assume that

$$\underbrace{V^* \otimes \dots \otimes V^*}_n \subseteq \underbrace{(V \otimes \dots \otimes V)^*}_n.$$

If  $\phi : V \rightarrow U$  is a linear map of vector spaces then the transpose  $\phi^* : U^* \rightarrow V^*$  of  $\phi$  is defined by the rule  $\phi^*(u^*)(v) = u^*(\phi(v))$ , where  $v \in V$ ,  $u^* \in U^*$ .

**Definition 1.** A pair  $(C, \Delta)$ , where  $C$  is a vector space over  $F$  and  $\Delta : C \rightarrow C \otimes C$  is a linear map, is called a coalgebra. The map  $\Delta$  is said to be a comultiplication of  $C$ . For an element  $a \in C$ , we will use the Sweedler's notation (see [1]), namely,  $\Delta(a) = \sum_{(a)} a_{(1)} \otimes a_{(2)}$ .

If  $(C, \Delta)$  is a coalgebra then the rule  $f \otimes g \mapsto fg$ , where

$$(fg)(a) = \rho(f \otimes g)(\Delta(a)) = \sum_{(a)} f(a_{(1)})g(a_{(2)}), \quad f, g \in C^*, \quad a \in C,$$

defines a product on  $C^*$ , so that  $C^*$  is an algebra. The algebra  $C^*$  is called the *dual algebra* of  $(C, \Delta)$ .

The dual algebra  $C^*$  has natural left and right actions on the initial coalgebra  $C$ . Namely,

$$\alpha \cdot a = \sum_{(a)} a_{(1)}\alpha(a_{(2)}), \quad a \cdot \alpha = \sum_{(a)} \alpha(a_{(1)})a_{(2)},$$

for  $\alpha \in C^*$ ,  $a \in C$ . Hence,  $C$  is a  $C^*$ -bimodule.

A linear map  $d : C \rightarrow C$  is called a *coderivation* of a coalgebra  $(C, \Delta)$  if

$$\Delta d = (d \otimes id + id \otimes d)\Delta,$$

i. e., for every  $a \in C$

$$\Delta(d(a)) = \sum_{(a)} d(a_{(1)}) \otimes a_{(2)} + a_{(1)} \otimes d(a_{(2)}).$$

The following statement is well-known.

**Lemma 1.** *Let  $d$  be a coderivation of a coalgebra  $(C, \Delta)$ . Then its transpose map  $d^*$  is a derivation of the dual algebra  $C^*$ , i. e.,*

$$d^*(fg) = d^*(f)g + fd^*(g)$$

*holds in the algebra  $C^*$  for all  $f, g \in C^*$ .*

A coalgebra equipped with a coderivation is called a *differential coalgebra*.

A coalgebra  $(C, \Delta)$  is said to be associative (coassociative) if

$$(\Delta \otimes id - id \otimes \Delta)\Delta = 0,$$

i. e., for every  $a \in C$

$$\sum_{(a)} (\Delta(a_{(1)}) \otimes a_{(2)} - a_{(1)} \otimes \Delta(a_{(2)})) = 0.$$

It is known that a coalgebra  $(C, \Delta)$  is associative if and only if its dual algebra  $C^*$  is associative. By this reason, the following definition of a coalgebra related to some variety of algebras was stated in [5].

**Definition 2.** *Let  $\mathcal{M}$  be a variety of algebras. Then a coalgebra  $(C, \Delta)$  is called an  $\mathcal{M}$ -coalgebra if its dual algebra  $C^*$  is an algebra of  $\mathcal{M}$ .*

Let  $V$  be a vector space, and let the linear map  $\tau : V \otimes V \rightarrow V \otimes V$  is defined by  $\tau(x \otimes y) = y \otimes x$ ,  $x, y \in V$ .

A coalgebra  $(C, \Delta)$  is commutative (cocommutative) if

$$\Delta = \tau\Delta,$$

i. e.,

$$\sum_{(a)} a_{(1)} \otimes a_{(2)} = \sum_{(a)} a_{(2)} \otimes a_{(1)}$$

for every  $a \in C$ .

Let  $(C, \Delta)$  be an arbitrary coalgebra. A vector subspace  $B$  of  $C$  is a *subcoalgebra* of  $(C, \Delta)$  if  $\Delta(B) \subseteq B \otimes B$ .

It is known (see [5]) that a vector space  $B$  of  $C$  is a subcoalgebra if and only if  $B$  is a submodule of the  $C^*$ -bimodule  $C$ . Therefore, the intersection of a family of subcoalgebras of  $(C, \Delta)$  is again a subcoalgebra of  $C$ .

Recall that the orthogonal complement

$$B^\perp = \{\alpha \in C^* \mid B \subseteq \ker \alpha\}$$

of a subcoalgebra  $B$  of  $(C, \Delta)$  is an ideal of the algebra  $C^*$ . Conversely, the orthogonal complement  $I^\perp$  of an ideal  $I$  of  $C^*$  is a subcoalgebra of  $(C, \Delta)$  (see [1, Proposition 1.4.9]).

Let  $S$  be a subset of  $C$ . The smallest subcoalgebra which contains  $S$  is called the *subcoalgebra generated by  $S$*  and denoted by  $Coalg(S)$ . In other words,  $Coalg(S)$  is the submodule of the  $C^*$ -bimodule  $C$ , generated by  $S$ . If  $S$  is a finite set then  $Coalg(S)$  is called finitely generated coalgebra.

**Definition 3.** *A coalgebra  $(C, \Delta)$  is called locally finite if every finitely generated subcoalgebra of  $C$  is finite-dimensional.*

Let  $(C, \Delta)$  be a coalgebra. Denote by  $Loc(C)$  the sum of all locally finite subcoalgebras of  $C$ . It is clear that  $Loc(C)$  is a locally finite coalgebra.

Let  $A$  be an algebra over a field  $F$  with a multiplication  $m : A \otimes A \rightarrow A$ , i. e.,  $m(a \otimes b) = ab$  for  $a, b \in A$ . Then  $m^* : A^* \rightarrow (A \otimes A)^*$  is the transpose map of  $m$ . A vector subspace  $V$  of  $A^*$  is called *good* if  $m^*(V) \subseteq \rho(V \otimes V)$ . If  $V$  is a good subspace then we can define the comultiplication  $\Delta_V$  on  $V$  by the rule  $\Delta_V = \rho^{-1}m^*$ .

Let  $A^\circ$  be the sum of all good subspaces of  $A^*$ . Then  $A^\circ$  is the largest good subspace of  $A^*$ , and hence the pair  $(A^\circ, \Delta^\circ)$  is a coalgebra with the comultiplication  $\Delta^\circ = \Delta_{A^\circ}$  [4, 5]. The coalgebra  $(A^\circ, \Delta^\circ)$  is called the *finite dual coalgebra* (or simply *dual coalgebra*) of  $A$ . For every  $a, b \in A$  and for every  $f \in A^\circ$  we have  $f(ab) = \sum_f f_{(1)}(a)f_{(2)}(b)$ , where  $\Delta^\circ(f) = \sum_f f_{(1)} \otimes f_{(2)}$ .

If  $A$  is finite dimensional then  $A^* \otimes A^* \cong \rho(A^* \otimes A^*) = (A \otimes A)^*$  and the dual space  $A^*$  is a coalgebra with the comultiplication  $\Delta = \rho^{-1}m^*$ . Therefore,  $(A^\circ, \Delta^\circ) = (A^*, \Delta)$ .

In [5], it was shown that the dual coalgebra  $A^\circ$  is a  $\mathcal{M}$ -coalgebra if  $A$  is an algebra of variety  $\mathcal{M}$ .

**Lemma 2.** *Let  $A$  be an algebra over a field  $F$ , and let  $(V, \Delta)$  be a coalgebra, where  $V \subseteq A^*$ . If  $f(ab) = \rho(\Delta(f))(a \otimes b)$  holds for every  $a, b \in A$  and for every  $f \in V$  then  $V$  is a good subspace of  $A^*$ .*

*Proof.* By the definition of  $m^*$ , we have  $m^*(f)(a \otimes b) = f(ab)$  for all  $a, b \in A$ ,  $f \in A^*$ . If  $f(ab) = \rho(\Delta(f))(a \otimes b)$  holds for  $f \in V$  then

$$m^*(f)(a \otimes b) = \rho(\Delta(f))(a \otimes b).$$

Consequently,  $m^*(f) = \rho(\Delta(f))$ . Thus  $m^*(V) \subseteq \rho(V \otimes V)$ .  $\square$

Let  $\mathcal{I}$  be the set of finite-codimensional ideals of an algebra  $A$ . Let us define

$$A_{\mathcal{I}}^* = \{\alpha \in A^* \mid \text{exists } I \in \mathcal{I} \text{ such that } I \subseteq \ker \alpha\}.$$

As it was shown in [5],  $A_{\mathcal{I}}^* = Loc(A^\circ)$ . It is known that  $A_{\mathcal{I}}^* = Loc(A^\circ) = A^\circ$  for every associative algebra  $A$ .

## 2. DUAL COALGEBRA OF THE DIFFERENTIAL POLYNOMIAL ALGEBRA $F[x]$

Let  $P_1 = F[x]$  be the polynomial algebra in one variable equipped with the standard derivation  $d = \frac{d}{dx}$ . Then

$$P_1^\circ = \{\alpha \in P_1^* \mid \text{exists } f(x) \in F[x], f(x) \neq 0 \text{ such that } f(x)F[x] \subseteq \ker \alpha\}.$$

Let  $\Delta^\circ = \Delta_{P_1^\circ}$ . For  $\alpha \in P_1^*$  and  $a \in P_1$  put  $\langle \alpha, a \rangle = \alpha(a)$ . Also, let us denote  $d^\circ = d^*|_{P_1^\circ}$ , where  $d^*$  is the transpose map of the derivation  $d$ . Let  $(P_1, d)^\circ$  be the sum of all good subspaces of  $P_1^*$  which are differential coalgebras with the coderivation  $d^\circ$ .

**Lemma 3.** *The inclusion  $d^\circ(P_1^\circ) \subseteq P_1^\circ$  holds. The triple  $(P_1^\circ, \Delta^\circ, d^\circ)$  is an associative and commutative differential coalgebra with the coderivation  $d^\circ$ . Moreover,  $(P_1^\circ, \Delta^\circ, d^\circ) = (P_1, d)^\circ$ .*

*Proof.* Let us show that  $d^\circ(P_1^\circ) \subseteq P_1^\circ$ . Suppose  $u \in P_1^\circ$ , then there exists  $I = f(x)P_1$ ,  $f(x) \neq 0$ , such that  $\langle u, I \rangle = 0$ . Consider  $J = I^2$ : this is an ideal of finite codimension in  $P_1$ , and  $\langle d^*(u), J \rangle = \langle u, d(J) \rangle \subseteq \langle u, I \rangle = 0$  since  $d(I^2) \subseteq Id(I) \subseteq I$ .

Obviously,  $(P_1^\circ, \Delta^\circ, d^\circ)$  is a differential coalgebra with coderivation  $d^\circ$ . Hence,  $(P_1^\circ, \Delta^\circ, d^\circ) = (P_1, d)^\circ$ .  $\square$

**Remark 1.** *The proof of Lemma 3 remains valid for an arbitrary associative commutative algebra  $A$  equipped with an arbitrary derivation  $D$ . Hence, the equality  $(A^\circ, \Delta^\circ, D^\circ) = (A, D)^\circ$  holds in general.*

**Theorem 1.** *Let  $F$  be a field of characteristic zero. Then the coalgebra  $(P_1^\circ, \Delta^\circ, d^\circ)$  does not contain non-zero finite-dimensional differentially closed subcoalgebras.*

*Proof.* Let  $B$  be a finite-dimensional differentially closed subcoalgebra of the differential coalgebra  $(P_1^\circ, \Delta^\circ, d^\circ)$ . Then the orthogonal complement  $B^\perp$  is a differentially closed ideal of  $P_1$ . Moreover, the ideal  $B^\perp$  has finite codimension. Since the characteristic of  $F$  is zero then  $P_1$  is a differentially simple algebra. Therefore, either  $B^\perp = 0$  or  $B^\perp = P_1$ . The first option is impossible since  $P_1$  is an infinite-dimensional algebra. Consequently,  $B^\perp = P_1$  and  $B = 0$ .  $\square$

Let  $W_1$  be the Witt algebra of index one, i. e.,  $W_1 = \text{Der}_F(P_1)$ ;  $P_1^*$  and  $W_1^*$  are isomorphic as vector spaces.

Recall that the vector space  $P_1$  is a Lie algebra relative to the operation

$$[f, g]_d = fd(g) - d(f)g,$$

this Lie algebra is isomorphic to the Witt algebra  $W_1$  under isomorphism  $f \mapsto fd$ . Thus, we can identify  $(P_1, [, ]_d)$  and  $W_1$ .

Define the new comultiplication  $\Delta_{d^\circ}^{(-)}$  on the vector space  $P_1^\circ$ :

$$\Delta_{d^\circ}^{(-)}(\alpha) = (id \otimes d^\circ - d^\circ \otimes id)\Delta^\circ(\alpha).$$

Then

$$\langle \alpha, [f, g]_d \rangle = \langle \alpha, fd(g) - d(f)g \rangle = \rho(\Delta_{d^\circ}^{(-)}(\alpha))(f \otimes g)$$

for  $\alpha \in P_1^\circ$  and  $f, g \in P_1$ . Therefore,  $P_1^\circ$  is a good subspace of  $W_1^*$ . Consequently, we can assume that  $(P_1^\circ, \Delta_{d^\circ}^{(-)})$  is a subcoalgebra of  $(W_1^\circ, \Delta_{W_1^\circ})$ .

In the algebra  $P_1$  we put  $x_i = x^{i+1}$ ,  $i = -1, 0, 1, \dots$ . Then we have  $x_i \cdot x_j = x_{i+j+1}$  in the algebra  $P_1$ .

We identify elements of  $P_1^*$  with sequences of elements of  $F$ . Namely, every  $\alpha \in P_1^*$  corresponds to  $(\alpha(x_n))_{n \geq -1}$ .

Following [18], we say that a sequence  $(a_n)_{n \geq -1}$  of elements of  $F$  is  $(F)$ -linearly recursive if there exist  $\beta_0, \beta_1, \dots, \beta_r \in F$ , not all equal to zero, and a number  $k$  such that  $\sum_{i=0}^r \beta_i a_{n+i} = 0$  for all  $n \geq k$ .

In the next statement, the base field  $F$  is of arbitrary characteristic.

**Theorem 2.** *Let  $V$  be the space of linearly recursive sequences. Then  $P_1^\circ = V$ . In particular, the space  $V$  is the dual coalgebra of  $P_1$ . Therefore, the coalgebra  $(V, \Delta_{d^\circ}^{(-)})$  is a Lie subcoalgebra of  $(W_1)^\circ$ .*

*Proof.* Let  $\alpha \in P_1^*$  and  $a_n = \alpha(x_n)$ , where  $n \geq -1$ .

Assume  $\alpha \in P_1^\circ$ . We show that the sequence  $(a_n)_{n \geq -1}$  is linearly recursive. There exists  $f(x) \in P_1$ ,  $f(x) \neq 0$ , such that  $\alpha(f(x)g(x)) = 0$  for all  $g(x) \in P_1$ . Let  $f(x) = \beta_r x_{r-1} + \dots + \beta_0 x_{-1}$ , where  $\beta_0, \beta_1, \dots, \beta_r \in F$ . Since  $f(x) \neq 0$ , not all  $\beta_0, \beta_1, \dots, \beta_r$  are zero. Then for all  $n \geq 0$  we have

$$\sum_{i=0}^r \beta_i a_{n+i} = \sum_{i=0}^r \beta_i \alpha(x_{n+i}) = \sum_{i=0}^r \beta_i \alpha(x^{n+i+1}) = \alpha(f(x)x^{n+1}) = 0.$$

Hence, the sequence  $(a_n)_{n \geq -1}$  is linearly recursive.

Let  $(a_n)_{n \geq -1}$  be a linearly recursive sequence. Then there exist  $\beta_0, \beta_1, \dots, \beta_r \in F$ , not all zero, and a number  $k$ , such that  $\sum_{i=0}^r \beta_i a_{n+i} = 0$  for all  $n \geq k$ . Put  $f(x) = \beta_r x_{k+r} + \dots + \beta_0 x_k$ . Then for all  $j \geq -1$  we get

$$\alpha(f(x)x_j) = \sum_{i=0}^r \beta_i \alpha(x_{k+i+j+1}) = \sum_{i=0}^r \beta_i a_{k+i+j+1} = 0,$$

since  $k+j+1 \geq k$ . Consequently,  $f(x)P_1 \subseteq \ker \alpha$ .

Since  $(P_1^\circ, \Delta_{d^\circ}^{(-)})$  is a subcoalgebra of  $(W_1^\circ, \Delta_{W_1^\circ})$  then the space  $V$  is a Lie subcoalgebra of  $(W_1)^\circ$ .  $\square$

By [18, Theorem 5], if  $F$  has characteristic 0 or characteristic  $p \neq 2$ , then  $W_1^\circ$  is the space of linearly recursive sequences. Hence,  $P_1^\circ = W_1^\circ$  by Theorem 2.

Therefore, the following statement holds.

**Corollary 1** (see also [19]). *Define the comultiplication*

$$\Delta_{d^\circ}^{(-)}(\alpha) = (id \otimes d^\circ - d^\circ \otimes id)\Delta^\circ(\alpha).$$

on the vector space  $P_1^\circ$ . If  $F$  is a field of characteristic  $\neq 2$  then  $(P_1^\circ, \Delta_{d^\circ}^{(-)}) = (W_1^\circ, \Delta_{W_1^\circ})$ .

Following [17], we define elements  $y_n$ , of  $W_1^*$ ,  $n \geq -1$ , by  $y_n(x_i) = \delta_{n,i}$ . Define  $Y = span(y_n \mid n \geq -1)$ , and let  $\Delta_Y$  be the restriction of  $m^*$  in  $P_1^*$  on the space  $Y$ . Then

$$\Delta_Y(y_n) = \sum_{i+j=n-1} y_i \otimes y_j.$$

By Lemma 2,  $Y$  is a good subspace of  $P_1^*$ . Consequently,  $(Y, \Delta_Y)$  is a subcoalgebra of  $(P_1^\circ, \Delta^\circ)$ .

Now following [18], define  $y_{a,j} \in P_1^*$ , for  $0 \neq a \in F$ ,  $j \geq -1$ , by putting  $y_{a,j}(x_i) = a^i \binom{i+1}{j+1}$ . Define  $Y_a = span(y_{a,j} \mid j \geq -1)$ , and let  $\Delta_{Y_a}$  be the restriction of  $m^*$  in  $P_1^*$  on the space  $Y_a$ . For the sake of uniformity, we set  $Y_0 = Y$ . Let us turn  $Y_a$  into a differential subcoalgebra of  $P_1^\circ$ .

**Lemma 4.** *The space  $Y_a$  is a good subspace of  $P_1^*$ , and  $(Y_a, \Delta_{Y_a}, d^\circ)$  is a differential coalgebra. Consequently,  $(Y_a, \Delta_{Y_a}, d^\circ) \subseteq (P_1, d)^\circ$ . If  $F$  is an algebraically closed field of characteristic zero then  $P_1^\circ = \bigoplus_{a \in F} Y_a$*

*Proof.* First let  $a = 0$ . We prove that  $d^\circ(y_k) = (k+2)y_{k+1}$  for  $k \geq -1$ . Indeed, if  $i > -1$  then

$$d^\circ(y_k)(x_i) = y_k(d((x_i))) = (i+1)y_k(x_{i-1}) = (i+1)y_{k+1}(x_i) = (k+2)y_{k+1}(x_i).$$

If  $i = -1$  then  $d^\circ(y_k)(x_{-1}) = y_k(d((1))) = 0$ . On the other hand  $y_{k+1}(x_{-1}) = \delta_{k+1,-1} = 0$ . Hence,  $d^\circ(y_k)(x_{-1}) = (k+2)y_{k+1}(x_{-1})$ .

It is clearly that  $d^\circ$  is a coderivation of  $(Y, \Delta_Y)$ . Therefore,  $(Y, \Delta_Y, d^\circ)$  is a good subspace of  $(P_1, d)^*$ . Consequently,  $(Y, \Delta_Y, d^\circ) \subseteq (P_1, d)^\circ$ .

Let  $0 \neq a \in F$ . Recall that for binomial coefficients we have

$$\binom{r+s+2}{k+1} = \sum_{i+j=k-1, i,j \geq -1} \binom{r+1}{i+1} \binom{s+1}{j+1},$$

where  $r, s, k \geq -1$ .

The comultiplication  $\Delta_{Y_a}$  on the space  $Y_a$  is given by

$$\Delta_{Y_a}(y_{a,n}) = \sum_{i+j=n-1, i,j \geq -1} ay_{a,i} \otimes y_{a,j}.$$

Indeed,

$$\begin{aligned} \rho(\Delta_{Y_a}(y_{a,k}))(x_r \otimes x_s) &= \sum_{i+j=k-1, i,j \geq -1} ay_{a,i}(x_r)y_{a,j}(x_s) = \\ &= \sum_{i+j=k-1, i,j \geq -1} a^{r+s+1} \binom{r+1}{i+1} \binom{s+1}{j+1} = a^{r+s+1} \binom{r+s+2}{k+1} = \\ &= y_{a,k}(x_{r+s+1}) = y_{a,k}(x_r \cdot x_s). \end{aligned}$$

Therefore,  $Y_a$  is a good subspace of  $P_1^*$ . Consequently,  $(Y_a, \Delta_{Y_a})$  is a subcoalgebra of  $(P_1^\circ, \Delta^\circ)$ .

Let us show that for the coderivation  $d^\circ$  we have  $d^\circ(y_{a,k}) = a^{-1}(k+2)y_{a,k+1}$ . Indeed, if  $i > -1$  then

$$\begin{aligned} d^\circ(y_{a,k})(x_i) &= y_{a,k}(d(x_i)) = (i+1)y_{a,k}(x_{i-1}) = a^{i-1}(i+1) \binom{i}{k+1} = \\ &= a^{i-1}(i+1) \frac{i!}{(k+1)!(i-k-1)!} = a^{-1}(k+2)a^i \frac{(i+1)!}{(k+2)!(i+1-k-2)!} = \\ &= a^{-1}(k+2)y_{a,k+1}(x_i). \end{aligned}$$

If  $i = -1$  then  $d^\circ(y_{a,k})(x_{-1}) = y_{a,k}(d(1)) = 0$ . On the other hand,  $y_{a,k+1}(x_{-1}) = a^{-1} \binom{0}{k+2} = 0$ . Therefore,  $d^\circ(y_{a,k})(x_{-1}) = a^{-1}(k+2)y_{a,k+1}(x_{-1})$ .

Consequently,  $(Y_a, \Delta_{Y_a}, d^\circ)$  is a good subspace of  $(P_1, d)^*$ , so  $(Y_a, \Delta_{Y_a}, d^\circ) \subseteq (P_1, d)^\circ$ .

If  $F$  is an algebraically closed field of characteristic zero then Theorems 4 and 5 from [18] imply  $W_1^\circ = \bigoplus_{a \in F} Y_a$ . Hence,  $P_1^\circ = \bigoplus_{a \in F} Y_a$ .  $\square$

An algebra  $(A, \circ)$  with a multiplication operation  $\circ$  is called a *Novikov algebra* (see [22, 23]) if  $A$  satisfies the following identities:

$$x \circ (y \circ z) - (x \circ y) \circ z = y \circ (x \circ z) - (y \circ x) \circ z \text{ (left symmetry),}$$

$$(x \circ y) \circ z = (x \circ z) \circ y \text{ (right commutativity).}$$

Define the operation of multiplication  $\circ$  on  $P_1$  by the rule  $f \circ g = fd(g)$ ,  $f, g \in P_1$ . Then  $(P_1, \circ)$  is a Novikov algebra (see [22]). Following [24], we will call it left-symmetric Witt algebra of index 1 and denote by  $\mathcal{L}_1$ . Let  $x_i = x^{i+1}$ ,  $i = -1, 0, 1, \dots$ . Then the equality  $x_i \circ x_j = (j+1)x_{i+j}$  holds in the algebra  $\mathcal{L}_1$ . It is known that the algebra  $\mathcal{L}_1$  is simple, if  $F$  is field of characteristic zero.

Define the comultiplication  $\Delta_N$  on  $P_1^\circ$  by the rule

$$\Delta_N(\alpha) = (id \otimes d^\circ)\Delta^\circ(\alpha).$$

As it was shown in [15], the coalgebra  $(P_1^\circ, \Delta_N)$  is a Novikov coalgebra. Obviously,  $P_1^\circ$  is a good subspace of  $\mathcal{L}_1^*$ . Therefore,  $(P_1^\circ, \Delta_N) \subseteq (\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ})$ .

The following Theorem holds.

**Theorem 3.** *Let  $\Delta_N = (id \otimes d^\circ)\Delta^\circ$ . Then  $(P_1^\circ, \Delta_N) \subseteq (\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ})$ . Therefore, the dual coalgebra  $(\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ})$  of  $\mathcal{L}_1$  is a non-zero Novikov coalgebra. If  $F$  is a field of characteristic zero then the coalgebra  $(\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ})$  does not contain non-zero proper finite-dimensional subcoalgebras.*

Let  $f, g \in \mathcal{L}_1$ . It is easy to see that in the algebra  $\mathcal{L}_1$  we have

$$[f, g] = f \circ g - g \circ f = fd(g) - d(f)g = [f, g]_d.$$

Consequently, the algebra  $(\mathcal{L}_1, [, ])$  is isomorphic to the Witt algebra  $W_1$ .

Define the new comultiplication  $\Delta_{\mathcal{L}_1^\circ}^{(-)} = (1 - \tau)\Delta_{\mathcal{L}_1^\circ}$  on the vector space  $\mathcal{L}_1^\circ$ . Then for every  $f, g \in P_1$  and  $\alpha \in \mathcal{L}_1^\circ$  we have

$$\langle \alpha, [f, g] \rangle = \langle \alpha, f \circ g - f \circ g \rangle = \rho(\Delta_{\mathcal{L}_1^\circ}^{(-)}(\alpha))(f \otimes g).$$

Therefore,  $(\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ}^{(-)})$  is a good subspace of  $(\mathcal{L}_1, [, ])^*$ , and we can assume that  $(\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ}^{(-)})$  is a good subspace of  $W_1^*$ . Consequently,  $(\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ}^{(-)}) \subseteq (W_1^\circ, \Delta_{W_1^\circ})$ .

Theorem 3 implies  $(P_1^\circ, \Delta_{d^\circ}) \subseteq (\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ}^{(-)})$ .

The following statement holds.

**Theorem 4.** *Let  $F$  be a field of characteristic  $\neq 2$ . Then  $(P_1^\circ, \Delta_N) = (\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ})$  and  $(W_1^\circ, \Delta_{W_1^\circ}) = (\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ}^{(-)})$ , where  $\Delta_{\mathcal{L}_1^\circ}^{(-)} = (1 - \tau)\Delta_{\mathcal{L}_1^\circ}$ .*

*Proof.* By Theorem 3, we have  $P_1^\circ \subseteq \mathcal{L}_1^\circ$ . It was shown that  $\mathcal{L}_1^\circ \subseteq W_1^\circ$ . Let  $F$  be a field of characteristic  $\neq 2$ . Then, by Theorem 2,  $P_1^\circ = W_1^\circ$ . Consequently,  $P_1^\circ = \mathcal{L}_1^\circ = W_1^\circ$ . Therefore,  $(W_1^\circ, \Delta_{W_1^\circ}) = (\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ}^{(-)})$ .

Let us prove that  $\Delta_{\mathcal{L}_1^\circ} = \Delta_N$ . Suppose  $f, g$  are arbitrary polynomials in  $P_1$ , and  $\alpha \in \mathcal{L}_1^\circ$ . Then

$$\langle \alpha, fd(g) \rangle = \rho(\Delta^\circ(\alpha))(f \otimes d(g)) = \rho((id \otimes d^\circ)\Delta^\circ(\alpha))(f \otimes g) = \rho(\Delta_N(\alpha))(f \otimes g).$$

On the other hand,

$$\langle \alpha, fd(g) \rangle = \langle \alpha, f \circ g \rangle = \rho(\Delta_{\mathcal{L}_1^\circ}(\alpha))(f \otimes g).$$

Hence,  $\Delta_N = \Delta_{\mathcal{L}_1^\circ}$ , so  $(\mathcal{L}_1^\circ, \Delta_{\mathcal{L}_1^\circ}) = (P_1^\circ, \Delta_N)$ . □

### 3. DUAL COALGEBRA OF THE KANTOR CONSTRUCTION OF POLYNOMIAL ALGEBRA IN ONE VARIABLE

Let  $G$  be the Grassmann algebra with identity. The algebra  $G = G_0 + G_1$  is a  $\mathbb{Z}_2$ -graded algebra. Let  $J = J_0 + J_1$  be a  $\mathbb{Z}_2$ -graded algebra. Then  $G(J) = J_0 \otimes G_0 + J_1 \otimes G_1$  is a subalgebra of  $J \otimes G$  and it is called the *Grassmann envelope* of  $J$ .

An algebra  $J$  is called a *Jordan superalgebra* if its Grassmann envelope is a Jordan algebra, i. e.,  $G(J)$  satisfies the identities

$$xy = yx, \quad (x^2y)x = x^2(xy).$$

Recall the Kantor construction [25]. Let  $A$  be an associative commutative algebra over a field  $F$  with a derivation  $D$ . Denote by  $\bar{A}$  an isomorphic copy of the vector space  $A$  with an isomorphism  $a \mapsto \bar{a}$ . On the direct sum of the vector spaces  $J(A, D) = A + \bar{A}$  define a product  $(\cdot)$  by

$$a \cdot b = ab, \quad a \cdot \bar{b} = \bar{ab}, \quad \bar{a} \cdot b = \bar{ab}, \quad \bar{a} \cdot \bar{b} = aD(b) - D(a)b,$$

where  $a, b \in A$  and  $ab$  is the product of elements in  $A$ . Then  $J(A, D)$  is an Jordan superalgebra. The superalgebra  $J(A, D)$  is said to be a superalgebra of vector type.

Let us give the dual analogue of the Kantor construction for coalgebras. Suppose  $(C, \Delta, d)$  is an associative commutative differential coalgebra with a coderivation  $d$ . Let  $\overline{C}$  be an isomorphic copy of the vector space  $C$  with an isomorphism  $c \mapsto \overline{c}$ . Define the comultiplication  $\Delta_J$  on the direct sum of vector spaces  $J^c(C, \Delta_J, d) = C + \overline{C}$  by

$$\begin{aligned}\Delta_J(c) &= \sum_{(c)} c_{(1)} \otimes c_{(2)} + \overline{c_{(1)}} \otimes \overline{d(c_{(2)})} - \overline{d(c_{(1)})} \otimes \overline{c_{(2)}}, \\ \Delta_J(\overline{c}) &= \sum_{(c)} \overline{c_{(1)}} \otimes c_{(2)} + c_{(1)} \otimes \overline{c_{(2)}},\end{aligned}$$

where  $c \in C$  and  $\Delta(c) = \sum_{(c)} c_{(1)} \otimes c_{(2)}$ . It is shown in [15] that the dual algebra of the coalgebra  $J^c(C, \Delta_J, d)$  is a Jordan superalgebra of vector type  $J(C^*, d^*)$ , where  $d^*$  is the transpose map to  $d$ . Therefore,  $J^c(C, \Delta_J, d)$  is a Jordan supercoalgebra.

**Lemma 5.** *Let  $(P_1^\circ, \Delta^\circ, d^\circ)$  be the dual coalgebra of  $(P_1, d)$ . Then  $J^c(P_1^\circ, \Delta_J^\circ, d^\circ) = J(P_1, d)^\circ$ .*

*Proof.* Set  $J = J(P_1, d)$ , and let  $(J^\circ, \Delta_{J^\circ})$  be the dual supercoalgebra of  $J$ . It was noted above that  $J^c = J^c(P_1^\circ, \Delta_J^\circ, d^\circ)$  is a Jordan supercoalgebra. It is clear that  $J^c$  is a good subspace of  $J^*$ . Therefore,  $J^c \subseteq J^\circ$ .

Let us put  $\langle \alpha, a \rangle = \alpha(a)$  for  $\alpha \in J^*$ ,  $a \in J$ , as above. Since  $J^c = P_1^\circ + \overline{P_1^\circ}$ , where  $\overline{P_1^\circ}$  is an isomorphic copy of the vector space  $P_1^\circ$  with the isomorphism  $a \mapsto \overline{a}$ , we can assume  $\langle \alpha, a \rangle = \langle \overline{\alpha}, \overline{a} \rangle$  for  $\alpha \in P_1^\circ$  and  $a \in P_1$ .

The algebra  $J$  acts on  $J^*$  as on a bimodule by the rule

$$\langle \alpha \cdot a, b \rangle = \langle \alpha, ab \rangle, \quad \langle a \cdot \alpha, b \rangle = \langle \alpha, ba \rangle,$$

where  $\alpha \in J^*$ ,  $a, b \in J$ .

Since  $J = P_1 \oplus \overline{P_1}$ , then we can assume that  $J^* = P_1^* \oplus \overline{P_1^*}$  (e.g.,  $P_1^* \cong \overline{P_1^*} \subset J^*$ ). Moreover,  $P_1^*$  is a  $P_1$ -subbimodule of  $J^*$ . Indeed, let  $\alpha \in P_1^*$  and  $f \in P_1$ . Then

$$\langle \alpha \cdot f, \overline{P_1} \rangle = \langle \alpha, f \overline{P_1} \rangle \subseteq \langle \alpha, \overline{P_1} \rangle = 0.$$

Consequently,  $\alpha \cdot f \in P_1^*$ . Similarly,  $f \cdot \alpha \in P_1^*$ . Hence,  $P_1^*$  is a  $P_1$ -subbimodule of  $J^*$ . In the same way, one may show that  $\overline{P_1^*} \cong P_1^\perp \subset J^*$  is a  $P_1$ -subbimodule of  $J^*$ .

Denote by  $V$  and  $W$  the projections of  $J^\circ$  on the spaces  $P_1^*$  and  $\overline{P_1^*}$ , respectively. Since  $J^\circ$  is a  $P_1$ -subbimodule of  $J^*$  then  $V$  and  $W$  are also  $P_1$ -subbimodules of  $J^*$ .

Let us show that  $V$  is a good subspace of  $P_1^*$ . For  $\alpha \in V$  there exist  $\gamma \in J^\circ$  and  $\beta \in W$  such that  $\gamma = \alpha + \beta$ . Let  $\Delta_{J^\circ}(\gamma) = \sum_{\gamma} \gamma_{(1)} \otimes \gamma_{(2)}$ . Then  $\gamma \cdot P_1 \subseteq \sum_{\gamma} \langle \gamma_{(1)}, P_1 \rangle \gamma_{(2)}$ . Consequently, the space  $\gamma \cdot P_1$  is finite-dimensional. The space  $\alpha \cdot P_1$  is the projection of  $\gamma \cdot P_1$  to  $V$ , therefore,  $\alpha \cdot P_1$  is finite-dimensional. Similarly the space  $P_1 \cdot \alpha$  is finite-dimensional. By Corollary 2.5 from [5] we get that  $V$  a good subspace of  $P_1^*$ .

Consequently,  $V \subseteq P_1^\circ$ . Since  $P_1^\circ \subseteq J^c \subseteq J^\circ$  then  $P_1^\circ \subseteq V$ . Therefore,  $P_1^\circ = V$ . Then  $V \subseteq J^\circ$  and  $W \subseteq J^\circ$ . Hence, the coalgebra  $J^\circ$  is a  $\mathbb{Z}_2$ -graded space, where  $V = (J^\circ)_0$ ,  $W = (J^\circ)_1$ . Moreover,  $\overline{V} = \overline{P_1^\circ} \subseteq W$ .

Assume  $w \in W$ . Then  $\Delta_{J^\circ}(w) = \sum_i v_i \otimes w_i + w_i \otimes v_i$ , where  $v_i \in V$ ,  $w_i \in W$ , and for all  $a \in P_1$  we have

$$\langle w, \bar{a} \rangle = \langle w, \bar{1} \cdot a \rangle = \left\langle \sum_i \langle w_i, \bar{1} \rangle v_i, a \right\rangle = \left\langle \sum_i \langle w_i, \bar{1} \rangle \bar{v}_i, \bar{a} \right\rangle.$$

Since  $W \subseteq \overline{P_1}^*$ , we have  $w = \sum_i \langle w_i, \bar{1} \rangle \bar{v}_i$ . Consequently,  $W = \overline{V}$ , and thus  $J^c(P_1^\circ, \Delta_J^\circ, d^\circ) = J^\circ$ .  $\square$

It is known (see [21]) that a Jordan superalgebra is simple if and only if it is simple as an algebra. Also, recall that the superalgebra  $J(A, D)$  is simple if and only if the algebra  $A$  is a differentially simple algebra [26].

Therefore, the following Theorem is true.

**Theorem 5.** *Let  $P_1 = F[x]$  be the polynomial algebra in the variable  $x$ , and  $d = \frac{d}{dx}$  is the derivation with respect to the variable  $x$ . Consider the Jordan superalgebra  $J(P_1, d)$  of vector type. Then  $J(P_1, d)^\circ = J^c(P_1^\circ, \Delta_J^\circ, d^\circ)$ . Moreover, if  $F$  is a field of characteristic 0 then  $J(P_1, d)^\circ$  does not contain non-zero finite-dimensional subcoalgebras.*

#### REFERENCES

- [1] M. Sweedler, *Hopf algebras*, W.A. Benjamin, Inc., New York, 1969.
- [2] V.G. Drinfeld, *Quantum groups*, In Proc. Int. Cong. Math. (Berkely, 1986), page 798-820. Amer. Math. Soc., Providence, RI, 1987.
- [3] V.G. Drinfeld, *Hamiltonian structures on Lie groups, Lie bialgebras and the geometric meaning of classical Yang-Baxter equations*, Dokl. Akad. Nauk SSSR **268** :2 (1983), 285–287.
- [4] W. Michaelis, *Lie coalgebras*, Adv. Math., **38** (1980), 1-54.
- [5] J. Anquella, T. Cortes, F. Montaner, *Nonassociative coalgebras*, Comm. in Algebra, **22**:12 (1994), 4693-4716.
- [6] V.N. Zhelyabin, *Structurable coalgebras*, Algebra i Logika **35**:5 (1996), 529–542; translation in Algebra and Logic **35**:5 (1996), 296–304.
- [7] V.N. Zhelyabin, *Embedding of Jordan copairs into Lie coalgebras*, Comm. in Algebra, **35**:2 (2007), 561-576.
- [8] G. Santos Filho, L. Murakami, I. Shestakov, *Locally finite coalgebras and the locally nilpotent radical I*, Linear Algebra and its Applications, **621** (2021), 235-253.
- [9] A. M. Slinko, *Local finiteness of coalgebraic Lie coalgebras*, Comm. Algebra. **23**:5 (1995), 1165-1170.
- [10] V.N. Zhelyabin, *The Kantor-Koecher-Tits construction for Jordan coalgebras*, Algebra i Logika **35**:2 (1996), 173–189; translation in Algebra and Logic **35**:2 (1996), 96–104.
- [11] V.N. Zhelyabin, *Jordan (super)coalgebras and Lie (super)coalgebras*, Sib. Mat. Zh. **44**:1 (2003), 87–111; translation in Siberian Math. J. **44**:1 (2003), 73–92.
- [12] M.E. Goncharov, V.N. Zhelyabin, *Mikheev's construction for Mal'tsev coalgebras*, Algebra Logika **51**:5 (2012), 668–671; translation in Algebra Logic **51**:5 (2012), 445–447.
- [13] M.E. Goncharov, V.N. Zhelyabin, *Embedding of Mal'tsev coalgebras into Lie coalgebras with triality*, Algebra Logika **52**:1 (2013), 34–56; translation in Algebra Logic **52**:1 (2013), 24–40.
- [14] D.Kh. Kozybaev, *Right-alternative and right-symmetric coalgebras*, South-Kazakhstan Science and Education. **19**:12 (2000), 155–163.
- [15] D. Kozybaev, U. Umirbaev, V. Zhelyabin, *Some examples of nonassociative coalgebras and supercoalgebras*, Linear Algebra and its Applications **643** (2022), 235-257.
- [16] G. Santos Filho, L. Murakami, and I. Shestakov, *Locally finite coalgebras and the locally nilpotent radical II*, Comm. in Algebra, **49**:12 (2021), 5472–5482.
- [17] W. Michaelis, *An example of a non-zero Lie coalgebra  $M$  for which  $\text{Loc}(M) = 0$* , J. Pure Appl. Algebra. **68** (1990), 341-348.
- [18] W.D. Nichols, *The structure of the dual Lie coalgebra of the Witt algebra*, J. Pure Appl. Algebra. **68** (1990), 359-364.
- [19] W.D. Nichols, *On Lie and associative duals*, J. Pure Appl. Algebra. **87** (1993), 313-320.

- [20] G. Griffing, *The dual coalgebra of certain infinite-dimensional Lie algebras*, *Comm. Algebra* **30**:12 (2002), 5715–5724.
- [21] V.N. Zhelyabin, *Dual coalgebras of Jordan bialgebras and superalgebras*, *Sib. Mat. Zh.* **46**:6 (2005), 1302–1315; translation in *Siberian Math. J.* **46**:6 (2005), 1050–1061.
- [22] I. M. Gel'fand, I. Ya. Dorfman, *Hamiltonian operators and algebraic structures related to them*, *Funktsional. Anal. i Prilozhen* **13**:4 (1979), 3–30.
- [23] I.M. Balinskii, S.P. Novikov, *Poisson brackets of hydrodynamic type, Frobenius algebras and Lie algebras*, *Dokl. Akad. Nauk SSSR* **283**:5 (1985), 1036–1039.
- [24] D. Kozybaev, U. Umirbaev, *Identities of the left-symmetric Witt algebras*, *International Journal of algebra and computation*, **26**:2 (2016), 435–450.
- [25] I.L. Kantor, *Jordan and Lie superalgebras determined by a Poisson algebra*, *Algebra and Analysis*, Tomsk, 1989; translation *Amer. Math. Soc. Transl. Ser. 2*, *Amer. Math. Soc.*, Providence, RI, **151** (1992), 55–80.
- [26] D. King, K. McCrimmon, *The Kantor doubling process revisited*, *Comm. in Algebra*, **23**:1 (1995), 357–372.

VICTOR N. ZHELYABIN  
SOBOLEV INSTITUTE OF MATHEMATICS,  
PR. KOPTYUGA, 4,  
630090, NOVOSIBIRSK, RUSSIA  
*E-mail address*: vicnic@math.nsc.ru

PAVEL S. KOLESNIKOV  
SOBOLEV INSTITUTE OF MATHEMATICS,  
PR. KOPTYUGA, 4,  
630090, NOVOSIBIRSK, RUSSIA  
*E-mail address*: pavelisk77@gmail.com