

## Different estimation methods and validation for the extension exponential distribution

Nouadri R. and Seddik-Ameur N.

Laboratory of probability and statistics LaPS

Badji Mokhtar University Annaba-Algeria

E-mail : rochdi.nouadri@univ-annaba.org

naciraseddik@yahoo.fr

**Abstract:** The extension of the exponential distribution of Nadarajah and Haghghi is particularly interesting because It can describe data with mode set at zero while the hazard rate can be increasing, decreasing and constant. Moreover the hazard function can increase when the probability density function decreases monotonically which is not possible with other models. In this work, different estimation methods such as the maximum product of spacing (*MPS*), the Cramer-von-Mises (*CM*), the Anderson-Darling (*AD*), the Right-tail, the Left-tail (*RAD, LAD*) Anderson-Darling and the Kolmogorov-Smirnov (*KS*) methods are proposed as alternatives to maximum likelihood estimation. Using a new technique based on initial data, a criteria goodness-of-fit test that can validate effectively this model is provided. Thousands of samples are generated to compare between the method performances and the practicality of the proposed criteria test. Real data applications are used for illustration purpose.

**Keywords:** Anderson-Darling method, Cramer-von-Mises method, maximum likelihood estimation, maximum product of spacing method, Right-tail Anderson-Darling method, goodness-of-fit test.

**AMS classification:** 62G10-62N05-62F12

### 1. INTRODUCTION

The recent statistical literature mainly focuses on the introduction of new models capable of describing observations when classical models cannot do so. There are different forms of generalizations either by adding shape parameters, by exponentiation the cumulative distribution or based on generator distributions, one can see Cordeiro et al., (2014), Alzaatreh et al., (2016), Bourguignon et al., (2014), Alzaghal et al., (2013) and others. Nevertheless, Nadarajah and haghghi (2011) proposed a simple and very interesting model. With two parameters, this one offers a good alternative to gamma, Weibull and the exponentiated exponential distributions for which the hazard functions are only constant or decreasing when the probability density function (pdf) is monotonically decreasing while the rate failure of the new distribution can be increasing. Also, this distribution is able to describe data with mode fixed at zero when Its hazard rates can be increasing, decreasing and constant which is not realized for other models. Making a simple change of variable, the new model is reduced to a truncated Weibull distribution. The proposed distribution has more applications than classical models like the fit of the daily rainfall data (for example) which have a monotonically decreasing probability density (pdf) function when the empirical hazard rate failure (hrf) is increasing. The authors studied mathematical properties such the moments, the Bonferroni and Lorenz curves, the entropy and the order statistics. For estimating the unknown parameters, they proposed the maximum likelihood (*MLE*) and the moment estimation (*MME*)

methods which cannot give the explicit forms of the estimators, so iterative methods are required and as It well known the results are performed for big sample sizes. It should be noted also, that the validation of this model is not investigate yet till now which motivate us to propose a new technique for testing the fit of this distribution.

In this work, we propose, firstly, different estimation methods which can be used for small and big sample sizes such as the maximum product of spacing (*MPS*), the Cramer-von-Mises (*CM*) method, Anderson-Darling (*AD*) method, Right-tail and Left-tail Anderson-Darling (*RAD, LAD*) methods and the method of Kolmogorov-Smirnov (*KS*) to estimate the unknown parameters as alternatives to maximum likelihood estimation method. On the other hand, to have reliable results in any statistical analysis, It must be necessary to verify if the observed data are really described by the model chosen. Using an extension of the chi-square statistic introduced by Nikulin (1973) and Rao and Robson (1974) for continuous models, we provide a criteria goodness-of-fit test which takes into account the unknown parameters and the initial data for the extension exponential distribution. This new technique recovers all the information given by the observations and can validate effectively the model without comparison with its competitors. Recently, this method was performed for some new distributions, see Seddik-Ameur and Chouia (2014), Treidi and Seddik-Ameur (2016), Tilbi and Seddik-Ameur (2017). An importante simulation study is given to calculate the different estimators, their biases, their mean square errors and their estimated average widths. A comparison study between the method performances is also provided. Beside classical model selection criteria such as AIC, BIC, LL and Kolmogorov-Smirnov statistic, we provided all the elements of the new statistic test which are used to distinguish between this distribution and its competitors. Real data applications are used for illustration purpose.

## 2. THE EXTENSION EXPONENTIAL DISTRIBUTION

Depending on two parameters  $\alpha$  and  $\lambda$  (shape and scale), the extension of the exponential distribution introduced by Nadarajah and Haghighi (2011) is characterized by the cumulative probability function

$$F(t) = 1 - \exp(1 - (1 + \lambda t)^\alpha)$$

The probability density function (pdf) is given by

$$f(t) = \alpha\lambda(1 + \lambda t)^{\alpha-1} \exp(1 - (1 + \lambda t)^\alpha) \quad \alpha > 0 \text{ and } \lambda > 0.$$

The hazard and the cumulative hazard functions are

$$h(t) = \alpha\lambda(1 + \lambda t)^{\alpha-1}$$

$$H(t) = -\log(S(t)) = (1 + \lambda t)^\alpha - 1$$

Different shapes of the pdf and the hazard rate functions are represented in Figure 1 and Figure 2.




FIGURE 1. probability density function of the extension exponential distribution




FIGURE 2. hazard function of the extension exponential distribution

The interest of this pdf is to have the mode zero and different hazard shapes (increasing, decreasing and constant), furthermore the extension exponential distribution pdf takes only decreasing shapes whereas the hazard function can be increasing (see Figure 3).




FIGURE 3. density and hazard function of the extension exponential distribution with  $\alpha > 1$

The authors studied mathematical properties such as the moments, the Bonferoni and Lorenz curves which have applications in several fields, the entropy and the order statistics. For estimating the unknown parameters, they proposed the maximum likelihood and the moment estimation methods.

### 3. MAXIMUM LIKELIHOOD ESTIMATION

In this section we resume the maximum likelihood estimators of the unknown parameters in order to compare with those of the proposed methods in this work.

We consider a sample of  $n$  independent identically distributed variables  $t_1, t_2, \dots, t_n$  from the extension exponential distribution with the parameter vector  $\theta = (\alpha, \lambda)$ , the likelihood function is given by

$$L(t_1, t_2, \dots, t_n; \theta) = \prod_{i=1}^n f(t_i; \theta) = \prod_{i=1}^n \alpha \lambda (1 + \lambda t_i)^{\alpha-1} \exp(1 - (1 + \lambda t_i)^\alpha)$$

The log-likelihood function is:

$$l = \ln(L(t_1, t_2, \dots, t_n; \theta)) = \sum_{i=1}^n [\ln(\alpha) + \ln(\lambda) + (\alpha - 1) \ln(1 + \lambda t_i) + (1 - (1 + \lambda t_i)^\alpha)]$$

$$l = n + n \ln(\alpha) + n \ln(\lambda) + (\alpha - 1) \sum_{i=1}^n \ln(1 + \lambda t_i) - \sum_{i=1}^n (1 + \lambda t_i)^\alpha$$

The maximum likelihood estimators  $MLE$  are obtained by equating to zero the following first derivatives of the log-likelihood function with respect to the parameters:

$$\begin{aligned}\frac{\partial l}{\partial \alpha} &= \frac{n}{\alpha} + \sum_{i=1}^n \ln(1 + \lambda t_i) - \sum_{i=1}^n (1 + \lambda t_i)^\alpha \ln(1 + \lambda t_i) = 0 \\ \frac{\partial l}{\partial \lambda} &= \frac{n}{\lambda} + (\alpha - 1) \sum_{i=1}^n t_i (1 + \lambda t_i)^{-1} - \alpha \sum_{i=1}^n t_i (1 + \lambda t_i)^{\alpha-1} = 0\end{aligned}$$

As the explicit forms of the  $MLE$  ( $\hat{\alpha}_{MLE}, \hat{\lambda}_{MLE}$ ) cannot be derived, one can use numerical methods for solving the obtained equations.

#### 4. PRODUCT OF SPACING ESTIMATION METHOD MPS

Introduced by Cheng and Amin (1979, 1983), the product of spacing estimation method ( $MPS$ ) is an alternative to the maximum likelihood estimation method. This method was also proposed by Ramnaby (1984) as an approximation to the Kullback-Leibler measure of information.

We consider  $n$  independent and identically distributed sample drawn from a population with unknown parameters  $\theta$ . The  $MPS$  estimation method is based on the differences between the successive increasing empirical repartition functions  $F(t_i, \theta)$ , and where  $F(t_0) = 0$  and  $F(t_{n+1}) = 1$ .

$$P_i(\theta) = F(t_i, \theta) - F(t_{i-1}, \theta)$$

The  $MPS$  estimators are obtained by maximizing the quantity  $M(\theta)$  :

$$M(\theta) = \frac{1}{n+1} \sum_{i=1}^{n+1} \ln(P_i(\theta))$$

For the extension exponential distribution, this function is given by:

$$M(\theta) = \frac{1}{n+1} \sum_{i=1}^{n+1} \ln((1 - \exp(1 - (1 + \lambda t_i)^\alpha)) - (1 - \exp(1 - (1 + \lambda t_{i-1})^\alpha)))$$

$$M(\theta) = \frac{1}{n+1} \sum_{i=1}^{n+1} \ln(\exp(1 - (1 + \lambda t_{i-1})^\alpha) - \exp(1 - (1 + \lambda t_i)^\alpha))$$

So, the  $MPS$  estimators are the solutions of the following equations

$$\begin{aligned}\frac{\partial M(\theta)}{\partial \alpha} &= \frac{1}{n+1} \sum_{i=1}^{n+1} \frac{D1(t_i, \theta) - D1(t_{i-1}, \theta)}{P_i} = 0 \\ \frac{\partial M(\theta)}{\partial \lambda} &= \frac{1}{n+1} \sum_{i=1}^{n+1} \frac{D2(t_i, \theta) - D2(t_{i-1}, \theta)}{P_i} = 0\end{aligned}$$

where  $D1$  and  $D2$  are the first derivatives of the repartition function with respect to the unknown parameters  $\alpha$  and  $\lambda$ :

$$D1 = \frac{dF(t)}{d\alpha} = (1 + \lambda t)^\alpha \ln(1 + \lambda t) \exp(1 - (1 + \lambda t)^\alpha)$$

$$D2 = \frac{dF(t)}{d\lambda} = \alpha t(1 + \lambda t)^{\alpha-1} \exp(1 - (1 + \lambda t)^\alpha)$$

Cheng and Amin (1983) showed that the obtained *MPS* estimators are as efficient and consistent as the maximum likelihood estimators (*MLE*) under more general conditions.

## 5. CRAMER VON MISES ESTIMATION METHOD

Among minimum type estimation methods proposed in the statistical literature, the Cramer-Von-Mises *CM* method estimation consists in minimizing the distances between theoretical and empirical repartition functions (Cramer-Von-Mises statistic) of the studied model. This method is shown to give the smaller bias estimators than the other minimum distance estimators (see Mac-Donald, 1971).

For the extension of the exponential distribution, this statistic is defined by

$$C(a, \lambda) = \frac{1}{12} + \sum_{i=1}^n \left( (1 - \exp(1 - (1 + \lambda t_i)^a)) - \frac{2i-1}{2n} \right)^2$$

To calculate the *CM* estimators ( $\widehat{a}_{CV}, \widehat{\lambda}_{CV}$ ) of the unknown parameters, we cancel the first partial derivatives of this statistic  $C(a, \lambda)$  with respect to the unknown parameters of the distribution, which amounts to solving the following equations:

$$\begin{aligned} \sum_{i=1}^n \left( (1 - \exp(1 - (1 + \lambda t_i)^a)) - \frac{2i-1}{2n} \right) D1 &= 0 \\ \sum_{i=1}^n \left( (1 - \exp(1 - (1 + \lambda t_i)^a)) - \frac{2i-1}{2n} \right) D2 &= 0 \end{aligned}$$

where  $D1$  and  $D2$  are as given above.

## 6. ANDERSON-DARLING ESTIMATION METHOD

Anderson-Darling method is also a minimum type estimation method, so the estimators ( $\widehat{a}_{AD}, \widehat{\lambda}_{AD}$ ) are obtained by minimizing the well-known Anderson-Darling statistic  $A^2$  suggested by Anderson and Darling (1952) with respect to the unknown parameters. Boos (1981) also discussed the properties of the *AD* estimators and presented its computational form as

$$A^2(a, \lambda) = -n - \frac{1}{n} \sum_{i=1}^n (2i-1) [\ln(F(t_i)) - \ln(1 - F(t_{i^*}))]$$

$$A^2(a, \lambda) = -n - \frac{1}{n} \sum_{i=1}^n (2i-1) [\ln(1 - \exp(1 - (1 + \lambda t_i)^a)) - \ln(1 - (1 - \exp(1 - (1 + \lambda t_{i^*})^a)))]$$

where  $i^* = n - i + 1$ .

For our model, the estimators ( $\widehat{a}_{AD}, \widehat{\lambda}_{AD}$ ) are obtained by the resolution of the following equations:

$$\sum_{i=1}^n (2i-1) \left[ \frac{D1(t_i)}{(1 - \exp(1 - (1 + \lambda t_i)^a))} - \frac{D1(t_{i^*})}{(1 - (1 - \exp(1 - (1 + \lambda t_{i^*})^a)))} \right] = 0$$

$$\sum_{i=1}^n (2i-1) \left[ \frac{D2(t_i)}{(1 - \exp(1 - (1 + \lambda t_i)^a))} - \frac{D2(t_{i^*})}{(1 - (1 - \exp(1 - (1 + \lambda t_{i^*})^a)))} \right] = 0$$

Later, Luceno (2006) discussed modifications of the standard  $AD$  statistics. The most used statistic is the Right-tail  $AD$  statistics (see Dey et al., (2014), Louzada et al., (2016) and Ramos and Louzada (2016)). The Right tail Anderson-Darling estimators ( $\hat{a}_{RAD}, \hat{\lambda}_{RAD}$ ) are obtained by minimizing the following expression:

$$RAD = \frac{n}{2} - 2 \sum_{i=1}^n \ln(F(t_i)) - \frac{1}{n} \sum_{i=1}^n (2i-1)(1 - (F(t_{i^*})))$$

$$RAD = \frac{n}{2} - 2 \sum_{i=1}^n \ln((1 - \exp(1 - (1 + \lambda t_i)^a))) - \frac{1}{n} \sum_{i=1}^n (2i-1)(1 - (1 - \exp(1 - (1 + \lambda t_{i^*})^a)))$$

which implies the resolution of the following equation system:

$$-2 \sum_{i=1}^n \frac{D1(t_i)}{(1 - \exp(1 - (1 + \lambda t_i)^a))} + \frac{1}{n} \sum_{i=1}^n (2i-1) \frac{D1(t_{i^*})}{(1 - ((1 - \exp(1 - (1 + \lambda t_{i^*})^a)))} = 0$$

$$-2 \sum_{i=1}^n \frac{D2(t_i)}{(1 - \exp(1 - (1 + \lambda t_i)^a))} + \frac{1}{n} \sum_{i=1}^n (2i-1) \frac{D2(t_{i^*})}{(1 - (1 - \exp(1 - (1 + \lambda t_{i^*})^a)))} = 0$$

The Left tail Anderson-Darling estimators ( $\hat{a}_{LAD}, \hat{\lambda}_{LAD}$ ) are obtained by minimizing the following expression

$$LAD = \frac{-3n}{2} + 2 \sum_{i=1}^n \ln(F(t_i)) - \frac{1}{n} \sum_{i=1}^n (2i-1)(1 - (F(t_{i^*})))$$

$$LAD = \frac{-3n}{2} + 2 \sum_{i=1}^n \ln((1 - \exp(1 - (1 + \lambda t_i)^a))) - \frac{1}{n} \sum_{i=1}^n (2i-1)(1 - (1 - \exp(1 - (1 + \lambda t_{i^*})^a)))$$

## 7. KOLMOGOROV-SMIRNOV ESTIMATION METHOD

Let us consider an independent and identically distributed sample drawn from a population with unknown vector parameters  $\theta$ . Weber et al. (2006) considered an estimation technique so-called as the minimum Kolmogorov-Smirnov estimation ( $KS$ ), whose purpose is to minimize the value of the famous Kolmogorov-Smirnov statistic which represents the biggest value of the difference between the theoretical and the corresponding empirical repartition functions of the proposed model:

$$KS = d(F_n, F) = \max_{x \in \mathfrak{R}} |F_n(t) - F(t)|$$

If  $F(t)$  is the extension exponential distribution, we have

$$KS = d(F_n, F) = \max_{t \in \mathfrak{R}} | F_n(t) - (1 - \exp(1 - (1 + \lambda t)^a)) |$$

where  $n$  is the sample size,  $F(t)$  is the hypothesized distribution (cdf), and  $F_n(t)$  is the empirical cdf, a step-function that increases by  $1/n$  at each data value. The  $KS$  statistic has been widely used to fit data to continuous populations. The interest of this statistic includes the straightforward computation of the test statistic and its distribution-free characteristic.

The computational formulas for computing  $KS$  are:

$$D^+ = \max_{i=1,2,\dots,n} \left( \frac{i}{n} - (1 - \exp(1 - (1 + \lambda t_{(i)})^a)) \right)$$

$$D^- = \max_{i=1,2,\dots,n} \left( (1 - \exp(1 - (1 + \lambda t_{(i)})^a)) - \frac{i-1}{n} \right)$$

and

$$KS = \max \{ D^+, D^- \}$$

where  $t_{(1)}, t_{(2)}, \dots, t_{(n)}$  are the order statistics. The parameter values of that minimize the  $KS$  statistic are noted  $(\hat{\alpha}_{KS}, \hat{\lambda}_{KS})$ .

## 8. MODEL VALIDATION

The validation of the model chosen to describe collected data for the analysis is very important. The most common used goodness-of-fit tests are those based on empirical distribution functions (EDF) such as the Kolmogorov-Smirnov test and the famous chi-square test based on Pearson statistic  $X^2$ . Nevertheless the theory showed that the chi-square goodness-of-fit test cannot be applied when the distribution is not specified and the distribution of the statistic test depends on the method used for the estimation of the unknown parameters. So different modifications were proposed in the statistical literature. In this work, we use the technique introduced by Nikulin (1973) and Rao and Robson (1974) to construct a criteria test  $Y^2$  to validate the extension exponential distribution.

Let us consider the null hypothesis that a sample  $T_1, T_2, \dots, T_n$  follows the extension exponential distribution  $P(T_i < t) = F(t, \theta)$  with the unknown parameter vector  $\theta = (\alpha, \lambda)^T$ .

The statistic  $Y^2$  is defined as the sum of the Pearson statistic  $X^2$  and a quadratic form  $Q$  which depends on the hypothesized model. Based on maximum likelihood estimators on initial data, this statistic follows a chi-square distribution with  $r - 1$  degrees of freedom where  $r$  represents the number of grouped classes.

$$Y^2(\hat{\theta}) = X_n^2(\hat{\theta}) + Q_n(\hat{\theta})$$

The computational formula of  $Q(\hat{\theta})$  is given as (see Van der Vaart, 1998):

$$Q_n(\hat{\theta}) = \frac{1}{n} L^T(\hat{\theta}_n) \left( I(\hat{\theta}_n) - J(\hat{\theta}_n) \right)^{-1} L(\hat{\theta}_n)$$

where  $I(\hat{\theta}_n)$  and  $J(\hat{\theta}_n)$  are the estimated information matrices respectively for initial data and grouped data into the  $r$  classes  $I_j$  chosen:

$$I_j = [a_j, a_{j+1}[; \quad \text{for } j = 1, \dots, r-1$$

As the grouped intervals must be equiprobable, so the limits  $a_j$  are obtained as

$$a_j = F^{-1}(j/r)$$

$$a_j = F^{-1}(j/r) = \frac{(1 - (\log(1 - \frac{j}{r}))^{1/\alpha} - 1}{\lambda}$$

If  $v_1, v_2, \dots, v_r$  are the theoretical frequencies of grouped data in these intervals

$$v_j = \text{card}(i, T_i \in I_j, i = 1, \dots, n)$$

So

$$X_n^2(\hat{\theta}) = X_n^T(\hat{\theta})X_n(\hat{\theta})$$

with

$$X_n(\hat{\theta}) = \left( \frac{v_1 - np_1(\theta)}{\sqrt{np_1(\theta)}}, \dots, \frac{v_r - np_r(\theta)}{\sqrt{np_r(\theta)}} \right)$$

and

$$p_j = F(a_j) - F(a_{j-1}) = (1 - \exp(1 - (1 + \lambda a_j)^\alpha)) - (1 - \exp(1 - (1 + \lambda a_{j-1})^\alpha)) \quad \text{with } j = 1, \dots, r$$

The element of the quadratic form  $Q$  for the extension exponential distribution are obtained as follow:

$$L = \left( L_1 = \sum_{j=1}^r \frac{v_j}{p_j(\theta)} \frac{dp_j(\theta)}{d\alpha}, \quad L_2 = \sum_{j=1}^r \frac{v_j}{p_j(\theta)} \frac{dp_j(\theta)}{d\lambda} \right)$$

$$J(\theta) = B^T(\theta)B(\theta) \quad \text{with } B(\theta) = \left( \frac{1}{\sqrt{p_j(\theta)}} \frac{dp_j(\theta)}{d\alpha}, \frac{1}{\sqrt{p_j(\theta)}} \frac{dp_j(\theta)}{d\lambda} \right)$$

The components of Fisher information matrix are

$$I_{11} = \left( \frac{-1}{n} \right) \sum_{i=1}^n - \left( \frac{1}{\alpha^2} + (1 + \lambda t_i)^\alpha (\ln(1 + \lambda t_i))^2 \right)$$

$$I_{22} = \left( \frac{-1}{n} \right) \sum_{i=1}^n - \left( \frac{1}{\lambda^2} + \frac{(\alpha - 1)t_i^2}{(1 + \lambda t_i)^2} + (1 + \lambda t_i)^{\alpha-2} (\alpha - 1) \alpha t_i^2 \right)$$

$$I_{12} = \left( \frac{-1}{n} \right) \sum_{i=1}^n \frac{t_i}{1 + \lambda t_i} + \frac{t(1 + \lambda t_i)^\alpha}{(1 + \lambda t_i)} - \alpha t_i \ln(1 + \lambda t_i) (1 + \lambda t_i)^{\alpha-1}$$

$$I_{21} = I_{12} = \left( \frac{-1}{n} \right) \sum_{i=1}^n \frac{t_i}{1 + \lambda t_i} + \frac{t(1 + \lambda t_i)^\alpha}{(1 + \lambda t_i)} - \alpha t_i \ln(1 + \lambda t_i) (1 + \lambda t_i)^{\alpha-1}$$

Thus all the elements of the statistic test  $Y^2$  for extension exponential distribution are provided. The applicability of this test is shown by an intensive simulation study.

## 9. SIMULATION STUDY

**Estimation:**

To evaluate the performances of the different methods proposed in this work, we generated  $N = 10,000$  samples with different sizes ( $n = 15, 25, 40, 80$  and  $n = 150$ ) from the extension exponential distribution. We compute the estimators of the unknown parameters, their biases, their mean square errors and their estimated average widths by different methods with the following values of the parameters  $\alpha = 0.2, \lambda = 0.1$ . The results are summarized in the following tables (Table1, 2, 3, 4, 5, 6 and 7).

n	$(\hat{\alpha}, \hat{\lambda})$	$MSE(\hat{\theta})$	$bias(\hat{\theta})$	$AW$
15	(0.2252 0.1162)	(0.00368 0.01256)	(0.02527 0.01620)	(0.1188 0.2196)
25	(0.2195 0.1148)	(0.00265 0.01129)	(0.01954 0.01482)	(0.1008 0.2082)
40	(0.2105 0.1105)	(0.00115 0.00540)	(0.01054 0.01055)	(0.0664 0.1440)
80	(0.2054 0.1040)	(0.00049 0.00212)	(0.00542 0.00407)	(0.0433 0.0902)
150	(0.2023 0.1028)	(0.00023 0.00101)	(0.00230 0.00283)	(0.0297 0.0622)

Table 1: *MLE* estimators for parameter values  $\alpha = 0.2, \lambda = 0.1$

n	$(\hat{\alpha}, \hat{\lambda})$	$MSE(\hat{\theta})$	$bias(\hat{\theta})$	$AW$
15	(0.1890 0.2097)	(0.00205 0.05728)	(0.01093 0.10977)	(0.0887 0.4687)
25	(0.1902 0.1785)	(0.00164 0.03929)	(0.00976 0.07854)	(0.0793 0.3885)
40	(0.1918 0.1441)	(0.00089 0.01032)	(0.00811 0.04416)	(0.0584 0.1991)
80	(0.1944 0.1217)	(0.00044 0.00301)	(0.00555 0.02177)	(0.0411 0.1075)
150	(0.1963 0.1116)	(0.00022 0.00121)	(0.00360 0.01165)	(0.0290 0.0681)

Table 2: *MPS* estimators for parameter values  $\alpha = 0.2, \lambda = 0.1$

n	$(\hat{\alpha}, \hat{\lambda})$	$MSE(\hat{\theta})$	$bias(\hat{\theta})$	$AW$
15	(0.2425 0.1425)	(0.02460 0.05036)	(0.04252 0.04250)	(0.3074 0.4398)
25	(0.2193 0.1270)	(0.00393 0.01534)	(0.01931 0.02707)	(0.1228 0.2427)
40	(0.2128 0.1142)	(0.00225 0.00663)	(0.01280 0.01424)	(0.0929 0.1595)
80	(0.2054 0.1074)	(0.00077 0.00270)	(0.00549 0.00747)	(0.0543 0.1018)
150	(0.2032 0.1033)	(0.00038 0.00124)	(0.00322 0.00332)	(0.0382 0.0690)

Table 3: *CvM* estimators for parameter values  $\alpha = 0.2, \lambda = 0.1$

n	$(\hat{\alpha}, \hat{\lambda})$	$MSE(\hat{\theta})$	$bias(\hat{\theta})$	$AW$
15	(0.2179 0.1631)	(0.00545 0.03055)	(0.01796 0.06314)	(0.1446 0.3425)
25	(0.2090 0.1329)	(0.00228 0.01512)	(0.00908 0.03295)	(0.0935 0.2410)
40	(0.2055 0.1184)	(0.00122 0.00643)	(0.00557 0.01844)	(0.0684 0.1571)
80	(0.2021 0.1093)	(0.00054 0.00235)	(0.00215 0.00930)	(0.0455 0.0950)
150	(0.2012 0.1049)	(0.00028 0.00117)	(0.00127 0.00499)	(0.0327 0.0670)

Table 4: *AD* estimators for parameter values  $\alpha = 0.2, \lambda = 0.1$

n	$(\hat{\alpha}, \hat{\lambda})$	$MSE(\hat{\theta})$	$bias(\hat{\theta})$	AW
15	(0.2237 0.1801)	(0.00921 0.07814)	(0.02378 0.08012)	(0.1880 0.5478)
25	(0.2115 0.1270)	(0.00232 0.01405)	(0.01152 0.02700)	(0.0944 0.2323)
40	(0.2070 0.1197)	(0.00133 0.00739)	(0.00708 0.01972)	(0.0714 0.1684)
80	(0.2032 0.1092)	(0.00055 0.00280)	(0.00325 0.00922)	(0.0459 0.1037)
150	(0.2018 0.1040)	(0.00028 0.00123)	(0.00186 0.00406)	(0.0327 0.0687)

Table 5: RAD estimators for parameter values  $\alpha = 0.2, \lambda = 0.1$

n	$(\hat{\alpha}, \hat{\lambda})$	$MSE(\hat{\theta})$	$bias(\hat{\theta})$	AW
15	(0.2227 0.1652)	(0.00519 0.06445)	(0.02277 0.06527)	(0.1412 0.4975)
25	(0.2243 0.1373)	(0.00610 0.02161)	(0.02433 0.03737)	(0.1530 0.2881)
40	(0.2120 0.1213)	(0.00271 0.00849)	(0.01201 0.02133)	(0.1020 0.1805)
80	(0.2057 0.1095)	(0.00097 0.00301)	(0.00576 0.00951)	(0.0610 0.1075)
150	(0.2030 0.1046)	(0.00047 0.00135)	(0.00303 0.00465)	(0.0424 0.0720)

Table 6: LAD estimators for parameter values  $\alpha = 0.2, \lambda = 0.1$

n	$(\hat{\alpha}, \hat{\lambda})$	$MSE(\hat{\theta})$	$bias(\hat{\theta})$	AW
15	(0.2417 0.1501)	(0.01127 0.06134)	(0.04173 0.05015)	(0.2080 0.4854)
25	(0.2229 0.1222)	(0.00441 0.01555)	(0.02292 0.02227)	(0.0411 0.2444)
40	(0.2136 0.1127)	(0.00204 0.00677)	(0.01368 0.01274)	(0.0885 0.1612)
80	(0.2065 0.1066)	(0.00084 0.00282)	(0.00652 0.00665)	(0.0568 0.1040)
150	(0.2036 0.1033)	(0.00040 0.00130)	(0.00366 0.00336)	(0.0392 0.0706)

Table 7: KS estimators for parameter values  $\alpha = 0.2, \lambda = 0.1$

From numerical experiments, It is observed that the biases and the MSE decrease as the sample size increases and this is for all methods. Regarding tables (1,2,3,4,5,6 and 7), we can see that for little sample sizes, the MPS, the AD and the RAD methods provided very good results but the MPS estimators are the best ones compared to all the other methods. For big sample sizes, all methods gave approximately the same results, however the AD and the RAD estimators are better than all of them and even that of the maximum likelihood MLE method.

#### criteria test

To show the practicability of the statistic test provided in this work, we test the null hypothesis  $H_0$  that samples are drawn from the extension exponential distribution. The number of rejection of the null hypothesis  $H_0$  for thousands of samples from different sizes ( $n = 15, 25, 40, 80$  and  $150$ ) are computed and compared with those of their theoretical ones ( $\epsilon = 0, 01; 0, 05$  and  $0, 1$ ). The results are presented in Table 8.

n	$\epsilon = 0.05$	$\epsilon = 0.01$	$\epsilon = 0.1$
15	(0.03879)	(0.01697)	(0.1230)
25	(0.04918)	(0.01513)	(0.1080)
40	(0.05638)	(0.01503)	(0.1068)
80	(0.05204)	(0.01222)	(0.1022)
150	(0.05073)	(0.01124)	(0.1004)

Table 8. critical chi-square and their corresponding empirical values

As expected, the obtained empirical values are very close to their corresponding theoretical ones which implies the feasibility of the proposed goodness-of-fit test to validate the extension exponential distribution.

## 10. APPLICATIONS ON REAL DATA SETS

In this section, we applied the obtained results to two real data sets.

**Example 1** The first one concerns the failure times of the air conditioning system of an air plane studied firstly by Linhart and Zucchini (1986), and Gupta and Kundu (2001) who fitted these data respectively to Weibull and exponentiated exponential distributions. Recently Dinesh et al.(2017) showed that the modified Weibull distribution is more appropriated. The observed values are given below: 23, 261, 87, 7, 120, 14, 62, 47, 225, 71, 246, 21, 42, 20, 5, 12, 120, 11, 3, 14, 71, 11, 14, 11, 16, 90, 1, 16, 52, 95.

Using the estimation methods given above, we compute the values of the unknown parameters considering different distributions capable to describe these data namely the extension exponential, Weibull, exponentiated exponential and modified Weibull distributions. The results are given in table 8.

The cumulative density function of modified Weibull distribution considered here is:

$$F_{MW} = \frac{2F_W}{1 + F_W} = \frac{2(1 - e^{-(\frac{t}{\lambda})^\alpha})}{2 - e^{-(\frac{t}{\lambda})^\alpha}}$$

Distributions	parameters	MLE	CVM	KS	AD	LAD	RAD	MPS
<i>Extension exp</i>	$\alpha$	0.5985	0.4866	0.6947	0.5233	0.5531	0.5081	0.5057
	$\lambda$	4.3385	6.3226	3.8403	5.4853	5.0388	5.8283	5.7787
<i>Weibull</i>	$\alpha$	0.8535	0.7854	0.9000	0.7994	0.8775	0.7560	0.7805
	$\lambda$	54.612	0.5183	0.4496	0.5324	0.4906	0.5265	0.5577
<i>M.Weibull</i>	$\alpha$	0.9869	0.8989	1.0260	0.9215	0.9909	0.8829	0.9023
	$\lambda$	84.097	0.8633	0.7095	0.8664	0.7808	0.8807	0.9050
<i>Exponentiated exp</i>	$\alpha$	0.8090	0.6979	0.8451	0.7082	0.8528	0.6243	0.6990
	$\lambda$	0.0145	1.3880	1.9274	1.3489	1.8123	1.1922	1.2200

Table 8. estimator values of the unknown parameters of competitor distributions of air conditioning data system of air planes

To differentiated between the competitor distributions, and based on the classical model selection criteria, we calculate the *AIC*, *BIC*, *LL*, *KS*, *CvM* and *AD* statistics, the results are given in Table 9. As It can be seen, for the maximum likelihood estimation method, only the values of *AIC*, *BIC* and *LL* of the modified Weibull model are the smallest ones. But for the other estimation methods, all criteria selection models *AIC*, *BIC*, *LL*, *KS*, *CvM* and *AD* of the extension exponential distribution give the smaller values. So, It can be said that the extension

exponential distribution might work better than all the other models, and this fact is proved by data plots of the different pdf distributions (see Figure 4).

	methods	extension exp	Weibull	Exponentiated exp	M.Weibull
AIC	MLE	(30.8527)	(31.5635)	(32.0910)	(30.6859)
	CvM	(31.3652)	(31.9047)	(32.5510)	(31.1571)
	KS	(31.6769)	(33.0019)	(33.9210)	(34.2721)
	LAD	(30.9216)	(31.9584)	(33.0875)	(30.8079)
	RAD	(31.1693)	(32.2839)	(33.2811)	(31.3592)
	AD	(31.0697)	(31.7777)	(32.4168)	(30.9512)
	MPS	(31.2078)	(32.0362)	(32.595)	(31.186)
BIC	MLE	(33.6551)	(34.3659)	(34.8934)	(33.4883)
	CvM	(34.1676)	(34.7071)	(35.3534)	(33.9595)
	KS	(34.4793)	(35.8043)	(36.7234)	(34.2721)
	LAD	(33.7240)	(34.7608)	(35.8899)	(33.6103)
	RAD	(33.9717)	(35.0863)	(36.0835)	(34.1616)
	AD	(33.8721)	(34.5801)	(35.2191)	(33.7536)
	MPS	(34.0101)	(34.8385)	(35.3973)	(33.9883)
LL	MLE	(-13.4263)	(-13.7817)	(-14.0455)	(-13.3429)
	CvM	(-13.6826)	(-13.9523)	(-14.2755)	(-13.5785)
	KS	(-13.8384)	(-14.5009)	(-14.9605)	(-13.7348)
	LAD	(-13.4608)	(-13.9792)	(-14.5437)	(-13.4039)
	RAD	(-13.5846)	(-14.1419)	(-14.6405)	(-13.6796)
	AD	(-13.5348)	(-13.8888)	(-14.2084)	(-13.4756)
	MPS	(-13.6039)	(-14.0181)	(-14.2975)	(-13.5930)
KS	MLE	(0.1318)	(0.1533)	(0.1719)	(0.1427)
	CvM	(0.1206)	(0.1229)	(0.1289)	(0.1202)
	KS	(0.1089)	(0.1120)	(0.1135)	(0.1080)
	LAD	(0.1215)	(0.1312)	(0.1339)	(0.1237)
	RAD	(0.1157)	(0.1303)	(0.1367)	(0.1234)
	AD	(0.1203)	(0.1331)	(0.1411)	(0.1267)
	MPS	(0.1198)	(0.1393)	(0.1595)	(0.5943)

Table 9: AIC, BIC, LL and KS statistic values

For testing the null hypothesis  $H_0$  that these data follow the extension exponential model, we use the criteria test constructed in this work. We grouped data into  $r = 8$  intervals, after calculating the limits of these intervals, we obtain:

$$L = \left( L_1 = -2.5073, L_2 = -0.6273 \right)$$

$$J(\hat{\theta}) = \begin{bmatrix} 2.1928 & 0.1684 \\ 0.1684 & 0.0145 \end{bmatrix}$$

$$I_n(\hat{\theta}) = \begin{bmatrix} 7.5394 & 0.4651 \\ 0.4651 & 0.0330 \end{bmatrix}$$

we deduce the value of  $Y^2 = 9.0753$ . For level of significance  $\epsilon = 0.05$ , the chi-square critical value is  $X_{r-1}^2 = 14.07$ . As  $Y^2 = 9.0753 < X_{r-1}^2 = 14.07$ , the null hypothesis  $H_0$  cannot be rejected, therefore the failure times of the air conditioning system of an air plane is effectively modeled by the extension exponential

distribution. The pdf plots and the empirical distribution function plots given in Figure 4, show that the extension exponential distribution is best suited to model these data.




FIGURE 4. empirical pdf and cdf versus theoretical alternative distributions

From these results, we can say that the extension exponential distribution fit these data better than all its alternatives.

### Example 2

Here we consider a classical flood data of the Floyd River given in Mudholkar and Huston (1996). In the paper of Akisente et al., (2008), the authors calculated the log-likelihood of these data for different distributions and the obtained values are respectively -376.35 for the exponentiated Weibull distribution, -382.13 for the beta-Weibull distribution and -365.45 for the beta-Pareto model. In this subsection, we propose the extension exponential distribution and its competitors to describe these data. We compute the different estimators of the unknown parameters (Table 10) and the criteria selection models AIC, BIC, LL and KS (Table 11).

Distributions	parameters	MLE	CVM	KS	AD	LAD	RAD	MPS
<i>Extension exp</i>	$\alpha$	0.6093	1.2844	0.8757	0.8301	4.0602	0.6793	0.5295
	$\lambda$	0.3358	1.1373	0.2053	0.2355	0.0364	0.3358	0.4625
<i>Weibull</i>	$\alpha$	0.8714	1.1196	1.0977	1.0011	1.2663	0.8886	0.8088
	$\lambda$	6.1888	5.0977	5.6478	5.5515	4.8755	5.4194	6.3671
<i>Gamma</i>	$\alpha$	0.9195	1.2261	1.1642	1.0716	1.4655	0.8615	0.8074
	$n$	0.1358	0.2498	0.2131	0.1970	0.3194	0.1523	0.1109
<i>Exponentiated exp</i>	$\alpha$	0.9688	1.2485	1.1758	1.0996	1.5061	0.8690	0.8357
	$\lambda$	0.1445	0.2337	0.2026	0.1963	0.2783	0.1617	0.1213

Table 10. Different estimators of the unknown parameters of competitor distributions for Floyd river data

	methods	Extension exp	Weibull	Gamma	Exponentiated exp
AIC	MLE	(225.9119)	(229.4550)	(231.0047)	(231.167)
	CvM	(246.0783)	(243.3696)	(238.8346)	(237.5375)
	KS	(229.0334)	(237.9247)	(234.7716)	(234.005)
	AD	(228.4938)	(232.8804)	(233.7901)	(233.7134)
	LAD	(571.5404)	(265.8108)	(247.2532)	(243.5087)
	RAD	(226.4296)	(230.1609)	(232.2683)	(232.7168)
	MPS	(226.4262)	(230.0223)	(231.5818)	(231.7741)
BIC	MLE	(229.2391)	(232.7821)	(234.3318)	(234.4941)
	CvM	(249.4054)	(246.6968)	(242.1618)	(240.8647)
	KS	(232.3605)	(241.2518)	(238.0988)	(237.3321)
	AD	(231.8209)	(236.2075)	(237.1172)	(237.0406)
	LAD	(574.8675)	(269.1379)	(250.5803)	(246.8356)
	RAD	(229.7567)	(233.4880)	(235.5954)	(236.0439)
	MPS	(229.7533)	(233.3495)	(234.9089)	(235.1012)
LL	MLE	(-110.956)	(-112.7275)	(-113.5023)	(-113.5835)
	CvM	(-121.0392)	(-119.6848)	(-117.4173)	(-116.7688)
	KS	(-112.5167)	(-116.9624)	(-115.3858)	(-115.0025)
	AD	(-112.2469)	(-114.4402)	(-114.895)	(-114.8567)
	LAD	(-283.7702)	(-130.9054)	(-121.6266)	(-119.7543)
	RAD	(-111.2148)	(-113.0804)	(-114.1341)	(-114.3584)
	MPS	(-111.2131)	(-113.0112)	(-113.7909)	(-113.887)
KS	MLE	(0.1366)	(0.1270)	(0.1471)	(0.1502)
	CvM	(0.1067)	(0.1015)	(0.1003)	(0.0996)
	KS	(0.1034)	(0.0782)	(0.0768)	(0.0764)
	AD	(0.1172)	(0.1059)	(0.0950)	(0.0916)
	LAD	(0.1360)	(0.1264)	(0.1200)	(0.1179)
	RAD	(0.1409)	(0.1425)	(0.1360)	(0.1332)
	MPS	(0.6684)	(0.6715)	(0.7197)	(0.7314)

Table 11: AIC, BIC, LL and KS statistic values for Floyd river data

Compared to the LL values obtained for the different models cited above, we find the value LL=(-110.956) for the extension exponential distribution which proves that this model is better than all the alternative distributions.

**NRR statistic** On the other hand, we suppose the null hypothesis  $H_0$  that the Floyd river data belong to the extension exponential distribution. If we choose  $r = 8$  grouping classes, the intermediate calculation of the criteria test  $Y^2$  are as follow

$$L = \left( L_1 = 10.2593, L_2 = 5.3251 \right)$$

$$J(\theta)(\hat{\theta}) = \begin{bmatrix} 4.4775 & 4.0355 \\ 4.0355 & 3.9302 \end{bmatrix}$$

$$I_n(\hat{\theta}) = \begin{bmatrix} 6.9144 & 5.1848 \\ 5.1848 & 5.9629 \end{bmatrix}$$

We obtain  $Y^2 = 6,6724$ , so for the significance level  $\epsilon = 0.05$ , the critical value  $X_{r-1}^2 = 14.07$ , therefore we can validate the extension exponential distribution, even the curves of the empirical function distributions and their pdf plots (Figure 5) corroborate the test results.




FIGURE 5. empirical pdf and cdf versus theoretical alternative distributions

### Conclusion:

As the explicit forms of the maximum likelihood ( $MLE$ ) estimators cannot be derived, we have used different estimation methods as alternatives to calculate the unknown parameters of the extension exponential distribution. Based on an important simulation study, we have showed that the maximum product spacing ( $MPS$ ) method gives the best results for little and moderate samples but for big samples the Anderson-Darling method  $AD$  and the right tail Anderson-Darling method  $RAD$  are better than all the other methods even the  $MLE$ . A new statistic which can validate effectively the extension exponential distribution is provided. By using this criteria test, we will not need to compare this model with the possible alternatives in the analysis.

### REFERENCES

- [1] Akinsete A., Famoye F., Lee C.(2008), "The beta-Pareto distribution". Statistics, 42(6), 547-563.
- [2] Alzaatreh A., Lee C., Famoye F., Ghosh I."The generalized Cauchy family of distributions with applications". Journal of Statistical Distributions and Applications 3 (1), 1-16.
- [3] Alzaghaf A., Felix F., Carl L. (2013). "Exponentiated T-X Family of Distributions with Some Applications". International Journal of Statistics and Probability. 2(3). 31-49.
- [4] Anderson.T.W. and Darling.D.A.(1952) , "Asymptotic theory of certain "goodness of fit" criteria based on stochastic processes". The Annals of Mathematical Statistics, 23(2) , 193-212.

- [5] Boos.D.D. (1981), "Minimum distance estimators for location and goodness of fit". *Journal of the American Statistical Association*, 76 , 663–670.
- [6] Bourguignon M., Silva R.B., Cordeiro G.M.(2014)."The Weibull-G Family of Probability Distributions".*Journal of Data Science* 12, 53-68.
- [7] Cheng R.,Amin N. (1983), "Estimating parameters in continuous univariate distributions with a shifted origin". *Journal of the Royal Statistical Society, Series B (Methodological)*, 45 , 394–403.
- [8] Cordeiro M., Castro M. (2011), "A new family of generalized distributions". *Journal of Statistical Computation and Simulation*. 81(7). 883-898.<http://dx.doi.org/10.1080/00949650903530745>.
- [9] Dey.S. , Dey.T. and Kundu.D.(2014), "Two-parameter Rayleigh distribution: different methods of estimation". *American Journal of Mathematical and Management Sciences*, 33 , 55–74.
- [10] Gupta, R. D. and kundu, D.(2001), "Exponentiated Exponential Family:An alternative to Gamma and Weibull Distributions", *Biometrical Journal*,43(1),pp.117-130.
- [11] kumar D.,Singh U.,Singh S.k , Mukherjee S.(2017),"The New Probability Distribution: An Aspect to a Life Time Distribution".*International Journal of Mathematical Science Letters*.
- [12] Linhart H. and Zucchini W. (1986),*Model selection*,John Wiley Sons. 12.
- [13] Louzada.F., Ramos.P.L. and Perdon.G.S.(2016) , "Different estimation procedures for the parameters of the extended exponential geometric distribution for medical data". *Computational and Mathematical Methods in Medicine*. Article ID 8727951, 12 pages. doi:10.1155/2016/8727951.
- [14] Luceno A.(2006), "Fitting the generalized Pareto distribution to data using maximum goodness-of-fit estimators". *Computational Statistics & Data Analysis*, 51 , 904–917.
- [15] MacDonald P. D. M. (1971), "Comment on an estimation procedure for mixtures of distributions by Choi and Bulgren". *Journal of Royal Statistical Society: Series B*,33(2), 326-329.
- [16] Mudholkar G.S., Huston A.D.(1996),"The exponentiated Weibull family:Some properties and a flood data application".*Communications in Statistics-Theory and Methods* Taylor and Francis.25(12), 3059-3083.
- [17] Nadarajah S., Haghighi F."An extension of the exponential distribution".*A Journal of Theoretical and Applied Statistics* Volume 45, 2011 - Issue 6.
- [18] Nikulin M. S.(1973), "Chi-square test for normality," in: *International Vilnius Conference on Probability Theory and Mathematical Statistics*, Vol. 2 , pp. 119–122.
- [19] Ramnaby B. (1984), "The Maximum Spacing Method: An Estimation Method Related to the Maximum Likelihood Method". *Scandinavian Journal of Statistics*, 11, 93-112.
- [20] Ramos P. and Louzada F.(2016), "The generalized weighted Lindley distribution", *Properties, estimation and applications*. *CogentMathematics*, 3 , 1–18.
- [21] Ramos P., Louzada F.(2016), "The generalized weighted Lindley distribution", *Properties, estimation and applications*. *CogentMathematics*, 3 ,1–18.
- [22] Rao K. C.,Robson D.S ,(1974). "A Chi-square statistic for goodness-of-fit for tests with in the exponential family".*Communications in statistics* 3:1139-53.
- [23] Seddik-Ameur N.,Chouia S.(2014)."A Goodness-of-fit Test for a Competing Risk Model".*International Journal of Applied Mathematics and Statistics*,Vol. 52 , Issue Number: 2.
- [24] Tilbi D.,Seddik-Ameur N.(2017)."Chi-squared goodness-of-fit tests for the generalized Rayleigh distribution".*Journal of Statistical Theory and Practice* volume 11, pages594–603 .
- [25] Treidi W., Seddik-Ameur N."NRRStatistic for the extension Weibull distribution".*Global Journal of Pure and Applied Mathematics*. ISSN 0973-1768Volume 12, Number 4 (2016), pp. 2809–2818.
- [26] Van der Vaart, A.w.(1998).*Asymptotic statistics*.Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge, UK: Cambridge University Press.
- [27] Weber.M. D., Leemis.L. M. and Kincaid.R. K.(2006) , "Minimum Kolmogorov-Smirnov test statistic parameter estimates". *Journal of Statistical Computation and Simulation*, Vol. 76, No. 3 , 195–206.