

СИБИРСКИЕ ЭЛЕКТРОННЫЕ МАТЕМАТИЧЕСКИЕ ИЗВЕСТИЯ

Siberian Electronic Mathematical Reports

<http://semr.math.nsc.ru>

Том 20, №2, стр. 1001–1012 (2023)

УДК ????

DOI 10.33048/semi.2023.20.061

MSC ??X??

LAGUERRE EXPANSIONS OF C -REGULARIZED SEMIGROUPS FUNCTIONS

Y. BAJJOU, A. BLALI, AND A. EL AMRANI

ABSTRACT. The aim of this paper is to approximate the exponentially bounded C -regularized semigroups function by the Laguerre series, recalling the notions and the results used.

Keywords: Laguerre functions, C -regularized semigroup, C_0 -semigroup.

Mathematics Subject Classification: 47D60, 47A10, 46A32

1. INTRODUCTION AND PRELIMINARIES

The series expansion of Laguerre orthogonal polynomials have been an important tool in mathematical physics, in problems involving the integration of Helmholtz's equation in parabolic coordinates, in the theory of the Hydrogen atom, in the theory of propagation of electromagnetic waves along transmission lines terminated by a lumped inductance [8]. The study of sufficient conditions for the convergence of Laguerre series has been the subject of numerous works, for more details see [14], [3], [13], [8] and [2].

In 2014, Abadias and Miana studied in their article [1] the Laguerre expansion of C_0 -semigroups and Resolvent Operators. In this work we will be interested in Laguerre expansion of C -regularized semigroups function, starting with reminding the notations, concepts and results used.

Throughout this paper E denotes a non-trivial complex Banach space, $\mathfrak{F}(E, F)$ denotes the set of all applications from E to another Banach space F , $B(E)$ denotes the space of all bounded linear operators from E into itself, and $L_{loc}^1(E)$ the set of all $f \in \mathfrak{F}(\mathbb{R}, E)$ locally integrable. For a closed linear operator A on E , $\mathcal{D}(A)$, $R(A)$ and $\rho(A)$ denote its domain, range and resolvent set, respectively. $\mathcal{D}(A)$ equipped with the graph norm $\|x\|_{\mathcal{D}(A)} = \|x\|_E + \|Ax\|_E$ becomes a Banach space. Throughout

BAJJOU, Y., BLALI, A., EL AMRANI, A. LAGUERRE EXPANSIONS OF C -REGULARIZED SEMIGROUPS FUNCTIONS.

© 2023 BAJJOU, Y., BLALI, A., EL AMRANI, A..

The work is supported by RFFI (grant 03-01-11111).

Received April, 20, 2023, published November, xx, 2023.

this paper, $C \in B(E)$ will be an injective operator. The C -resolvent set of A , denoted by $\rho_C(A)$, is defined by $\rho_C(A) := \{\lambda \in \mathbb{C} \mid R(C) \subseteq R(\lambda I - A) \text{ and } \lambda I - A \text{ is injective}\}$ and if $\lambda \in \rho_C(A)$ then we denote by $R_C(\lambda, A) = (\lambda I - A)^{-1}C$ the C -resolvent.

1.1. Laguerre functions and Laguerre expansions on Banach spaces. For all $n \in \mathbb{N}$, and arbitrary real $\alpha > -1$ the classical Laguerre polynomial is defined by Rodrigues formula:

$$(\forall x \in \mathbb{R}) \quad \phi_{n,\alpha}(x) = e^x \frac{x^{-\alpha}}{n!} \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha}),$$

$\phi_{n,\alpha}$ is a polynomial with the degree n , the same parity as n , whose highest monomial degree is $\frac{(-1)^n}{n!} x^n$ and have real coefficients. Furthermore, they verify the following condition of orthogonality with respect to gamma density $x \mapsto x^\alpha e^{-x}$ on $[0, +\infty[$:

$$\int_{\mathbb{R}^+} \phi_{n,\alpha}(x) \phi_{m,\alpha}(x) x^\alpha e^{-x} dx = \delta_{n,m} \frac{\Gamma(n + \alpha + 1)}{n!},$$

where $\delta_{n,m}$ is the Kronecker delta. We also have recurrence relations, differential equations and the estimates:

$$(1) \quad (\forall x \in \mathbb{R}^+) \quad (\exists c_x > 0) \quad (\exists n_0 \in \mathbb{N}) \quad (\forall n \geq n_0), \quad |\phi_{n,\alpha}(x)| \leq c_x n^{\frac{\alpha}{2}}.$$

For more details of the classical theory of orthogonal polynomials see [8], [14], [3], [13], [1] and [2].

The Laguerre functions on $\mathbb{R}^+$ are defined by:

$$(2) \quad \begin{aligned} \varphi_{n,\alpha}(x) &= \sqrt{\frac{n!}{\Gamma(n + \alpha + 1)}} \phi_{n,\alpha}(x) x^{\frac{\alpha}{2}} e^{-\frac{x}{2}} = \\ &= \frac{1}{\sqrt{n! \Gamma(n + \alpha + 1)}} x^{\frac{-\alpha}{2}} e^{\frac{x}{2}} \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha}), \end{aligned}$$

$(\varphi_{n,\alpha})_{n \in \mathbb{N}}$ is an orthonormal basis in the Hilbert space $L^2(\mathbb{R}^+)$ and satisfies some recurrence relations, equalities and inequalities. For more details see [14], [3], [1], [13] and [2].

For $n \in \mathbb{N}$ and arbitrary real $\alpha > -1$, we denote by $\psi_{n,\alpha}$, the function on $\mathbb{R}^+$ defined by:

$$(3) \quad (\forall x \in \mathbb{R}^+) \quad \psi_{n,\alpha}(x) = \frac{n!}{\Gamma(n + \alpha + 1)} x^\alpha e^{-x} \phi_{n,\alpha}(x) = \frac{1}{\Gamma(n + \alpha + 1)} \frac{d^n}{dx^n} (x^{n+\alpha} e^{-x}),$$

$\psi_{n,\alpha}$ satisfies recurrence relations and differential equations, for example:

$$(4) \quad (\forall (n, m) \in \mathbb{N}^2) \quad \psi_{n,\alpha}^{(m)} = \psi_{n+m,\alpha-m}.$$

And the following useful inequality:

$$(5) \quad (\forall n \geq 1) \quad \|\psi_{n,\alpha}\|_1 \leq \frac{c_\alpha}{n^{\frac{\alpha}{2}}}.$$

For more details see [1] and [8].

The most important property of the family $(\psi_{n,\alpha})_{n \in \mathbb{N}}$ is that if $f : \mathbb{R}_+^* \rightarrow E$ be a differentiable function such that $\int_0^{+\infty} e^{-t} t^\alpha \|f(t)\|^2 dt < +\infty$, then the series $\sum_{n \in \mathbb{N}} c_n(f) \phi_{n,\alpha}(t)$ converges pointwise to f on $\mathbb{R}_+^*$, where

$$c_n(f) = \int_0^{+\infty} \psi_{n,\alpha}(t) f(t) dt.$$

For more details see [8], [3] and [13].

1.2. C -regularized semigroup. A family of operators $(T(t))_{t \geq 0}$ in $B(E)$ is called exponentially bounded C -regularized semigroup or exponentially bounded C -semigroup on E , if

- (1) $T(t+s)C = T(t)T(s)$ for all $t, s \in \mathbb{R}^+$.
- (2) $T(0) = C$.
- (3) The function $t \mapsto T(t)x$ is continuous on $\mathbb{R}^+$ for any $x \in E$.
- (4) $(\exists M \geq 0) (\exists \omega \geq 0) : \|T(t)\| \leq Me^{\omega t}$, for all $t \geq 0$ (exponentially bounded condition).

Its generator W is defined by

$$\mathcal{D}(W) = \left\{ x \in E : \lim_{s \rightarrow 0^+} \frac{T(s)x - Cx}{s} \text{ exists in } R(C) \right\}$$

and

$$(\forall x \in \mathcal{D}(W)) \quad Wx = C^{-1} \lim_{s \rightarrow 0^+} \frac{T(s)x - Cx}{s}.$$

In all that follows, $(T(t))_{t \geq 0} \subset B(E)$ is exponentially bounded C -regularized semigroup on E with generator $(W, \mathcal{D}(W))$ such that

- (6) $(\exists M > 0) (\exists \omega \geq 0) : (\forall t \geq 0) \|T(t)\| \leq Me^{\omega t}$.

We present some known facts about C -semigroup and its generator, which will be used in the sequel (see [7], [5], [6], [7], [9], [10], [11], [12], [15] and [16] for more details):

- By the property 1, we conclude that $T(t)T(s) = T(s)T(t)$ for all $t, s \geq 0$, this means that $T(t)x \in \mathcal{D}(W)$ and $WT(t)x = T(t)Wx$, for all $t \geq 0$ and $x \in \mathcal{D}(W)$.
- $\int_0^t T(s)x ds \in \mathcal{D}(W)$ and $W \int_0^t T(s)x ds = T(t)x - Cx$ for every $x \in E$ and $t \geq 0$, which implies that for each $x \in \mathcal{D}(W)$, $u := T(\cdot)x$ is of class $C^1(\mathbb{R}^+, E)$ and solves the Abstract Cauchy problem

$$((ACP(W, x, 0)_1) \left\{ \begin{array}{l} u'(t) = Wu(t), \quad t \in \mathbb{R}^+ \\ u(0) = Cx. \end{array} \right.$$

- $\lim_{h \rightarrow 0^+} \frac{1}{h} \int_t^{t+h} T(s)x ds = T(t)x$ for all $x \in E$ and $t \geq 0$, this means that W is a closed linear operator with $R(C) \subset \overline{D(W)}$.
- The C -resolvent operator $R_C(\lambda, W)$ is analytic in the C -resolvent set $\rho_C(W)$ and

$$(7) \quad \frac{d^n}{d\lambda^n} (R_C(\lambda, W)) = (-1)^n n! (\lambda I - W)^{-n-1} C \text{ for all } n \in \mathbb{N}.$$

- $W = C^{-1}WC$, $(\omega, +\infty) \subset \rho_C(W)$ and $R_C(\lambda, W)x = \int_0^{+\infty} e^{-\lambda t} T(t)x dt$ for $\lambda > \omega$ and $x \in E$. For every $\lambda > \omega$ and $n \in \mathbb{N}$, $R(C) \subset D((\lambda I - W)^{-n})$ and

$$(8) \quad (\lambda I - W)^{-n} Cx = \frac{1}{(n-1)!} \int_0^{+\infty} t^{n-1} e^{-\lambda t} T(t)x dt,$$

which implies

$$(9) \quad \|(\lambda - \omega)^n (\lambda I - W)^{-n} C\| \leq M.$$

please delete this parenthesis

- Let $\alpha > 0$ and $\lambda > 0$. We have $\int_0^{+\infty} t^{\alpha-1} e^{-\lambda t} dt = \frac{\Gamma(\alpha)}{\lambda^\alpha}$ and the family of operators $(e^{-\omega t} T(t))_{t \geq 0}$ is uniformly bounded C -semigroup with the generator $W - \omega I$. Then we can define the fractional power of C -resolvent operator (see [5] for more details) as below :

(10)

$$(\lambda I - (W - \omega I))^{-\alpha} Cx = ((\lambda + \omega)I - W)^{-\alpha} Cx := \frac{1}{\Gamma(\alpha)} \int_0^{+\infty} t^{\alpha-1} e^{-\lambda t} e^{-\omega t} T(t) x dt,$$

for all $x \in E$. With a simple verification, $(\lambda I - (W - \omega I))^{-\alpha} C$ is bounded linear operator.

2. MAIN RESULTS

Theorem 1. Let $(T(t))_{t \in \mathbb{R}^+}$ be an exponentially bounded C -semigroup in a Banach space E with generator $(W, \mathcal{D}(W))$, $q \in \mathbb{N}$ such that $q > 2$ and $\alpha > -1$. Then

- (1) For any $n \in \mathbb{N}$ and $x \in E$, we have

$$\int_0^{+\infty} \psi_{n,\alpha}(t) e^{-\omega t} T(t) x dt = (\omega I - W)^n ((\omega + 1)I - W)^{-n-\alpha-1} Cx.$$

- (2) For $x \in \mathcal{D}(W)$, we have :

- (a) $(t > 0)$ $T(t)x = e^{\omega t} \sum_{n=0}^{+\infty} (\omega I - W)^n ((\omega + 1)I - W)^{-n-\alpha-1} \phi_{n,\alpha}(t) Cx$.
- (b) For each $t > 0$ there is $n_0 \in \mathbb{N}$ such that for all integer n with $n \geq n_0$ and $x \in \mathcal{D}(W^q)$, we have

$$\| T(t)x - e^{\omega t} \sum_{m=0}^n (\omega I - W)^m ((\omega + 1)I - W)^{-m-\alpha-1} Cx \phi_{m,\alpha}(t) \| \leq$$

$$\leq \frac{c_{t,\alpha,q} \| (W - \omega I)^q x \|}{n^{\frac{q}{2}-1}}.$$

please delete this comma

where $c_{t,\alpha,q}$ is a constant which depends only on $t > 0$, α and q .

Proof.

Throughout the proof, α is an arbitrary real such that $\alpha > -1$.

- (1) Let $n \in \mathbb{N}$ and $x \in E$. The function $t \mapsto \psi_{n,\alpha}(t) T(t)x$ is continuous on $\mathbb{R}^+$ and integrable in the sens of Bochner. Put $I := \int_0^{+\infty} \psi_{n,\alpha}(t) (e^{-\omega t} T(t)) x dt$,

then

$$\begin{aligned}
 I &:= \int_0^{+\infty} \psi_{n,\alpha}(t)(e^{-\omega t}T(t))x dt \\
 &= \frac{1}{\Gamma(n+\alpha+1)} \int_0^{+\infty} \frac{d^n}{dt^n} (t^{n+\alpha}e^{-t})(e^{-\omega t}T(t))x dt \\
 &= \frac{1}{\Gamma(n+\alpha+1)} \left(\left[\frac{d^{n-1}}{dt^{n-1}} (t^{n+\alpha}e^{-t})(e^{-\omega t}T(t))x \right]_0^{+\infty} - \right. \\
 &\quad \left. - \int_0^{+\infty} \frac{d^{n-1}}{dt^{n-1}} (t^{n+\alpha}e^{-t}) \frac{d}{dt} (e^{-\omega t}T(t))x dt \right) \\
 &= \frac{1}{\Gamma(n+\alpha+1)} \left(0 - \int_0^{+\infty} \frac{d^{n-1}}{dt^{n-1}} (t^{n+\alpha}e^{-t})(W - \omega I)(e^{-\omega t}T(t))x dt \right) \\
 &= \frac{-1}{\Gamma(n+\alpha+1)} \left((W - \omega I) \int_0^{+\infty} \frac{d^{n-1}}{dt^{n-1}} (t^{n+\alpha}e^{-t})(e^{-\omega t}T(t))x dt \right) \\
 &= (-1)^n (W - \omega I)^n \left(\frac{1}{\Gamma(n+\alpha+1)} \int_0^{+\infty} (t^{n+\alpha}e^{-t})(e^{-\omega t}T(t))x dt \right) \\
 &= (\omega I - W)^n \left(\frac{1}{\Gamma(n+\alpha+1)} \int_0^{+\infty} (t^{(n+\alpha+1)-1}e^{-t})(e^{-\omega t}T(t))x dt \right) \\
 &= (\omega I - W)^n (I - (W - \omega I))^{-n-\alpha-1} Cx \\
 &= (\omega I - W)^n ((\omega + 1)I - W)^{-n-\alpha-1} Cx
 \end{aligned}$$

(2) Let $x \in \mathcal{D}(W)$.

- (a) Let's remember that $e^{-\omega(\cdot)}T(\cdot)x : \mathbb{R}^+ \rightarrow E$ is in $C^1(\mathbb{R}^+, E)$ and

$$\begin{aligned}
 \int_0^{+\infty} e^{-t} t^\alpha \|e^{-\omega t}T(t)x\|^2 dt &\leq \int_0^{+\infty} e^{-t} t^\alpha \|e^{-\omega t}T(t)\|^2 \|x\|^2 dt \\
 &\leq M^2 \|x\|^2 \int_0^{+\infty} e^{-t} t^\alpha dt \\
 &< +\infty.
 \end{aligned}$$

Now, we apply the most important properties of $(\psi_{n,\alpha})_{n \in \mathbb{N}}$ to get the series $\sum_{n \in \mathbb{N}} c_n(e^{-\omega(\cdot)}T(\cdot)x)\phi_{n,\alpha}(\cdot)$, where

$$\begin{aligned}
 c_n((e^{-\omega(\cdot)}T(\cdot))x) &= \int_0^{+\infty} \psi_{n,\alpha}(t)(e^{-\omega t}T(t))x dt \\
 &= (\omega I - W)^n ((\omega + 1)I - W)^{-n-\alpha-1} Cx
 \end{aligned}$$

converges pointwise to $e^{-\omega(\cdot)}T(\cdot)x$ for $t \in \mathbb{R}^+$. Therefore

$$\begin{aligned}
 T(t)x &= e^{\omega t}(e^{-\omega t}T(t)x) \\
 &= e^{\omega t} \sum_{n=0}^{+\infty} (\omega I - W)^n ((\omega + 1)I - W)^{-n-\alpha-1} Cx \phi_{n,\alpha}(t).
 \end{aligned}$$

- (b) Let $t \geq 0$ and $q \in \mathbb{N}$ such that $q > 2$.

Since $\lim_{n \rightarrow +\infty} \left(\frac{n}{n-q}\right)^{\frac{\alpha}{2}} = 1$, we have

$$(\exists N_1 \in \mathbb{N}) (\forall n \geq \max(q+1, N_1)), \left(\frac{n}{n-q}\right)^{\frac{\alpha}{2}} \leq 2.$$

The Riemann series $\sum_{m \geq q+1} \frac{1}{(m-q)^{\frac{q}{2}}}$ is convergent, hence

$$\sum_{m=n+1}^{+\infty} \frac{1}{(m-q)^{\frac{q}{2}}} \sim_{n \rightarrow +\infty} \frac{1}{(\frac{q}{2}-1)} \frac{1}{(n-q)^{\frac{q}{2}-1}}.$$

We have

$$\frac{1}{(\frac{q}{2}-1)} \frac{1}{(n-q)^{\frac{q}{2}-1}} \sim_{n \rightarrow +\infty} \frac{1}{(\frac{q}{2}-1)} \frac{1}{n^{\frac{q}{2}-1}},$$

thus

$$(\exists N_2 \in \mathbb{N}) (\forall n \geq \max(q+1, N_1, N_2)), \quad \sum_{m=n+1}^{+\infty} \frac{1}{m^{\frac{q}{2}}} \leq \frac{1}{(\frac{q}{2}-1)} \frac{1}{n^{\frac{q}{2}-1}}.$$

By (1),

$$(\exists N_3 \in \mathbb{N}) (\exists c_t > 0) : (\forall n \geq \max(N_1, N_2, N_3, q+1)) \mid \phi_{n,\alpha}(t) \mid \leq c_t n^{\frac{\alpha}{2}}.$$

Put $n_0 = \max(N_1, N_2, N_3, q+1)$, then for $n \geq n_0$ and $x \in D(W^q)$

$$\begin{aligned} J &:= \int_0^{+\infty} \psi_{n,\alpha}(t)(e^{-\omega t}T(t))x dt \\ &= \int_0^{+\infty} \frac{d^q}{dt^q}(\psi_{n-q,\alpha+q}(t))(e^{-\omega t}T(t))x dt \quad (\text{according to (4)}) \\ &= \left[\frac{d^{q-1}}{dt^{q-1}}(\psi_{n-q,\alpha+q}(t))(e^{-\omega t}T(t))x \right]_0^{+\infty} - \\ &\quad - \int_0^{+\infty} \frac{d^{q-1}}{dt^{q-1}}(\psi_{n-q,\alpha+q}(t)) \frac{d}{dt}(e^{-\omega t}T(t))x dt \\ &= 0 - \int_0^{+\infty} \frac{d^{q-1}}{dt^{q-1}}(\psi_{n-q,\alpha+q}(t))(W - \omega I)(e^{-\omega t}T(t))x dt \\ &= -(W - \omega I) \int_0^{+\infty} \frac{d^{q-1}}{dt^{q-1}}(\psi_{n-q,\alpha+q}(t))(e^{-\omega t}T(t))x dt \\ &= (\omega I - W) \int_0^{+\infty} \frac{d^{q-1}}{dt^{q-1}}(\psi_{n-q,\alpha+q}(t))(e^{-\omega t}T(t))x dt \\ &= (\omega I - W)^q \int_0^{+\infty} \psi_{n-q,\alpha+q}(t)(e^{-\omega t}T(t))x dt \\ &= \int_0^{+\infty} \psi_{n-q,\alpha+q}(t)(e^{-\omega t}T(t))(\omega I - W)^q x dt. \end{aligned}$$

So

$$\begin{aligned}
\| (\omega I - W)^n ((\omega + 1)I - W)^{-n-\alpha-1} Cx \| &= \left\| \int_0^{+\infty} \psi_{n,\alpha}(t) (e^{-\omega t} T(t)) x dt \right\| \\
&= \left\| \int_0^{+\infty} (\psi_{n-q,\alpha+q}(t) (e^{-\omega t} T(t)) (W - \omega I)^q x) dt \right\| \\
&\leq \int_0^{+\infty} | \psi_{n-q,\alpha+q}(t) | \| e^{-\omega t} T(t) \| dt \| (W - \omega I)^q x \| \\
&\leq \| (W - \omega I)^q x \| M \| \psi_{n-q,\alpha+q} \|_1 \\
&\leq \| (W - \omega I)^q x \| \frac{c_{\alpha+q}}{(n-q)^{\frac{\alpha+q}{2}}} \text{ (according to (5))}.
\end{aligned}$$

On the other hand, we are looking to estimate the quantity

$$\begin{aligned}
(11) \quad & e^{\omega t} \left\| \sum_{m=0}^{+\infty} (\omega I - W)^m ((\omega + 1)I - W)^{-m-\alpha-1} Cx \phi_{m,\alpha}(t) - \right. \\
& \left. - \sum_{m=0}^n (\omega I - W)^m ((\omega + 1)I - W)^{-m-\alpha-1} Cx \phi_{m,\alpha}(t) \right\|.
\end{aligned}$$

We have

$$\begin{aligned}
(11) &\leq \sum_{m=n+1}^{+\infty} \| e^{\omega t} (\omega I - W)^m ((\omega + 1)I - W)^{-m-\alpha-1} Cx \phi_{m,\alpha}(t) \| \\
&\leq e^{\omega t} \sum_{m=n+1}^{+\infty} \| (\omega I - W)^m ((\omega + 1)I - W)^{-m-\alpha-1} Cx \| | \phi_{m,\alpha}(t) | \\
&\leq e^{\omega t} \sum_{m=n+1}^{+\infty} \| (W - \omega I)^q x \| \frac{c_{\alpha+q}}{(m-q)^{\frac{\alpha+q}{2}}} c_t m^{\frac{\alpha}{2}} \\
&\leq \| (W - \omega I)^q x \| e^{\omega t} c_{\alpha+q} \sum_{m=n+1}^{+\infty} \frac{m^{\frac{\alpha}{2}}}{(m-q)^{\frac{\alpha+q}{2}}} \\
&\leq \| (W - \omega I)^q x \| c'_t c_{\alpha+q} \sum_{m=n+1}^{+\infty} \frac{(m-q)^{\frac{\alpha}{2}}}{(m-q)^{\frac{\alpha+q}{2}}} \frac{m^{\frac{\alpha}{2}}}{(m-q)^{\frac{\alpha}{2}}} \\
&\leq \| (W - \omega I)^q x \| c'_t c_{\alpha+q} \sum_{m=n+1}^{+\infty} \frac{(m-q)^{\frac{\alpha}{2}}}{(m-q)^{\frac{\alpha+q}{2}}} \left(\frac{m}{m-q} \right)^{\frac{\alpha}{2}} \\
&\leq \| (W - \omega I)^q x \| c'_t c_{\alpha+q} \sum_{m=n+1}^{+\infty} \frac{1}{(m-q)^{\frac{q}{2}}} \times 2 \\
&\leq \| (W - \omega I)^q x \| c'_t c_{\alpha+q} \frac{1}{(\frac{q}{2}-1)} \frac{1}{n^{\frac{q}{2}-1}},
\end{aligned}$$

where

$$\begin{aligned}
&\| T(t)x - e^{\omega t} \sum_{m=0}^n (\omega I - W)^m ((\omega + 1)I - W)^{-m-\alpha-1} Cx \phi_{m,\alpha}(t) \| \leq \\
&\leq \frac{c_{t,\alpha,q} \| (W - \omega I)^q x \|}{n^{\frac{q}{2}-1}}.
\end{aligned}$$

Theorem 2. *Let $(T(t))_{t \in \mathbb{R}^+}$ be an exponentially bounded C -semigroup in Banach space E with generator $(W, \mathcal{D}(W))$ and $\alpha \in \mathbb{R}$.*

(1) *For $x \in E$, $\alpha > 0$ and $n \in \mathbb{N}$, we have*

$$\int_0^{+\infty} \psi_{n,\alpha}(t) R_C(t, W - \omega I) x dt = \int_0^{+\infty} \frac{s^n}{(s+1)^{n+\alpha+1}} e^{-\omega s} T(s) x ds.$$

(2) *For $x \in E$ and $\alpha > 1$, we have*

$$R_C(t, W - \omega I) x = \sum_{n=0}^{+\infty} \left(\int_0^{+\infty} \frac{s^n e^{-\omega s}}{(s+1)^{n+\alpha+1}} T(s) x ds \right) \phi_{n,\alpha}(t), \text{ for all } t > 0.$$

Proof.

(1) Let $x \in E$, $\alpha > 0$ and $n \in \mathbb{N}$.

For all $t > 0$, $t + \omega \in \rho_C(W)$, so the function $t \mapsto R_C(t, W - \omega I) = R_C(t + \omega, W)$ is analytic in $\mathbb{R}_+^*$.

In the other hand, by equation (9), we know that $\| R_C(t + \omega, W) \| \leq \frac{M}{t}$ for all $t > 0$, so

$$\begin{aligned} \int_0^{+\infty} \| \psi_{n,\alpha}(t) R_C(t, W - \omega I) x \| dt &= \int_0^{+\infty} | \psi_{n,\alpha}(t) | \| (R_C(t, W - \omega I) x \| dt \\ &\leq \int_0^{+\infty} \frac{n!M}{\Gamma(n+\alpha+1)t} e^{-t} t^\alpha | \phi_{n,\alpha}(t) | dt \\ &\leq \frac{n!M}{\Gamma(n+\alpha+1)} \int_0^{+\infty} e^{-t} t^{\alpha-1} | \phi_{n,\alpha}(t) | dt \\ &< +\infty. \end{aligned}$$

Put $H := \Gamma(n + \alpha + 1) \int_0^{+\infty} \psi_{n,\alpha}(t) R_C(t, W - \omega I) x dt$, then

$$\begin{aligned}
H &= \Gamma(n + \alpha + 1) \int_0^{+\infty} \psi_{n,\alpha}(t) R_C(t, W - \omega I) x dt \\
&= \int_0^{+\infty} \frac{d^n}{dt^n} (e^{-t} t^{n+\alpha}) R_C(t, W - \omega I) x dt \\
&= \underbrace{\left[\frac{d^{n-1}}{dt^{n-1}} (e^{-t} t^{n+\alpha}) R_C(t, W - \omega I) x \right]_0^{+\infty}}_{=0} - \\
&\quad - \int_0^{+\infty} \frac{d^{n-1}}{dt^{n-1}} (e^{-t} t^{n+\alpha}) \frac{d}{dt} (R_C(t, W - \omega I) x) dt \\
&= \left[-\frac{d^{n-2}}{dt^{n-2}} (e^{-t} t^{n+\alpha}) \frac{d}{dt} (R_C(t, W - \omega I) x) \right]_0^{+\infty} + \\
&\quad + (-1)^2 \int_0^{+\infty} \frac{d^{n-2}}{dt^{n-2}} (e^{-t} t^{n+\alpha}) \frac{d^2}{dt^2} (R_C(t, W - \omega I) x) dt \\
&= (-1)^n \int_0^{+\infty} e^{-t} t^{n+\alpha} \frac{d^n}{dt^n} (R_C(t, W - \omega I) x) dt \quad (\text{integration by parts}) \\
&= (-1)^n \int_0^{+\infty} e^{-t} t^{n+\alpha} (-1)^n n! ((t + \omega)I - W)^{-n-1} C x dt \\
&= \int_0^{+\infty} e^{-t} t^{n+\alpha} (n!) \frac{1}{n!} \int_0^{+\infty} s^n e^{-ts} e^{-\omega s} T(s) x ds dt \\
&= \int_0^{+\infty} s^n e^{-\omega s} T(s) x \left(\int_0^{+\infty} t^{n+\alpha} e^{-t} e^{-ts} dt \right) ds \quad (\text{Fubini's Theorem}) \\
&= \int_0^{+\infty} s^n e^{-\omega s} T(s) x \left(\int_0^{+\infty} t^{n+\alpha} e^{-(s+1)t} dt \right) ds \\
&= \int_0^{+\infty} s^n e^{-\omega s} T(s) x \left(\int_0^{+\infty} \frac{u^{n+\alpha}}{(s+1)^{n+\alpha+1}} e^{-u} \frac{du}{s+1} \right) ds \quad (u = (s+1)t) \\
&= \int_0^{+\infty} \frac{s^n}{(s+1)^{n+\alpha+1}} e^{-\omega s} T(s) x \left(\int_0^{+\infty} u^{n+\alpha} e^{-u} du \right) ds \\
&= \int_0^{+\infty} \frac{s^n}{(s+1)^{n+\alpha+1}} e^{-\omega s} T(s) x \Gamma(n + \alpha + 1) ds \\
&= \Gamma(n + \alpha + 1) \int_0^{+\infty} \frac{s^n}{(s+1)^{n+\alpha+1}} e^{-\omega s} T(s) x ds,
\end{aligned}$$

hence the result.

(2) Let $x \in E$ and $\alpha > 1$.

The function $t \mapsto R_C(t, W - \omega I)x = R_C(t + \omega, W)x$ is differentiable in

$\mathbb{R}_+^*$ (because it's analytic in $\mathbb{R}_+^*$), and

$$\begin{aligned}
 \int_0^{+\infty} t^\alpha e^{-t} \| R_C(t, W - \omega I)x \|^2 dt &= \int_0^{+\infty} t^\alpha e^{-t} \| ((t + \omega)I - W)^{-1}Cx \|^2 dt \\
 &\leq \int_0^{+\infty} t^\alpha e^{-t} \frac{M^2}{t^2} \| x \|^2 dt \text{ (equation (9))} \\
 &\leq M^2 \| x \|^2 \int_0^{+\infty} t^{\alpha-2} e^{-t} dt \\
 &= M^2 \| x \|^2 \Gamma(\alpha - 1) \text{ (because } \alpha > 1) \\
 &< +\infty.
 \end{aligned}$$

Thus, the series $\sum_{n \in \mathbb{N}} c_n(R_C(., W - \omega I)x)\phi_{n,\alpha}$ converges pointwise to $R_C(., W - \omega I)x$ on $\mathbb{R}_+^*$, where

$$\begin{aligned}
 c_n(R_C(., W - \omega I)x) &= \int_0^{+\infty} \psi_{n,\alpha}(t) R_C(t, W - \omega I)x dt \\
 &= \int_0^{+\infty} \frac{s^n}{(s+1)^{n+\alpha+1}} e^{-\omega s} T(s)x ds \text{ (according to Theorem 2(1))}.
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 (\forall t > 0) \quad R_C(t, W - \omega I)x &= ((t + \omega)I - W)^{-1}Cx = \\
 &= \sum_{n=0}^{+\infty} \left(\int_0^{+\infty} \frac{s^n e^{-\omega s}}{(s+1)^{n+\alpha+1}} T(s)x ds \right) \phi_{n,\alpha}(t).
 \end{aligned}$$

Example 1. Let $m : \mathbb{R} \rightarrow \mathbb{R}^-$ be an even measurable function. In the Banach space $L^1(\mathbb{R})$, we consider the family $T := (T(t))_{t \geq 0} \subset \mathfrak{F}(L^1(\mathbb{R}))$ defined by $\forall t \geq 0$, $T(t) : L^1(\mathbb{R}) \rightarrow L^1(\mathbb{R})$, $f \mapsto T(t)(f) : \mathbb{R} \rightarrow \mathbb{R}$, $s \mapsto T(t)(f)(s) = e^{t.m(s)} f(-s)$. Clearly, $(T(t))_{t \geq 0} \subset B(L^1(\mathbb{R}))$. If we put $T(0) = C$, then the family of operators $(T(t))_{t \geq 0}$ is uniformly bounded C -regularized semigroup with generator $(W, \mathcal{D}(W))$ defined by

$$W : \mathcal{D}(W) = \{f \in L^1(\mathbb{R}) / m.f \in L^1(\mathbb{R})\} \rightarrow L^1(\mathbb{R}), \quad f \mapsto W(f) = m.f.$$

Theorem 1 gives for $f \in \mathcal{D}(W)$ and $s, t \in \mathbb{R}^+$:

$$\begin{aligned}
 T(t)(f)(s) &= e^{\omega t} \sum_{n=0}^{+\infty} (\omega I - W)^n ((\omega + 1)I - W)^{-n-\alpha-1} C(f)(s) \phi_{n,\alpha}(t) \\
 &= e^{0 \times t} \sum_{n=0}^{+\infty} (0 \times I - W)^n ((0 + 1)I - W)^{-n-\alpha-1} C(f)(s) \phi_{n,\alpha}(t) \\
 &= \sum_{n=0}^{+\infty} (-m(s))^n (1 - m(s))^{-n-\alpha-1} C(f)(s) \phi_{n,\alpha}(t).
 \end{aligned}$$

$$\text{So } T(t)(f) = \sum_{n=0}^{+\infty} \phi_{n,\alpha}(t) (-m)^n (1 - m)^{-n-\alpha-1} C(f).$$

Theorem 2 gives

$$R_C(t, W)(f)(.) = \sum_{n=0}^{+\infty} \left(\int_0^{+\infty} \frac{s^n}{(s+1)^{n+\alpha+1}} e^{sm(.)} ds \right) C(f)(.) \phi_{n,\alpha}(t).$$

Example 2. The space $X = c_0 = \left\{ (x_k)_{k \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}} \mid \lim_{k \rightarrow +\infty} x_k = 0 \right\}$, equipped with the norm $\| (x_k)_{k \in \mathbb{N}} \|_{\infty} = \max_{k \in \mathbb{N}} |x_k|$ becomes a Banach space. For each $n \in \mathbb{N}$,

let $e_n = (\delta_{n,k})_{k \in \mathbb{N}}$ be an element of X . Since for all $x = (x_k)_{k \in \mathbb{N}} \in X$, $x = \sum_{k=0}^{+\infty} x_k e_k$, we have $X = \text{span}\{e_n/n \in \mathbb{N}\}$. Considering the family of operators $(T(t))_{t \geq 0}$ defined by:

$$\text{for all } t \in \mathbb{R}^+, \text{ for all } x = \left(\sum_{k=0}^{+\infty} x_k e_k \right) \in X, \quad T(t)x = \sum_{k=0}^{+\infty} e^{-k^2 t} x_k e_k$$

$(T(t))_{t \geq 0}$ is a uniformly bounded C_0 -semigroup ($\|T(t)\| \leq 1$) with generator $(W, \mathcal{D}(W))$ such that $\mathcal{D}(W) = \{x = (x_k)_{k \in \mathbb{N}} \in X / (k^2 x_k)_{k \in \mathbb{N}} \in X\}$ and $(\forall x \in \mathcal{D}(W)) \quad Wx = \sum_{k=0}^{+\infty} -k^2 x_k e_k$. Theorem 1 gives for $x = (x_k)_{k \in \mathbb{N}} \in \mathcal{D}(W)$ and for all $t \in \mathbb{R}^+$:

$$\begin{aligned} T(t)(x) &= e^{\omega t} \sum_{n=0}^{+\infty} (\omega I - W)^n ((\omega + 1)I - W)^{-n-\alpha-1}(x) \phi_{n,\alpha}(t) \\ &= e^{0 \times t} \sum_{n=0}^{+\infty} (0 \times I - W)^n ((0 + 1)I - W)^{-n-\alpha-1}(x) \phi_{n,\alpha}(t) \\ &= \sum_{n=0}^{+\infty} (k^{2n} (1 + k^2)^{-n-\alpha-1} x_k)_{k \in \mathbb{N}} \phi_{n,\alpha}(t) \\ &= \sum_{n=0}^{+\infty} \sum_{k=0}^{+\infty} k^{2n} (1 + k^2)^{-n-\alpha-1} x_k e_k \phi_{n,\alpha}(t). \end{aligned}$$

Theorem 2 gives

$$R_C(t, W)x = \sum_{n=0}^{+\infty} \sum_{k=0}^{+\infty} \left(\int_0^{+\infty} \frac{s^n}{(s+1)^{n+\alpha+1}} e^{-k^2 s} ds \right) x_k e_k \phi_{n,\alpha}(t).$$

Acknowledgments The authors are greatly indebted to the editor and the referee for many valuable comments and suggestions which improved the initial draft of this paper.

REFERENCES

- [1] L. Abadias, P.J. Miana, C_0 -semigroups and resolvent operators approximated by Laguerre expansions, J. Approx. Theory, **213** (2017), 1–22. Zbl 1357.33012
- [2] R. ASKEY, S. WAINGER, Mean convergence of expansions in Laguerre and Hermite series, Amer. J. Math., **87** (1965), 695–708. Zbl 0125.31301
- [3] R. deLaubenfels, C -semigroups and the Cauchy problem, J. Funct. Anal., **111**:1 (1993), 44–61. Zbl 0895.47029
- [4] R. DELAUBENFELS, Existence families, functional calculi and evolution equations, Lecture Notes in Mathematics, **1570**, Springer-Verlag, Berlin, 1994. Zbl 0811.47034
- [5] H.O. Fattorini, A note on fractional derivatives of semigroups and cosine functions, Pac. J. Math., **109**:2 (1983), 335–347. Zbl 0495.47026
- [6] M. Kaltenback, F.L. Schwenninger, Generalisations of semigroups of operators in the view of linear relations, Analysis und Scientific Computing der Technischen Universität Wien, Matr. Nr., **0625539**, Vienna University of Technology, Wien.

- [7] M. Kostić, *(a, k)-regularized C-resolvent families: regularity and local properties*, Abstr. Appl. Anal., **2009** (2009), Article ID 858242. Zbl 1200.47059
- [8] N.N. Lebedev, *Special function and their applications*, Prentice Hall, Englewood Cliffs, 1965. Zbl 0131.07002
- [9] M. Li, Q. Zheng, J. Zhang, *Regulrized resolvent families*, Taiwanese J. Math., **11**:1 (2007), 117–133. Zbl 1157.45006
- [10] C. Martínez, M. Sanz, L. Marco, *Fractional powers of operators*, J. Math. Soc. Japan, **40**:2 (1988), 331–347. Zbl 0628.47006
- [11] M. Mosallanezhad, M. Janfada, *On mixed C-semigroups of operators on Banach spaces*, Filomat, **30**:10 (2016), 2673–2682. Zbl 1465.47035
- [12] P. Preda, A. Pogan, C. Preda, *The Perron problem for C-semigroups*, Math. J. Okayama Univ., **46** (2004), 141–151. Zbl 1091.47036
- [13] J.A. SHOHAT, *On the developement of functions in series of orthogonal polynomials*, The university of Pennsylvania, 1935.
- [14] G. Szegő, *Orthogonal polynomials*, American Mathematical Society, New York, 1939. Zbl 0023.21505
- [15] Q. Zheng, *Controllability of a class of linear systems in Banach spaces*, Proc. Am. Math. Soc., **123**:4 (1995), 1241–1251. Zbl 0833.93029
- [16] Q. Zheng, Y. Zhao, *An analyticity criterion for regularized semigroups*, Proc. Am. Math. Soc., **130**:8 (2002), 2271–2274. Zbl 1015.47030

YOUSSEF BAJJOU

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE,
DHAR EL MAHRAZ FACULTY OF SCIENCES,
SIDI MOHAMED BEN ABDELLAH UNIVERSITY,
FES ATLAS, MOROCCO
Email address: youssefbajjou2017@gmail.com

AZIZ BLALI

DEPARTMENT OF MATHEMATICS,
ÉCOLE NORMALE SUPÉRIEURE,
SIDI MOHAMED BEN ABDELLAH UNIVERSITY,
B.P. 5206 BENSOUDA-FES, MOROCCO
Email address: aziz.blali@usmba.ac.ma

ABDELKHALEK EL AMRANI

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE,
DHAR EL MAHRAZ FACULTY OF SCIENCES,
SIDI MOHAMED BEN ABDELLAH UNIVERSITY,
FES ATLAS, MOROCCO
Email address: abdelkhalek.elamrani@usmba.ac.ma