

SPLITTING PARTIALLY COMMUTATIVE LIE ALGEBRAS
INTO DIRECT SUMS

E. N. POROSHENKO

Communicated by P. P. PETROV (filled by Editorial board)

Abstract: In this work, we prove that partially commutative, partially commutative metabelian, or partially commutative nilpotent Lie algebra splits into the direct sum of two subalgebras if and only if the defining graph G of this algebra is such that $\bar{G}$ is not connected.

Keywords: partially commutative (metabelian, nilpotent) Lie algebra, direct sum.

1 Preliminaries

Research into partially commutative structures started in late sixties of the last century. In [1], the notion of a partially commutative monoid was introduced. Further, partially commutative groups (also known as right-angled Artin groups) were studied the most heavily among all partially commutative structures (see, for example [2, 3, 4]). There are so many papers on partially commutative groups that it is impossible to list all of them here. But quite many papers are mentioned in the surveys [5, 6].

POROSHENKO, E.N., SPLITTING PARTIALLY COMMUTATIVE LIE ALGEBRAS INTO DIRECT SUMS.

© 2023 POROSHENKO E. N..

Поступила 1 января 2023 г., опубликована 31 декабря 2023 г.

Partially commutative algebras (both associative and Lie algebras) were studied not so actively. But there are also some results obtained for them, see for example [6].

Splitting an algebra to a direct sum is a very important tool for an investigation into this algebra. If an algebra can be decomposed to a direct sum then one can study each summand separately which is obviously easier than studying the entire algebra. In this paper, we find a criterion for partially commutative Lie algebras in some varieties to be decomposable into a direct sum. This is more or less an analogue of the result for partially commutative groups obtained in [7] for a certain class of varieties of groups.

In this paper by a graph we mean an undirected graph without loops. Denote such graph by $G = \langle A; E \rangle$, where $A = \{a_1, a_2, \dots, a_n\}$ is the finite set of its vertices and $E \subseteq A \times A$ is the set of its edges. We assume below that $n \geq 2$.

If vertices a and b are connected in G we write $a \leftrightarrow b$. Analogously, if $A_1 \subseteq A$ and $a \in A$ is adjacent to all vertices in A_1 then we use the notation $a \leftrightarrow A_1$. Finally, if $A_1, A_2 \subseteq A$ then $A_1 \leftrightarrow A_2$ means $a_1 \leftrightarrow a_2$ for any $a_1 \in A_1$ and $a_2 \in A_2$. Let H be an arbitrary undirected graph. By $V(H)$ and $E(H)$ denote the set of the vertices and the set of the edges of this graph respectively. Next, let $V_1 \subseteq V(H)$. By $H(V_1)$ denote the subgraph of H , generated by the set V_1 .

In this paper, we work with Lie R -algebras, i.e. Lie algebras over R , where R is a domain and we write just “Lie algebra” instead of “Lie R -algebra” for short.

Consider a variety $\mathfrak{M}$ of Lie algebras. A *partially commutative Lie algebra in $\mathfrak{M}$ with a defining graph G* is a Lie algebra $\mathcal{L}_R(A; G)$ defined by

$$\mathcal{L}_R(A; G) = \langle A \mid [x_i, x_j] = 0 \iff \{x_i, x_j\} \in E; \mathfrak{M} \rangle$$

in $\mathfrak{M}$. Thus, in this algebra, the variety identities and the defining relations hold together.

By $\mathcal{L}(A; G)$, $\mathcal{M}(A; G)$, and $\mathcal{N}_m(A; G)$ denote the partially commutative, partially commutative metabelian, and partially commutative nilpotent of degree m Lie R -algebras defined by the graph G respectively. Let us use the common notation $L(A; G)$ for one of the algebras $\mathcal{L}(A; G)$, $\mathcal{M}(A; G)$, or $\mathcal{N}_m(A; G)$. We denote by $F(A)$ the absolutely free algebra over R with the set of generators A , i.e. the algebra of non-commutative non-associative polynomials in the set A such that these polynomials do not have monomials of cumulative degree 0.

Denote by $[u]$ a non-associative monomial in A , i.e. a finite product of elements in A with a parenthesizing defining the order of multiplications on it.

Definition 1. *Multi-degree* of a non-associative monomial $\alpha[u]$ is the vector $\bar{\delta} = (\delta_1, \delta_2, \dots, \delta_n)$ where δ_i is the number of occurrences of a_i in $[u]$.

Definition 2. A non-zero element g of $F(A)$ is called *multi-homogeneous* if g can be represented as a linear combination of Lie monomials of the same multi-degree $\bar{\delta} = (\delta_1, \delta_2, \dots, \delta_n)$.

For a multi-homogeneous element $g \in F(A)$, if $g = \sum_{i=1}^k \alpha_i [u_i]$, where $\alpha_i \in R \setminus \{0\}$ and $[u_i]$ are non-zero monomials in $F(A)$, then set $\text{mdeg}(g) = \text{mdeg}([u_i])$, where $1 \leq i \leq k$.

Let $[u] \in F(A)$ be a monomial such that $\text{mdeg}([u]) = (\delta_1, \delta_2, \dots, \delta_n)$. The set $\{a_i \mid \delta_i \neq 0\}$ is called a *support* and is denoted by $\text{supp}([u])$. For a polynomial $g = \sum_j \alpha_j [u_j]$ in $F(A)$ set $\text{supp}(g) = \bigcup_j \text{supp}([u_j])$.

The notions of the multi-degree and the support can be defined for elements in $L(A; G)$ in an obvious way.

Since identities and relations of $L(A; G)$ are multi-homogeneous, the following statement holds. If g_i 's are multi-homogeneous polynomials of mutually distinct multi-degrees and $0 = \sum_i g_i$ in $L(A; G)$ then $g_i = 0$ in $L(A; G)$ for any i .

Length of a non-associative monomial $[u]$ is $\sum_{i=1}^n \delta_i$, where $(\delta_1, \delta_2, \dots, \delta_n) = \text{mdeg}([u])$. For $h \in L(A; G)$ consider a representation of h as a linear combination of Lie monomials. Of course, these monomials may have different lengths. Let i be any integer positive number. By $\omega_i(h)$ denote the part of this linear combination consisting of all monomials of length i . Then $h = \sum_{i=1}^r \omega_i(h)$ where r is the maximum among lengths of monomials in this linear combination. Furthermore, set $O_k(h) = \sum_{i=1}^k \omega_i(h)$. Finally, denote by $o_k(h)$ the element $h - O_k(h)$.

Let $f, g \in L(A; G)$. We write $f \sim g$ if $\alpha f = \beta g$ in $L(A; G)$ for some $\alpha, \beta \in R \setminus \{0\}$.

We will use the following theorem on centralizers in $\mathcal{L}(A; G)$ [8].

Theorem 3. Let $g \in \mathcal{L}(A; G)$, $H = \overline{G}(\text{supp}(g))$ and $H_1, \dots, H_p$ be connected components of H . Then

- (1) there is a decomposition $g = \sum_{i=1}^p g_i$, where $\text{supp}(g_i) = A(H_i)$ for $i = 1, 2, \dots, p$;
- (2) $C(g)$ consists of all elements of the form $h = \sum_{i=1}^p h_i + h'$ such that for any $i = 1, 2, \dots, p$ if $h_i \neq 0$ then $g_i \sim h_i$ and if $h' \neq 0$ then $\text{supp}(g) \leftrightarrow \text{supp}(h')$.

In metabelian Lie algebras we assume the left-normed parenthesizing and omit all parentheses except the outer pair.

Let us order A in an arbitrary way. For any multi-degree $\bar{\delta} = (\delta_1, \delta_2, \dots, \delta_n)$ consider the set $A(\bar{\delta}) = \{a_i \mid \delta_i \neq 0\}$. Let $H_0, H_1, \dots, H_s$ be the connected components of $G(A(\bar{\delta}))$. Without loss of generality we can assume that the smallest vertex of $A(\bar{\delta})$ lies in $A(H_0)$. Denote this vertex by b . Consider the set of all monomials of the form $[a_{i_1}, a_{i_2}, \dots, a_{i_r}]$, where

- (1) $\text{mdeg}([a_{i_1}, a_{i_2}, \dots, a_{i_r}]) = \bar{\delta}$, in particular $r = \sum_{i=1}^n \delta_i$;
- (2) $a_{i_1} > a_{i_2}, a_{i_2} \leq a_{i_3} \leq \dots \leq a_{i_r}$, so $a_{i_2} = b$;
- (3) a_{i_1} is the largest element of one of the sets $A(H_1), A(H_2), \dots, A(H_n)$.

Denote this set by $B_{\bar{\delta}}(A; G)$ and consider the set $\mathfrak{B}(A; G) = \bigcup_{\bar{\delta}} B_{\bar{\delta}}(A; G)$. The following theorem holds [9].

Theorem 4. *The set $\mathfrak{B}(A; G)$ is a basis of the partially commutative metabelian Lie algebra $\mathcal{M}(A; G)$.*

2 Main Part

Consider a graph G with the set of vertices $A = \{a_1, a_2, \dots, a_n\}$. Assume that $L(A; G) = L_1 \oplus L_2$ be a decomposition of $L(A; G)$ into a direct sum of two subalgebras.

For any $h \in L(A; G)$ the pair $(h_1, h_2) \in L_1 \times L_2$ is *corresponding to h* if $h = h_1 + h_2$. For any h there is a unique pair corresponding to this element. Denote the elements of this pair by $(h)_1$ and $(h)_2$ respectively.

Lemma 5. *Let G be a graph with the set of vertices $A = \{a_1, a_2, \dots, a_n\}$, where $n \geq 2$ and let $L(A; G) = L_1 \oplus L_2$. If $a_i \leftrightarrow a_j$, where $a_i, a_j \in A$, $i \neq j$, then the following statements hold.*

- (1) *If $L(A; G) = \mathcal{L}(A; G)$ then $a_j \notin \text{supp}((a_i)_1)$.*
- (2) *If $L(A; G) = \mathcal{N}_m(A; G)$ for $m \geq 2$ then $a_j \notin \text{supp}(O_{m-1}((a_i)_1))$.*
- (3) *If $L(A; G) = \mathcal{M}(A; G)$ then $a_j \notin \text{supp}(O_2((a_i)_1))$.*

Proof. Let a_i be an arbitrary element in A . Note that if we change $(a_i)_1$ by $(a_i)_2$ in statements (1)–(3) then we obtain the equivalent statements.

Suppose that $(a_i)_1 = \alpha_i a_i + g_i$ and $a_i \notin \text{supp}(\omega_1(g_i))$. Then $(a_i)_2 = (1 - \alpha_i)a_i - g_i$. So, $\text{supp}((a_i)_1) = \text{supp}(g_i) \cup \{a_i\} = \text{supp}((a_i)_2) \cup \{a_i\}$.

Since $(a_i)_l \in L_l$ for $l = 1, 2$, we have $[(a_i)_1, (a_i)_2] = 0$. Therefore,

$$\begin{aligned} 0 &= [\alpha_i a_i + g_i, (1 - \alpha_i)a_i - g_i] = \\ &= -[\alpha_i a_i, g_i] + [g_i, (1 - \alpha_i)a_i] = \\ &= [g_i, a_i]. \end{aligned} \tag{1}$$

Let $L(A; E) = \mathcal{L}(A; E)$. Then (1) implies by Theorem 3 that $a_i \leftrightarrow \text{supp}(g_i)$. Since $a_j \leftrightarrow a_i$ we obtain $a_j \notin \text{supp}(g_i)$, and so $a_j \notin \text{supp}((a_i)_1)$. Thus, the first statement holds.

Denote by $\mathcal{L}_k(A; G)$ the subset of $\mathcal{L}(A; G)$ consisting of all elements g such that $o_k(g) = 0$. Similarly, $\mathcal{M}_k(A; G)$ is the subset of $\mathcal{M}(A; G)$ such that $o_k(g) = 0$ for any $g \in \mathcal{M}_k(A; G)$.

The restriction of the natural homomorphism $\mathcal{L}(A; G) \rightarrow \mathcal{N}_m(A; G)$ to the map $\mathcal{L}_m(A; G) \rightarrow \mathcal{N}_m(A; G)$ is obviously a bijection as well as the restriction of the natural homomorphism $\mathcal{L}(A; G) \rightarrow \mathcal{M}(A; G)$ to the map $\mathcal{L}_3(A; G) \rightarrow \mathcal{M}_3(A; G)$. Since $\mathcal{N}_m(A; G)$ and $\mathcal{M}(A; G)$ are homogeneous (1) implies

$$[O_k(g_i), a_i] = 0$$

for any positive integer k . Consequently, we can use Theorem 3 to complete the proofs of the second and the third statements similarly we did for the first one. $\square$

Lemma 6. *Let G be a graph with the set of vertices $A = \{a_1, a_2, \dots, a_n\}$, where $n \geq 2$. If $\overline{G}$ is connected and $L(A; G) = L_1 \oplus L_2$ then there exists $l \in \{1, 2\}$ such that for any $i \in \{1, 2, \dots, n\}$ the equation $(a_i)_l = a_i + g_i$, where $a_i \notin \text{supp}(\omega_1(g_i))$, holds.*

Proof. For any $i = 1, 2, \dots, n$ we can write $(a_i)_1 = \alpha_i a_i + g_i$, where $a_i \notin \text{supp}(\omega_1(g_i))$ (see the proof of Lemma 5). Then $(a_i)_2 = (1 - \alpha_i)a_i - g_i$.

Since $[(a_i)_1, (a_j)_2] = [(a_j)_1, (a_i)_2] = 0$ for any $i, j \in \{1, 2, \dots, n\}$, we have

$$\begin{aligned} & \begin{cases} [\alpha_i a_i + g_i, (1 - \alpha_j)a_j - g_j] = 0 \\ [\alpha_j a_j + g_j, (1 - \alpha_i)a_i - g_i] = 0; \end{cases} \Leftrightarrow \\ \Leftrightarrow & \begin{cases} \alpha_i(1 - \alpha_j)[a_i, a_j] + \alpha_i[g_j, a_i] + (1 - \alpha_j)[g_i, a_j] - [g_i, g_j] = 0 \\ -\alpha_j(1 - \alpha_i)[a_i, a_j] + \alpha_j[g_i, a_j] + (1 - \alpha_i)[g_j, a_i] + [g_i, g_j] = 0. \end{cases} \quad (2) \end{aligned}$$

For any i there exists j such that $a_i \leftrightarrow a_j$. Otherwise, there would be no edge incident to a_i in $\overline{G}$ and $\{a_i\}$ would be a connected component of $\overline{G}$ that would contradict to the condition of the lemma.

Since $L(A; G)$ is homogeneous we obtain

$$\begin{cases} \alpha_i(1 - \alpha_j)[a_i, a_j] + \alpha_i[\omega_1(g_j), a_i] + (1 - \alpha_j)[\omega_1(g_i), a_j] - [\omega_1(g_i), \omega_1(g_j)] = 0 \\ -\alpha_j(1 - \alpha_i)[a_i, a_j] + \alpha_j[\omega_1(g_i), a_j] + (1 - \alpha_i)[\omega_1(g_j), a_i] + [\omega_1(g_i), \omega_1(g_j)] = 0. \end{cases} \quad (3)$$

for any $i, j \in \{1, 2, \dots, n\}$.

Next, $[a_i, a_j] \neq 0$ because $a_i \leftrightarrow a_j$. By Lemma 5 and by choosing g_i and g_j the products $[\omega_1(g_j), a_i]$, $[\omega_1(g_i), a_j]$, and $[\omega_1(g_i), \omega_1(g_j)]$ are linear combinations of monomials of the form $[a_p, a_q]$, where $\{p, q\} \neq \{i, j\}$. Consequently, α_i and α_j must satisfy the following system

$$\begin{cases} \alpha_i(1 - \alpha_j) = 0 \\ \alpha_j(1 - \alpha_i) = 0. \end{cases} \quad (4)$$

because conversely, $\alpha_i(1 - \alpha_j)[a_i, a_j]$ in the first equation of (3) or $\alpha_j(1 - \alpha_i)[a_i, a_j]$ in the second equation of the same system cannot cancel.

From the first equation of the system (4) either $\alpha_i = 0$ or $1 - \alpha_j = 0$. In the former case, $1 - \alpha_i = 1$ and by the second equation of the system $\alpha_j = 0$. So, we obtain $(g_i)_2 = a_i - g_i$ and $(g_j)_2 = a_j - g_j$. In the latter case, $1 - \alpha_j = 0$. Then $\alpha_j = 1$, therefore by the second equation of (4) $1 - \alpha_i = 0$ and we get $(g_i)_1 = a_i + g_i$ and $(g_j)_1 = a_j + g_j$.

Let $a_i \leftrightarrow a_j$. Since $\overline{G}$ is connected there is a sequence of vertices

$$a_i = a_{i_0}, a_{i_1}, \dots, a_{i_k} = a_j \quad (5)$$

such that a_{i_p} is adjacent to $a_{i_{p+1}}$ in the graph $\overline{G}$ (i.e. a_{i_p} is not connected to $a_{i_{p+1}}$ in the graph G) for $p = 0, 1, \dots, k - 1$.

We need to show that if $(a_i)_1 = a_i + g_i$ then $(a_j)_1 = a_j + g_j$ and if $(a_i)_2 = a_i - g_i$ then $(a_j)_2 = a_j - g_j$. Let us prove this statement by induction on k .

The basis ($k = 1$) has already been proved above.

Suppose that the statement holds for any sequence $a_{t_0}, s_{t_1}, \dots, a_{t_{k-1}}$ such that $a_{t_p} \leftrightarrow a_{t_{p+1}}$ for $p = 0, 1, \dots, k-2$. Consider the sequence (5). If $(a_i)_1 = a_i + g_i$ then $(a_{i_{k-1}})_1 = a_{i_{k-1}} + g_{i_{k-1}}$ by the inductive hypothesis. Therefore, $(a_{i_k})_1 = (a_j)_1 = a_j + g_j$ as above. The case $(a_i)_2 = a_i - g_i$ is considered analogously. $\square$

Lemma 6 implies that without loss of generality we can assume that a_i corresponds to the pair $(a_i + g_i, -g_i)$, where $a_i \notin \text{supp}(\omega_1(g_i))$, and a_j corresponds to the pair $(a_j + g_j, -g_j)$, where $a_j \notin \text{supp}(\omega_1(g_j))$. So, for any $a_i, a_j \in A$ system of equations (3) can be rewritten as

$$\begin{cases} [g_j, a_i] - [g_i, g_j] = 0 \\ [g_i, a_j] + [g_i, g_j] = 0. \end{cases} \quad (6)$$

Lemma 7. *Let G be a graph with the set of vertices $A = \{a_1, a_2, \dots, a_n\}$, where $n \geq 2$. If $\bar{G}$ is connected, $L(A; G) = L_1 \oplus L_2$ and g_i is defined as above then $\omega_1(g_i) = 0$.*

Proof. Let $a_i = (a_i + g_i, -g_i)$. It follows from (1) that $[g_i, a_i] = 0$, consequently $[\omega_1(g_i), a_i] = 0$. So, by Theorem 3 we get $a_i \leftrightarrow \text{supp}(\omega_1(g_i))$.

Next, we can rewrite (3) as follows

$$\begin{cases} [\omega_1(g_j), a_i] - [\omega_1(g_i), \omega_1(g_j)] = 0 \\ [\omega_1(g_i), a_j] + [\omega_1(g_i), \omega_1(g_j)] = 0. \end{cases} \quad (7)$$

Suppose that $\omega_1(g_i) \neq 0$. Let us show that in this case there is $j \neq i$ such that a_j is adjacent to a vertex in $\text{supp}(\omega_1(g_i))$. Take $a_r \in \text{supp}(\omega_1(g_i))$. Since $\bar{G}$ is connected, there is a sequence of vertices $a_r = a_{i_0}, a_{i_1}, \dots, a_{i_k} = a_i$ such that for any $p \in \{0, 1, \dots, k-1\}$ the vertices a_{i_p} and $a_{i_{p+1}}$ are adjacent in $\bar{G}$ or, equivalently, $a_{i_p} \leftrightarrow a_{i_{p+1}}$. Let s be the smallest number such that $a_{i_s} \notin \text{supp}(\omega_1(g_i))$. Such number exists since $a_i = a_{i_k} \leftrightarrow \text{supp}(\omega_1(g_i))$ and therefore $a_i \notin \text{supp}(\omega_1(g_i))$. On the other hand, $i_s \neq i$ because $a_{i_s} \leftrightarrow a_{i_{s-1}}$ while $a_i \leftrightarrow a_{i_{s-1}}$. We may assume that $j = i_s$.

Consider the second equation of (7). Note that $[\omega_1(g_i), \omega_1(g_j)]$ is a linear combination of elements of the form $[a_q, a_t]$ and since $a_j \notin \text{supp}(\omega_1(g_i)) \cup \text{supp}(\omega_1(g_j))$, neither a_q nor a_t can be equal to a_j . Consequently, the element $\beta[a_{i_{s-1}}, a_j]$ for $\beta \neq 0$ is among the summands in $[\omega_1(g_i), a_j]$ but no monomial of the form $\gamma[a_{i_{s-1}}, a_j]$ or $\gamma[a_j, a_{i_{s-1}}]$ for $\gamma \neq 0$ exists among the summands in $[\omega_1(g_i), \omega_1(g_j)]$. Since $[a_{i_{s-1}}, a_j] \neq 0$, $[\omega_1(g_i), a_j]$ cannot be equal to 0. Therefore $[\omega_1(g_i), a_j] + [\omega_1(g_i), \omega_1(g_j)] \neq 0$. We get a contradiction to (7).

So, if $L(A; G) = \mathcal{L}(A; G)$ then we are done. If $L(A; G) = \mathcal{M}(A; G)$ or $L(A; G) = \mathcal{N}_m(A; G)$ for $m \geq 2$, then the statement follows from the fact that the restrictions of the natural homomorphisms $\mathcal{L}(A; G) \rightarrow \mathcal{N}_m(A; G)$ and $\mathcal{L}(A; G) \rightarrow \mathcal{M}(A; G)$ to the maps $\mathcal{L}_m(A; G) \rightarrow \mathcal{N}_m(A; G)$ and $\mathcal{L}_3(A; G) \rightarrow \mathcal{M}(A; G)$ are bijections. $\square$

Lemma 8. *Let G be a graph with the set of vertices $A = \{a_1, a_2, \dots, a_n\}$, where $n \geq 2$. If $\overline{G}$ is connected, $\mathcal{L}(A; G) = L_1 \oplus L_2$, and g_i be defined as above then $g_i = 0$ for all $i \in \{1, 2, \dots, n\}$.*

Proof. As we have shown in Lemma 7, $g_i = o_1(g_i)$ for $i = 1, 2, \dots, n$. So, (6) can be rewritten as

$$\begin{cases} [o_1(g_j), a_i] - [o_1(g_i), o_1(g_j)] = 0 \\ [o_1(g_i), a_j] + [o_1(g_i), o_1(g_j)] = 0 \end{cases} \quad (8)$$

Suppose that $o_1(g_i) \neq 0$. Then there exists $j \neq i$ such that $a_j \notin \text{supp}(o_1(g_i))$ but it is adjacent to one of the vertices in $\text{supp}(o_1(g_i))$. The proof is similar to that in Lemma 7.

Summing the equations in (8) we get

$$[o_1(g_j), a_i] + [o_1(g_i), a_j] = 0. \quad (9)$$

By Theorem 3 implies

$$[o_1(g_i), a_j] \neq 0. \quad (10)$$

On the other hand, by (1) $a_j \notin \text{supp}([o_1(g_j)])$ and $a_i \notin \text{supp}([o_1(g_i)])$. Thus the multi-degrees of the summands of $[o_1(g_j), a_i]$ are not equal to the multi-degrees of the summands of $[o_1(g_i), a_j]$ we obtain a contradiction to (9). $\square$

Lemma 9. *Let G be a graph with the set of vertices $A = \{a_1, a_2, \dots, a_n\}$, where $n \geq 2$. If $\overline{G}$ is connected, $\mathcal{N}_m(A; G) = L_1 \oplus L_2$, where $m \geq 2$, and g_i be defined as above then $O_{m-1}(g_i) = 0$ for all $i \in \{1, 2, \dots, n\}$.*

Proof. To prove this statement by induction it suffices to show that if $O_k(g_i) = 0$ for all $i \in \{1, 2, \dots, n\}$ for some k such that $1 \leq k \leq m - 2$, then $\omega_{k+1}(g_i) = 0$ for all $i \in \{1, 2, \dots, n\}$.

Suppose that $O_k(g_i) = 0$ and $\omega_{k+1}(g_i) \neq 0$ for some $i \in \{1, 2, \dots, n\}$ and for some $k \in \{1, 2, \dots, m - 1\}$. Since $\mathcal{N}_m(A; E)$ homogeneous the second equation in (6) implies

$$[\omega_{k+1}(g_i), a_j] = 0. \quad (11)$$

Indeed, since $O_k(g_i) = O_k(g_j) = 0$, the product $[g_i, g_j] = [O_{k+1}(g_i), O_{k+1}(g_j)]$ is represented as a linear combination of monomials of degrees at least $(k + 1) + (k + 1) = 2k + 2 > k + 2$.

On the other hand, there exists $j \neq i$ such that $a_j \in \text{supp}(\omega_{k+1}(g_i))$. The proof is similar to that in Lemma 7. So, by Theorem 3 we get

$$[\omega_{k+1}(g_i), a_j] \neq 0 \quad (12)$$

that contradicts to (11). Therefore, $\omega_{k+1}(g_i) = 0$. So, $O_{m-1}(g_i) = 0$. $\square$

Lemma 10. *Let G be a graph with the set of vertices $A = \{a_1, a_2, \dots, a_n\}$, where $n \geq 2$. If $\mathcal{M}(A; G) = L_1 \oplus L_2$, and g_i be defined as above. Then $g_i = 0$ for all $i \in \{1, 2, \dots, n\}$.*

Proof. As we have shown in Lemma 7, $g_i = o_1(g_i)$ for $i = 1, 2, \dots, n$. So, as in Lemma 8, the system of equations (8) holds. Since the considered algebra is metabelian the second equation of (8) gives

$$[o_1(g_i), a_j] = 0. \quad (13)$$

On the other hand, suppose that $o_1(g_i) \neq 0$. Then $|\text{supp}(o_1(g_i))| \geq 2$. So there exists $a_j \in \text{supp}(o_1(g_i)) \setminus \{a_i\}$. Let us order the set A in such a way that a_j is the smallest element. There is the representation

$$o_1(g_i) = \sum_{\bar{\delta}} g_{i,\bar{\delta}}, \quad (14)$$

where $g_{i,\bar{\delta}} \in \mathcal{M}(A; G)$ is a multi-homogeneous element of multi-degree $\bar{\delta}$.

Let $g_{i,\bar{\delta}_0}$ be a summand in the right-hand side of (14) such that $a_j \in \text{supp}(g_{i,\bar{\delta}_0})$. Then we can write $g_{i,\bar{\delta}_0} = \sum_{p=1}^k \lambda_p [u_p]$, where $[u_p]$ are basis monomials with respect to the order described above and $\lambda_p \in R \setminus \{0\}$. Let $H_0, H_1, \dots, H_s$ ($s \geq 1$) be connected components of the graph $G(\text{supp}(g_{i,\bar{\delta}_0}))$. It can be assumed that $a_j \in H_0$. According to Theorem 4 each $[u_p]$ has the form $[u_p] = [a_{i_{p,1}}, a_j, a_{i_{p,3}}, \dots, a_{p,t}]$, where $a_{i_{p,1}} > a_j$, $a_j \leq a_{i_{p,3}} \leq \dots \leq a_{p,t}$, and $a_{i_{1,1}}, a_{i_{2,1}}, \dots, a_{i_{k,1}}$ are the largest elements of some of the sets $A(H_1), A(H_2), \dots, A(H_s)$. We have

$$[o_1(g_i), a_j] = \sum_{\bar{\delta}} [g_{i,\bar{\delta}}, a_j],$$

where

$$\begin{aligned} [g_{i,\bar{\delta}_0}, a_j] &= \sum_{p=1}^k \lambda_p [[u_p], a_j] = \sum_{p=1}^k \lambda_p [[a_{i_{p,1}}, a_j, a_{i_{p,3}}, \dots, a_{i_{p,t}}], a_j] = \\ &= \sum_{p=1}^k \lambda_p [a_{i_{p,1}}, a_j, a_j, a_{i_{p,3}}, \dots, a_{i_{p,t}}]. \end{aligned} \quad (15)$$

Then the graph $G(\text{supp}([u_p], a_j))$ coincides with the graphs $G(\text{supp}(g_{i,\bar{\delta}_0}))$ because $\text{supp}(g_{i,\bar{\delta}_0}) = \text{supp}([u_p]) = \text{supp}([u_p], a_j)$. In particular, these graphs have the same connected components. Therefore all monomials in (15) are pairwise distinct basis elements with respect to the order described above. So, $[g_{i,\bar{\delta}_0}, a_j] \neq 0$. Since $\mathcal{M}(G; H)$ is homogeneous we obtain $[o_1(g_i), a_j] \neq 0$ that contradicts to (13). Therefore, $o_1(g_i) = g_i = 0$. $\square$

Now we are ready to proof the following theorem.

Theorem 11. *Let G be a graph with the set of vertices $A = \{a_1, a_2, \dots, a_n\}$, where $n \geq 2$ and let $L(A; G)$ be a partially commutative Lie algebra $\mathcal{L}(A; G)$, or a partially commutative metabelian Lie algebra $\mathcal{M}(A; G)$, or a partially commutative nilpotent of degree $m \geq 2$ Lie algebra $\mathcal{N}_m(A; G)$. Then $L(A; G)$ splits into a direct sum of two non-zero subalgebras if and only if $\bar{G}$ is not connected.*

Proof. Suppose that $\overline{G}$ is connected and $L(A; G) = L_1 \oplus L_2$. Without loss of generality we can assume that a_i corresponds to $(a_i + g_i, -g_i)$, where $a_i \notin \text{supp}(\omega_1(g_i))$.

Let $L(A; G)$ is either $\mathcal{L}(A; G)$ or $\mathcal{M}(A; G)$. Then $g_i = 0$ by Lemma 8 or Lemma 10. Hence $a_i \in L_1$ for $i = 1, 2, \dots, n$. Thus, $L_1 = \mathcal{L}(A; G)$ and so $L_2 = 0$.

Let $L(A; G) = \mathcal{N}_m(A; G)$. Since a_i corresponds to $(a_i + g_i, -g_i)$, we obtain $g_i \in L_2$.

On the other hand, Lemma 9 implies that $O_{m-1}(g_i) = 0$. Consequently, $[(a_p)_1, (a_q)_1] = [a_p + g_p, a_q + g_q] = [a_p + \omega_m(g_p), a_q + \omega_m(g_q)] = [a_p, a_q]$ for any $p, q \in \{1, 2, \dots, n\}$. Now, let $[u(a_1, a_2, \dots, a_n)]$ be a non-commutative non-associative monomial of cumulative degree at least 3. Then

$$[u(a_1, a_2, \dots, a_n)] = [[u_1(a_1, a_2, \dots, a_n)], [u_2(a_1, a_2, \dots, a_n)]].$$

If the cumulative degrees of $[u_1(a_1, a_2, \dots, a_n)]$ and $[u_2(a_1, a_2, \dots, a_n)]$ are greater than 1, then by the inductive hypothesis $[u_l((a_1)_1, (a_2)_1, \dots, (a_n)_1)] = [u_l(a_1, a_2, \dots, a_n)]$ for $l = 1, 2$. Therefore,

$$\begin{aligned} [u((a_1)_1, (a_2)_1, \dots, (a_n)_1)] &= \\ &= [[u_1((a_1)_1, (a_2)_1, \dots, (a_n)_1)], [u_2((a_1)_1, (a_2)_1, \dots, (a_n)_1)]] = \\ &= [[u_1(a_1, a_2, \dots, a_n)], [u_2(a_1, a_2, \dots, a_n)]] = \\ &= [u(a_1, a_2, \dots, a_n)] \end{aligned}$$

in $\mathcal{N}_m(A; G)$. If $[u_2(a_1, a_2, \dots, a_n)] = a_q$ for some $q \in \{1, 2, \dots, n\}$ then

$$\begin{aligned} [u((a_1)_1, (a_2)_1, \dots, (a_n)_1)] &= [[u_1((a_1)_1, (a_2)_1, \dots, (a_n)_1)], (a_p)_1] = \\ &= [[u_1((a_1)_1, (a_2)_1, \dots, (a_n)_1)], a_p + g_p] = \\ &= [[u_1(a_1, a_2, \dots, a_n)], a_p + g_p] = \\ &= [u_1(a_1, a_2, \dots, a_n), a_p] = \\ &= [u(a_1, a_2, \dots, a_n)] \end{aligned}$$

in $\mathcal{N}_m(A; G)$.

The case $[u_1] = a_p$ for some $p \in \{1, 2, \dots, n\}$ is analogous. Therefore, for any non-commutative non-associative polynomial $f(a_1, \dots, a_n)$ such that $\omega_1(f(a_1, a_2, \dots, a_n)) = 0$ the equation

$$f((a_1)_1, (a_2)_1, \dots, (a_n)_1) = f(a_1, a_2, \dots, a_n)$$

holds. In particular it holds for any g_i . Consequently, $g_i \in L_1$ and so $g_i = 0$.

The converse is obvious. If $\overline{G}$ is not connected then $A = A_1 \sqcup A_2$, where there are no edges $\{a, b\}$ such that $a \in A_1$ and $b \in A_2$. Then $A_1 \leftrightarrow A_2$ and $L(A; G) = L(A_1; G(A_1)) \oplus L(A_2; G(A_2))$. $\square$

References

- [1] Cartier F., Foata D., *Problèmes combinatoires de computation et rearrangements*, Lecture Notes in Mathematics, 85, Springer-Verlag, Berlin, 1969.

- [2] Servatius H., *Automorphisms of graph groups*, J. Algebra, **126**, 1 (1989), 34–60.
- [3] Duchamp G., Krob D., *The lower central series of the free partially commutative group*, Semigroup Forum, **45** (1992), 385–394.
- [4] Duncan A. J., Kazachkov I. V., Remeslennikov V. N., *Parabolic and quasiparabolic subgroups of free partially commutative groups*, Journal of Algebra, **318**, 2 (2007), 918–932.
- [5] Charney R., *An introduction to right-angled Artin groups*, Geometriae Dedicata **125** (2007), 141–158.
- [6] Poroshenko E. N., Timoshenko E. I., *Partially commutative groups and Lie algebras*, Siberian Electronic Mathematical Repots, **18**, 1 (2021), 668–693.
- [7] Timoshenko E. I., *On the decomposition of partially commutative groups of varieties and their defining graphs*, Siberian Mathematical Journal, **62**, 4 (2021), 742–746.
- [8] Poroshenko E. N., *Centralizers in partially commutative Lie algebras*, Algebra and Logic, **51**, 4 (2012), 351–371.
- [9] Poroshenko E. N., Timoshenko E. I., *Universal Equivalence of Partially Commutative Metabelian Lie Algebras*, J. Algebra, (2013), 143–168.

EVGENY NIKOLAEVICH POROSHENKO
NOVOSIBIRSK STATE TECHNICAL UNIVERSITY,
PR. K. MARXA, 20,
630073, NOVOSIBIRSK, RUSSIA
E-mail address: auto_stoper@ngs.ru