

**ALGEBRAS OF BINARY FORMULAS FOR
 \aleph_0 -CATEGORICAL WEAKLY CIRCULARLY MINIMAL
THEORIES: PIECEWISE MONOTONIC CASE****B.SH. KULPESHOV** *Communicated by S.V. SUDOPLATOV*

Abstract: Algebras of binary isolating formulas are described for \aleph_0 -categorical 1-transitive non-primitive weakly circularly minimal theories of convexity rank greater than 1 having a non-trivial piecewise (non-strictly) monotonic function.

Keywords: weak circular minimality, algebra of binary formulas, \aleph_0 -categorical theory, circularly ordered structure, convexity rank.

1 Preliminaries

Algebras of binary formulas are a tool for describing relationships between elements of the sets of realizations of a type at the binary level with respect to the superposition of binary definable sets. A *binary isolating formula* is a formula of the form $\varphi(x, y)$ such that for some parameter a the formula $\varphi(a, y)$ isolates a complete type in $S(\{a\})$. The concepts and notations related to these algebras can be found in the papers [1, 2]. In recent years, algebras of binary formulas have been studied intensively and have been continued in the works [3]–[7].

KULPESHOV, B.SH., ALGEBRAS OF BINARY FORMULAS FOR \aleph_0 -CATEGORICAL WEAKLY CIRCULARLY MINIMAL THEORIES: PIECEWISE MONOTONIC CASE.

© 2023 KULPESHOV B.SH.

The work was supported by Science Committee of Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. BR20281002).

Received July, 17, 2023, Published October, 5, 2023.

Let L be a countable first-order language. Throughout we consider L -structures and assume that L contains a ternary relational symbol K , interpreted as a circular order in these structures (unless otherwise stated).

Let $\mathcal{M} = \langle M, \leq \rangle$ be a linearly ordered set. If we connect two endpoints of \mathcal{M} (possibly, $-\infty$ and $+\infty$), then we obtain a circular order. More formally, the *circular order* is described by a ternary relation K satisfying the following conditions:

- (co1) $\forall x \forall y \forall z (K(x, y, z) \rightarrow K(y, z, x))$;
- (co2) $\forall x \forall y \forall z (K(x, y, z) \wedge K(y, x, z) \Leftrightarrow x = y \vee y = z \vee z = x)$;
- (co3) $\forall x \forall y \forall z (K(x, y, z) \rightarrow \forall t [K(x, y, t) \vee K(t, y, z)])$;
- (co4) $\forall x \forall y \forall z (K(x, y, z) \vee K(y, x, z))$.

Sometimes we will identify \mathcal{M} and the universe M if a linear/circular order is fixed.

The notion of *weak circular minimality* was studied initially in [8]. Let $A \subseteq M$, where \mathcal{M} is a circularly ordered structure. The set A is called *convex* if for any $a, b \in A$ the following property is satisfied: for any $c \in M$ with $K(a, c, b)$, $c \in A$ holds, or for any $c \in M$ with $K(b, c, a)$, $c \in A$ holds. A *weakly circularly minimal structure* is a circularly ordered structure $\mathcal{M} = \langle M, K, \dots \rangle$ such that any definable (with parameters) subset of M is a union of finitely many convex sets in \mathcal{M} . Recall [9] that such a structure \mathcal{M} is called *circularly minimal* if any definable (with parameters) of M is a union of finitely many intervals and points in \mathcal{M} . Clearly, the weak circular minimality is a generalization of circular minimality. Notice also that any weakly o-minimal structure is weakly circular minimal. The converse, in general, fails. The study of weakly circularly minimal structures was continued in the papers [10]–[16].

Let \mathcal{M} be an \aleph_0 -categorical weakly circularly minimal structure, $G := \text{Aut}(\mathcal{M})$. Following the standard group theory terminology, the group G is called *k-transitive* if for any pairwise distinct $a_1, a_2, \dots, a_k \in M$ and pairwise distinct $b_1, b_2, \dots, b_k \in M$ there exists $g \in G$ such that $g(a_1) = b_1, g(a_2) = b_2, \dots, g(a_k) = b_k$. A *congruence* on \mathcal{M} is an arbitrary G -invariant equivalence relation on \mathcal{M} . The group G is called *primitive* if G is 1-transitive and there are no non-trivial proper congruences on \mathcal{M} .

Let \mathcal{M}, \mathcal{N} be circularly ordered structures. The *2-reduct* of \mathcal{M} is a circularly ordered structure with the same universe of \mathcal{M} and consisting of predicates for each \emptyset -definable relation on \mathcal{M} of arity ≤ 2 as well as of the ternary predicate K for the circular order, but does not have other predicates of arities more than two. We say that the structure \mathcal{M} is *isomorphic* to \mathcal{N} *up to binarity* or *binarily isomorphic* to \mathcal{N} if a 2-reduct of \mathcal{M} is isomorphic to a 2-reduct of \mathcal{N} .

Notation.

- (1) $K_0(x, y, z) := K(x, y, z) \wedge y \neq x \wedge y \neq z \wedge x \neq z$.

(2) $K(u_1, \dots, u_n)$ denotes a formula saying that all subtuples of the tuple $\langle u_1, \dots, u_n \rangle$ having the length 3 (in ascending order) satisfy K ; similar notations are used for K_0 .

(3) Let A, B, C be disjoint convex subsets of a circularly ordered structure \mathcal{M} . We write $K(A, B, C)$ if for any $a, b, c \in M$ with $a \in A, b \in B, c \in C$ we have $K(a, b, c)$. We extend naturally that notation using, for instance, the notation $K_0(A, d, B, C)$ if $d \notin A \cup B \cup C$ and $K_0(A, d, B) \wedge K_0(d, B, C)$ holds.

The following definition can be used in a circular ordered structure as well.

Definition 1. [17], [18] Let T be a weakly o-minimal theory, M be a sufficiently saturated model of T , $A \subseteq M$. The rank of convexity of the set A ($RC(A)$) is defined as follows:

- 1) $RC(A) = -1$ if $A = \emptyset$.
- 2) $RC(A) = 0$ if A is finite and non-empty.
- 3) $RC(A) \geq 1$ if A is infinite.
- 4) $RC(A) \geq \alpha + 1$ if there exist a parametrically definable equivalence relation $E(x, y)$ and an infinite sequence of elements $b_i \in A, i \in \omega$, such that:
 - For every $i, j \in \omega$ whenever $i \neq j$ we have $M \models \neg E(b_i, b_j)$;
 - For every $i \in \omega, RC(E(x, b_i)) \geq \alpha$ and $E(M, b_i)$ is a convex subset of A .
- 5) $RC(A) \geq \delta$ if $RC(A) \geq \alpha$ for all $\alpha < \delta$, where δ is a limit ordinal.

If $RC(A) = \alpha$ for some α , we say that $RC(A)$ is defined. Otherwise (i.e. if $RC(A) \geq \alpha$ for all α), we put $RC(A) = \infty$.

The rank of convexity of a formula $\phi(x, \bar{a})$, where $\bar{a} \in M$, is defined as the rank of convexity of the set $\phi(M, \bar{a})$, i.e. $RC(\phi(x, \bar{a})) := RC(\phi(M, \bar{a}))$.

The rank of convexity of an 1-type p is defined as the rank of convexity of the set $p(M)$, i.e. $RC(p) := RC(p(M))$.

In particular, a theory has convexity rank 1 if there are no definable (with parameters) equivalence relations with infinitely many infinite convex classes.

Let $f : I \rightarrow M$ be an \emptyset -definable function with $Dom(f) = I \subseteq M$, where I is an open convex set. We say that f is *monotonic-to-right (left) on I* if it preserves (reverses) the relation K_0 , i.e. for any $a, b, c \in I$ such that $K_0(a, b, c)$ we have $K_0(f(a), f(b), f(c))$ ($K_0(f(c), f(b), f(a))$). We also say that f is *piecewise monotonic-to-right (left) on M* if there exists an \emptyset -definable non-trivial equivalence relation $E(x, y)$ partitioning M into finitely many infinite convex classes so that f is monotonic-to-right on each E -class and f is not monotonic-to-left (right) on M/E , where by M/E we denote the set of representatives of E -classes in M .

Example. [10] Let $M := \langle M, K, E^2, f^1 \rangle$ be a circularly ordered structure, where M is a disjoint union of $\mathbb{Q}_1, \mathbb{Q}_2, \dots, \mathbb{Q}_6$, where \mathbb{Q}_i is a copy of the ordering of rational numbers \mathbb{Q} . The symbol E interprets an equivalence relation on M as follows: $E(a, b)$ iff there is $1 \leq i \leq 6$ with $a, b \in \mathbb{Q}_i$.

The symbol f interprets a function on M as follows: $f(Q_i) = Q_{i+3}$ for each $1 \leq i \leq 3$, $f(Q_j) = Q_{j-3}$ for each $4 \leq j \leq 6$, and $f(q) = -q$ for all $q \in Q$.

It can be proved that M is an \aleph_0 -categorical 1-transitive weakly circularly minimal structure, f is a bijection on M so that $f^2(a) = a$ for all $a \in M$, f is monotonic-to-left on each E -class and f is monotonic-to-right on M/E , i.e. f is piecewise monotonic-to-left on M .

Lemma 1. [10] *Let M be an \aleph_0 -categorical 1-transitive weakly circularly minimal structure, f be an \emptyset -definable function on M . Then f cannot be piecewise monotonic-to-right on M .*

The following theorem characterizes \aleph_0 -categorical 1-transitive non-primitive weakly circularly minimal structures of convexity rank greater than 1 having a non-trivial piecewise (non-strictly) monotonic function up to binarity:

Theorem 1. [11] *(piecewise monotonic case) Let M be an \aleph_0 -categorical 1-transitive non-primitive weakly circularly minimal structure of convexity rank greater than 1 having a non-trivial piecewise (non-strictly) monotonic function so that $dcl(a) \neq \{a\}$ for some $a \in M$. Then M is isomorphic up to binarity to $M_{s,m,k} := \langle M, K, f^1, E_1^2, \dots, E_s^2, E_{s+1}^2 \rangle$, where*

- M is a circularly ordered structure, M is densely ordered, $s \geq 1$, $k \geq 2$, k is even, k divides m , $m \geq 4$;
- E_{s+1} is an equivalence relation partitioning M into m infinite convex classes without endpoints, for every $1 \leq i \leq s$ the relation E_i is an equivalence relation partitioning every E_{i+1} -class into infinitely many infinite convex E_i -subclasses without endpoints so that the induced order on E_i -subclasses is dense without endpoints;
- f is a bijection on M so that $f^k(a) = a$ for any $a \in M$, for every $1 \leq i \leq s+1$ $f(E_i(M, a)) = E_i(M, f(a))$ and $\neg E_i(a, f(a))$, and f is piecewise monotonic-to-left on M , i.e. f is monotonic-to-left on each E_{s+1} -class and f is monotonic-to-right on M/E_{s+1} .

In [19] algebras of binary isolating formulas are described for \aleph_0 -categorical weakly circularly minimal theories with a primitive automorphism group. In [20] algebras of binary isolating formulas are described for \aleph_0 -categorical weakly circularly minimal theories of convexity rank 1 with a 1-transitive non-primitive automorphism group and a non-trivial definable closure. Here we describe algebras of binary isolating formulas for \aleph_0 -categorical weakly circularly minimal theories of convexity rank greater than 1 with a 1-transitive non-primitive automorphism group and having a non-trivial piecewise (non-strictly) monotonic function.

2 Results

Definition 2. [1] Let $p \in S_1(\emptyset)$ be non-algebraic. The algebra $\mathcal{P}_{\nu(p)}$ is said to be *deterministic* if $u_1 \cdot u_2$ is a singleton for any labels $u_1, u_2 \in \rho_{\nu(p)}$.

Generalizing the last definition, we say that the algebra $\mathcal{P}_{\nu(p)}$ is *m-deterministic* if the product $u_1 \cdot u_2$ consists of at most m elements for any labels $u_1, u_2 \in \rho_{\nu(p)}$. We also say that an m -deterministic algebra $\mathcal{P}_{\nu(p)}$ is *strictly m-deterministic* if it is not $(m - 1)$ -deterministic.

Example. Consider the structure $M_{1,4,2} := \langle M, K^3, f^1, E_1^2, E_2^2 \rangle$ from Theorem 1, where f is piecewise monotonic-to-left on M , E_1 is an equivalence relation partitioning M into infinitely many infinite convex classes, E_2 is an equivalence relation partitioning M into four infinite convex classes.

We assert that $Th(M_{1,4,2})$ has twelve binary isolating formulas:

$$\begin{aligned} \theta_0(x, y) &:= x = y, \theta_1(x, y) := K_0(x, y, f(x)) \wedge E_1(x, y), \\ \theta_2(x, y) &:= K_0(x, y, f(x)) \wedge \neg E_1(x, y) \wedge E_2(x, y), \\ \theta_3(x, y) &:= K_0(x, y, f(x)) \wedge \neg E_2(x, y) \wedge \neg E_2(f(x), y), \\ \theta_4(x, y) &:= K_0(x, y, f(x)) \wedge \neg E_1(f(x), y) \wedge E_2(f(x), y), \\ \theta_5(x, y) &:= K_0(x, y, f(x)) \wedge E_1(f(x), y), \\ \theta_6(x, y) &:= f(x) = y, \theta_7(x, y) := K_0(f(x), y, x) \wedge E_1(f(x), y), \\ \theta_8(x, y) &:= K_0(f(x), y, x) \wedge \neg E_1(f(x), y) \wedge E_2(f(x), y), \\ \theta_9(x, y) &:= K_0(f(x), y, x) \wedge \neg E_2(f(x), y) \wedge \neg E_2(x, y), \\ \theta_{10}(x, y) &:= K_0(f(x), y, x) \wedge \neg E_1(x, y) \wedge E_2(x, y), \\ \theta_{11}(x, y) &:= K_0(f(x), y, x) \wedge E_1(x, y), \end{aligned}$$

and

$$K_0(\theta_i(a, M), \theta_{i+1}(a, M), \theta_{i+2}(a, M)), \text{ where } 0 \leq i \leq 9,$$

$$K_0(\theta_{10}(a, M), \theta_{11}(a, M), \theta_0(a, M)), \quad K_0(\theta_{11}(a, M), \theta_0(a, M), \theta_1(a, M))$$

hold for any $a \in M$.

Define labels for these formulas as follows:

$$\text{label } k \text{ for } \theta_k(x, y) \text{ where } 0 \leq k \leq 11.$$

It easy to check that for the algebra $\mathfrak{P}_{M_{1,4,2}}$ the Cayley table has the following form:

·	0	1	2	3	4	5	6	7	8	9	10	11
0	{0}	{1}	{2}	{3}	{4}	{5}	{6}	{7}	{8}	{9}	{10}	{11}
1	{1}	{1}	{2}	{3}	{4}	{5}	{5}	{5, 6, 7}	{8}	{9}	{10}	{11, 0, 1}
2	{2}	{2}	{2}	{3}	{4}	{4}	{4}	{4}	...	{9}	{10, 11, 0, 1, 2}	{2}
3	{3}	{3}	{3}	...	{9}	{9}	{9}	{9}	{9}	...	{3}	{3}
4	{4}	{4}	...	{9}	...	{2}	{2}	{2}	{2}	{3}	{4}	{4}
5	{5}	{5, 6, 7}	{8}	{9}	{10}	{11, 0, 1}	{1}	{1}	{2}	{3}	{4}	{5}
6	{6}	{7}	{8}	{9}	{10}	{11}	{0}	{1}	{2}	{3}	{4}	{5}
7	{7}	{7}	{8}	{9}	{10}	{11}	{11}	{11, 0, 1}	{2}	{3}	{4}	{5, 6, 7}
8	{8}	{8}	{8}	{9}	{10}	{10}	{10}	{10}	...	{3}	{4, 5, 6, 7, 8}	{8}
9	{9}	{9}	{9}	...	{3}	{3}	{3}	{3}	...	{3}	{9}	{9}
10	{10}	{10}	...	{3}	...	{8}	{8}	{8}	{9}	{10}	{10}	{10}
11	{11}	{11, 0, 1}	{2}	{3}	{4}	{5, 6, 7}	{7}	{7}	{8}	{9}	{10}	{11}

By the Cayley table the algebra $\mathfrak{P}_{M_{1,4,2}}$ is not commutative.

Theorem 2. *The algebra $\mathfrak{P}_{M_{s,m,k}}$ of binary isolating formulas having a piecewise monotonic-to-left function on M has $2k(s + 1) + m$ labels, is strictly $(2s + 3)$ -deterministic and is not commutative.*

Proof. Indeed, since $f^k(a) = a$, we have the following isolating formulas:

$$f^l(x) = y \text{ for every } 0 \leq l \leq k - 1.$$

Since for every $1 \leq i \leq s$ the relation E_i is an equivalence relation partitioning every E_{i+1} -class into infinitely many infinite convex E_i -subclasses without endpoints so that the induced order on E_i -subclasses is dense without endpoints, we obtain the following binary isolating formulas:

$$K_0(f^l(x), y, f^{l+1}(x)) \wedge E_1(f^l(x), y), \text{ where } 0 \leq l \leq k - 1,$$

$$K_0(f^l(x), y, f^{l+1}(x)) \wedge \neg E_j(f^l(x), y) \wedge E_{j+1}(f^l(x), y),$$

$$\text{where } 0 \leq l \leq k - 1, 1 \leq j \leq s - 1,$$

$$K_0(f^l(x), y, f^{l+1}(x)) \wedge \neg E_s(f^l(x), y) \wedge \neg E_s(f^{l+1}(x), y), \text{ where } 0 \leq l \leq k - 1,$$

$$K_0(f^l(x), y, f^{l+1}(x)) \wedge \neg E_j(f^{l+1}(x), y) \wedge E_{j+1}(f^{l+1}(x), y),$$

$$\text{where } 0 \leq l \leq k - 1, 1 \leq j \leq s - 1,$$

$$K_0(f^l(x), y, f^{l+1}(x)) \wedge E_1(f^{l+1}(x), y), \text{ where } 0 \leq l \leq k - 1.$$

Since in this structure there exists additionally an equivalence relation $E_{s+1}(x, y)$ partitioning M into m infinite convex classes, additionally the following binary isolating formulas appear:

$$K_0(f^l(x), y, f^{l+1}(x)) \wedge \neg E_s(f^l(x), y) \wedge E_{s+1}(f^l(x), y),$$

$$K_0(f^l(x), y, f^{l+1}(x)) \wedge \neg E_s(f^{l+1}(x), y) \wedge E_{s+1}(f^{l+1}(x), y),$$

where $0 \leq l \leq k - 1$.

Also, the formulas $\theta^{l,i}(x, y)$ containing the conjunctive term $K_0(f^l(x), y, f^{l+1}(x))$ and extracting the i -th E_{s+1} -class to the right of E_{s+1} -class containing $f^l(x)$ for some $1 \leq i \leq m/k - 1$ (here also $0 \leq l \leq k - 1$) will be binary isolating formulas. For example, the formula $\theta^{l,1}(x, y)$ has the following form:

$$\theta^{l,1}(x, y) := K_0(f^l(x), y, f^{l+1}(x)) \wedge \neg E_{s+1}(f^l(x), y) \wedge$$

$$\forall t [K_0(f^l(x), t, y) \wedge \neg E_{s+1}(t, y) \rightarrow E_{s+1}(f^l(x), t)].$$

Thus, we obtain $k + k + 2k(s - 1) + k + 2k + k(m/k - 1) = 2k(s + 1) + m$ binary isolating formulas.

The formulas

$$\exists t [f^l(x) = t \wedge K_0(f^l(t), y, f^{l+1}(t)) \wedge E_1(f^l(t), y)]$$

and

$$\exists t [K_0(f^l(x), t, f^{l+1}(x)) \wedge E_1(f^l(x), t) \wedge f^l(t) = y],$$

where $0 \leq l \leq k - 1$, uniquely determine the formula

$$K_0(f^{2l \bmod k}(x), y, f^{2l+1 \bmod k}(x)) \wedge E_1(f^{2l \bmod k}(x), y).$$

The formula

$$\exists t [K_0(x, t, f(x)) \wedge E_1(x, t) \wedge K_0(f(t), y, f^2(t)) \wedge E_1(f(t), y)]$$

is compatible with the following formulas:

$$\begin{aligned} &K_0(x, y, f(x)) \wedge E_1(f(x), y), \\ &f(x) = y, \\ &K_0(f(x), y, f^2(x)) \wedge E_1(f(x), y). \end{aligned}$$

While the formula

$$\exists t[K_0(f(x), t, f^2(x)) \wedge E_1(f(x), t) \wedge K_0(t, y, f(t)) \wedge E_1(t, y)]$$

uniquely determines the formula $K_0(f(x), y, f^2(x)) \wedge E_1(f(x), y)$. Consequently, the algebra $\mathfrak{A}_{M_{s,m,k}}$ is not commutative.

And in general we consider the formula

$$\begin{aligned} &\exists t[K_0(f^{l_1}(x), t, f^{l_1+1}(x)) \wedge E_1(f^{l_1}(x), t) \wedge \\ &K_0(f^{l_2}(t), y, f^{l_2+1}(t)) \wedge E_1(f^{l_2}(t), y)], \end{aligned}$$

where $0 \leq l_1, l_2 \leq k-1$. Such a formula for even l_2 uniquely determines the formula

$$K_0(f^{l_1+l_2 \pmod k}(x), y, f^{l_1+l_2+1 \pmod k}(x)) \wedge E_1(f^{l_1+l_2 \pmod k}(x), y),$$

and for odd l_2 it is compatible with the following three formulas:

$$\begin{aligned} &K_0(f^{l_1+l_2 \pmod k}(x), y, f^{l_1+l_2+1 \pmod k}(x)) \wedge E_1(f^{l_1+l_2 \pmod k}(x), y), \\ &f^{l_1+l_2 \pmod k}(x) = y, \end{aligned}$$

$$K_0(f^{l_1+l_2-1 \pmod k}(x), y, f^{l_1+l_2 \pmod k}(x)) \wedge E_1(f^{l_1+l_2 \pmod k}(x), y).$$

Further consider the formula

$$\begin{aligned} &\exists t[K_0(f^{l_1}(x), t, f^{l_1+1}(x)) \wedge E_1(f^{l_1}(x), t) \wedge K_0(f^{l_2}(t), y, f^{l_2+1}(t)) \\ &\wedge \neg E_j(f^{l_2}(t), y) \wedge E_{j+1}(f^{l_2}(t), y)], \end{aligned}$$

where $0 \leq l_1, l_2 \leq k-1$, $1 \leq j \leq s$. It uniquely determines the formula

$$\begin{aligned} &K_0(f^{l_1+l_2 \pmod k}(x), y, f^{l_1+l_2+1 \pmod k}(x)) \wedge \neg E_j(f^{l_1+l_2 \pmod k}(x), y) \\ &\wedge E_{j+1}(f^{l_1+l_2 \pmod k}(x), y). \end{aligned}$$

On the other hand, the formula

$$\begin{aligned} &\exists t[K_0(f^{l_2}(x), t, f^{l_2+1}(x)) \wedge \neg E_j(f^{l_2}(x), t) \wedge E_{j+1}(f^{l_2}(x), t) \\ &\wedge K_0(f^{l_1}(t), y, f^{l_1+1}(t)) \wedge E_1(f^{l_1}(t), y)] \end{aligned}$$

uniquely determines the formula

$$\begin{aligned} &K_0(f^{l_1+l_2-1 \pmod k}(x), y, f^{l_1+l_2 \pmod k}(x)) \wedge \neg E_j(f^{l_1+l_2 \pmod k}(x), y) \\ &\wedge E_{j+1}(f^{l_1+l_2 \pmod k}(x), y). \end{aligned}$$

Consider now the formulas $\theta^{l_1, i}(x, y)$ and $\theta^{l_2, j}(x, y)$ for arbitrary $0 \leq l_1, l_2 \leq k-1$, $1 \leq i, j \leq m/k-1$. If $i+j \pmod{m/k} \neq 0$, it is easy to check that the formulas

$$\exists t[\theta^{l_1, i}(x, t) \wedge \theta^{l_2, j}(t, y)] \text{ and } \exists t[\theta^{l_2, j}(x, t) \wedge \theta^{l_1, i}(t, y)]$$

uniquely determine the formula $\theta^{l_1+l_2(\bmod k), i+j(\bmod m/k)}(x, y)$.

If $i + j \pmod{m/k} = 0$, these formulas are compatible with the following $2s + 3$ formulas:

$$\begin{aligned} & f^{l_1+l_2+1(\bmod k)}(x) = y, \\ & K_0(f^{l_1+l_2(\bmod k)}(x), y, f^{l_1+l_2+1(\bmod k)}(x)) \wedge E_1(f^{l_1+l_2+1(\bmod k)}(x), y), \\ & K_0(f^{l_1+l_2(\bmod k)}(x), y, f^{l_1+l_2+1(\bmod k)}(x)) \wedge \neg E_j(f^{l_1+l_2+1(\bmod k)}(x), y) \\ & \quad \wedge E_{j+1}(f^{l_1+l_2+1(\bmod k)}(x), y), \quad 1 \leq j \leq s, \\ & K_0(f^{l_1+l_2+1(\bmod k)}(x), y, f^{l_1+l_2+2(\bmod k)}(x)) \wedge E_1(f^{l_1+l_2+1(\bmod k)}(x), y), \\ & K_0(f^{l_1+l_2+1(\bmod k)}(x), y, f^{l_1+l_2+2(\bmod k)}(x)) \wedge \neg E_j(f^{l_1+l_2+1(\bmod k)}(x), y) \\ & \quad \wedge E_{j+1}(f^{l_1+l_2+1(\bmod k)}(x), y), \quad 1 \leq j \leq s. \end{aligned}$$

Thus, the algebra $\mathfrak{B}_{M_s, m, k}$ is strictly $(2s + 3)$ -deterministic. \square

References

- [1] S.V. Sudoplatov, *Classification of countable models of complete theories*, Novosibirsk: NSTU, 2018.
- [2] I.V. Shulepov, S.V. Sudoplatov, *Algebras of distributions for isolating formulas of a complete theory*, Sib. Elektron. Math. Izv., **11** (2014), 380–407. Zbl 1355.03025
- [3] S.V. Sudoplatov, *Algebras of distributions for semi-isolating formulas of a complete theory*, Sib. Elektron. Mat. Izv., **11** (2014), 408–433. Zbl 1355.03026
- [4] D.Yu. Emel'yanov, B.Sh. Kulpeshov, S.V. Sudoplatov, *Algebras of distributions for binary formulas in countably categorical weakly o-minimal structures*, Algebra Logic, **56**:1 (2017), 13–36. Zbl 1420.03082
- [5] K.A. Baikalova, D.Yu. Emel'yanov, B.Sh. Kulpeshov, E.A. Palyutin, S.V. Sudoplatov, *On algebras of distributions of binary isolating formulas for theories of abelian groups and their ordered enrichments*, Russ. Math., **62**:4 (2018), 1–12. Zbl 6955640
- [6] D.Yu. Emel'yanov, B.Sh. Kulpeshov, S.V. Sudoplatov, *Algebras of distributions of binary isolating formulas for quite o-minimal theories*, Algebra Logic, **57**:6 (2019), 429–444. Zbl 1468.03049
- [7] A.B. Altayeva, B.Sh. Kulpeshov, S.V. Sudoplatov, *Algebras of distributions of binary isolating formulas for almost ω -categorical weakly o-minimal theories*, Algebra Logic, **60**:4 (2021), 241–262. Zbl 1515.03172
- [8] B.Sh. Kulpeshov, H.D. Macpherson, *Minimality conditions on circularly ordered structures*, Math. Log. Q., **51**:4 (2005), 377–399. Zbl 1080.03023
- [9] D. Macpherson, C. Steinhorn, *On variants of o-minimality*, Ann. Pure Appl. Logic, **79**:2 (1996), 165–209. Zbl 0858.03039
- [10] B.Sh. Kulpeshov, *On \aleph_0 -categorical weakly circularly minimal structures*, Math. Log. Q., **52**:6 (2006), 555–574. Zbl 1112.03035
- [11] B.Sh. Kulpeshov, *Definable functions in the \aleph_0 -categorical weakly circularly minimal structures*, Sib. Math. J., **50**:2 (2009), 282–301. Zbl 1224.03017
- [12] B.Sh. Kulpeshov, *On indiscernibility of a set in circularly ordered structures*, Sib. Elektron. Mat. Izv., **12** (2015), 255–266. Zbl 1345.03073
- [13] B.Sh. Kulpeshov, V.V. Verbovskiy, *On weakly circularly minimal groups*, Math. Log. Q., **61**:1-2 (2015), 82–90. Zbl 1360.03077

- [14] B.Sh. Kulpeshov, A.B. Altayeva, *Binary formulas in countably categorical weakly circularly minimal structures*, Algebra Logic, **55**:3 (2016), 226–241. Zbl 1404.03035
- [15] B.Sh. Kulpeshov, *On almost binarity in weakly circularly minimal structures*, Eurasian Math. J., **7**:2 (2016), 38–49. Zbl 1463.03011
- [16] A.B. Altayeva, B.Sh. Kulpeshov, *Almost binarity of countably categorical weakly circularly minimal structures*, Math. Notes, **110**:6 (2021), 813–829. Zbl 7455561
- [17] B.Sh. Kulpeshov, *Weakly o-minimal structures and some of their properties*, J. Symb. Log., **63**:4 (1998), 1511–1528. Zbl 0926.03041
- [18] B.Sh. Kulpeshov, *A criterion for binarity of almost ω -categorical weakly o-minimal theories*, Sib. Math. J., **62**:6 (2021), 1063–1075. Zbl 7436440
- [19] D.Yu. Emel'yanov, B.Sh. Kulpeshov, S.V. Sudoplatov, *Algebras of binary formulas for compositions of theories*, Algebra Logic, **59**:4 (2020), 295–312. Zbl 1484.03053
- [20] B.Sh. Kulpeshov, S.V. Sudoplatov, *Algebras of binary formulas for weakly circularly minimal theories with non-trivial definable closure*, Lobachevskii J. Math., **43**:12 (2022), 3532–3540. Zbl 7676592

Beibut Shaiykovich Kulpeshov
Institute of Mathematics and Mathematical Modeling,
Shevchenko street, 28,
050010, Almaty, Kazakhstan
E-mail: kulpesh@mail.ru
Kazakh British Technical University,
Tole bi street, 59,
050000, Almaty, Kazakhstan
E-mail: b.kulpeshov@kbtu.kz
Novosibirsk State Technical University,
K. Marx avenue, 20,
630073, Novosibirsk, Russia
E-mail: kulpeshov@corp.nstu.ru